

Residual-controlled depth extension of finite-horizon Weil positivity

A spatial continuation past $\log 7$ for the shifted-zeta transfer

Edward Baker

Working draft, version 0.3 — 10 September 2026

Abstract

The finite-horizon Weil programme certifies lower bounds for the central Weil form $Q_{0,L}$ on test functions supported in an interval of total length L , and converts them into small-shift contraction of Suzuki’s shifted-zeta transfer. Two-sided certificates exist through $L = \log 7$. This paper extends the certified depth past $\log 7$ by a spatial method: the interval is split into an old part and a short new slab, a finite-rank continuation J from old to new inputs is chosen by a Galerkin principle, and the graph congruence turns $Q_{0,L}$ into a form whose Schur complement is the continuation energy H_J minus a stationarity residual. Because the continuation matches the join, the endpoint logarithm of the new-slab output cancels and the leakage of the transformed form into the infinite polynomial complements becomes small enough to certify. At $L_q = \log 7 + \frac{1}{4} \log(8/7)$ we prove, in the working normalization and with ball arithmetic, that $Q_{0,L_q} \succeq 2.99 \times 10^{-29} I$ while a ball Rayleigh quotient gives $\lambda_{\min}(Q_{0,L_q}) \leq 3.29 \times 10^{-29}$; that the residual satisfies $R^* F^{-1} R \preceq 0.78 H_J$ on the entire old form domain; and that $\|V_{\omega,L}\| \leq e^{-1.4 \times 10^{-29} \omega}$ for $0 < \omega \leq 9 \times 10^{-16}$ and $L \leq L_q$. A second quarter-step to $L_2 = \frac{1}{2} \log 56$ with 32-mode slabs certifies only the comparison $H_{J_2} \succeq 5.01 \times 10^{-8} A_2$ and the upper bound $\lambda_{\min}(Q_{0,L_2}) \leq 1.89 \times 10^{-30}$; we show that the obstruction is entirely the analytic floor of the three-interval complement, and a rebuild with 96-mode slabs then certifies $1.69 \times 10^{-30} \leq \lambda_{\min}(Q_{0,L_2}) \leq 1.89 \times 10^{-30}$, a residual factor 0.8, and contraction for $\omega \leq 2 \times 10^{-16}$. Appending slabs indefinitely degrades the complement floor, so a recursion must periodically merge its intervals. A conditional theorem states what per-step data give divergent depth with vanishing shift; an estimate of the verification dimension required by the polynomial-complement method places its practical horizon near $L = 3$; and a Fourier-side computation shows that the near-null vector of Q_{0,L_q} is a band-limited function tuned to vanish at the low zeros, so that the certificate constrains zeros only below height about 70. Normalization verification, independent review, and any all-depth argument remain open.

Status and dependencies. This is a working draft. Analytic conclusions use the definitions and finite-horizon framework of the Weil-depth manuscript [1]; the exact v0.3 source used here is preserved in this paper’s archive, and the v0.4 revision of [1] changes no constant used below. The author’s independent normalization verification is pending. “Certified” means a recorded guarded Arb sign computation together with its stated analytic reduction, within these working conventions; it does not mean an independent proof audit, formal verification, or any statement about the Riemann hypothesis. Version 0.2 of this draft, which consolidated the original research packets, is preserved unchanged under `archive/drafts/v0.2_2026-09-10/`; the review that motivated the present version is under `archive/reviews/`.

Contents

1	Introduction	4
1.1	The finite-horizon programme and the depth problem	4
1.2	Main results	4
1.3	The mechanism, in one paragraph	5
1.4	Limits of the method	6
1.5	What this paper does not show	6
1.6	Organization	6
2	Conventions and the finite-horizon framework	6
3	Spatial splitting, continuation and the direct floor	7
3.1	The cross block and the new-slab floor	7
3.2	The residual identity	8
3.3	The graph-scaled form and the direct floor	8
3.4	The endpoint logarithm	9
4	Finite certification	9
4.1	Analytic complement floors	9
4.2	The three guarded tests	10
4.3	Sufficient, not necessary	11
5	The first quarter-step	11
5.1	Verification space, continuation and archives	11
5.2	Certified inequalities	11
5.3	Sharpness, and the ladder	12
5.4	Positive-shift consequences	12
6	The second quarter-step	12
6.1	The three-interval construction	12
6.2	The 32-mode build: what is and is not certified	13
6.3	Diagnosis	13
6.4	The 96-mode build closes the step	13
6.5	Appended slabs degrade the floor	14
6.6	Beyond the radius-three representation	14
7	Towards an all-depth argument	15
7.1	The conditional theorem	15
7.2	Why the scalar comparison was dropped	15
7.3	Two invariants for a recursion	16
7.4	The verification dimension	16
7.5	What finite-horizon positivity at fixed L can see	16
7.6	Open problems	17
8	Records and reproducibility	17

A	Generator geometry and cumulative storage	19
A.1	The hyperbolic identity and local extension	19
A.2	Cumulative storage	19
A.3	Why the whole step is difficult	20
B	Full-output storage construction	20
C	A compact map of the logical dependencies	20

1 Introduction

1.1 The finite-horizon programme and the depth problem

Let $Q_{0,L}$ be the central Weil form on $L^2(0, L)$ in the normalization of [1], recalled in Section 2, and let $V_{\omega,L}$ be the finite-horizon realization of Suzuki's shifted-zeta transfer at shift ω . The evolution identity $\frac{d}{ds}\|V_{s,L}f\|^2 = -2Q_{s,L}[V_{s,L}f]$ together with the perturbation bound $\|Q_{s,L} - Q_{0,L}\| \leq C_L s^2$ converts a floor $Q_{0,L} \succeq mI$ into a contraction $\|V_{\omega,L}\| \leq e^{-m\omega/2}$ for $\omega \leq \sqrt{m/(2C_L)}$. A final diagonal argument in that framework needs horizons $L_j \rightarrow \infty$ and shifts $\omega_j \downarrow 0$ with contraction at every (L_j, ω_j) . The Weil-depth paper supplies two-sided enclosures of $\lambda_{\min}(Q_{0,L})$ at seven horizons through $\log 7$, where

$$1.37 \times 10^{-28} \leq \lambda_{\min}(Q_{0,\log 7}) \leq 6.81 \times 10^{-28}.$$

Its method controls the infinite polynomial complement of a global Legendre verification space by an analytic tail floor and a full-output leakage Gram; a positive finite compression alone would not suffice.

The depth problem is to continue past $\log 7$. The naive route, an absolute comparison of the cross block with the smallest old energy, is hopeless at these scales: the old energy is of order 10^{-28} and the elementary local extension estimate of Appendix A can require an increment of length $e^{-10^{31}}$. The route taken here is spatial. Write

$$a = \log 7, \quad h = \frac{1}{4} \log(8/7), \quad L_q = a + h = \frac{3}{4} \log 14 = 1.97929299721144396089 \dots, \quad (1.1)$$

and split Q_{0,L_q} at $x = a$ into the block form $\begin{pmatrix} A & B^* \\ B & F \end{pmatrix}$ with $A = Q_{0,a}$ the old form, $F = Q_{0,h}^\gamma$ the pure gamma form on the new slab (no arithmetic delay fits inside a slab of length $h < \log 2$), and B the complete cross block. A bounded continuation J from old inputs to new inputs gives the *continuation energy* and *residual*

$$H_J = A + B^*J + J^*B + J^*FJ, \quad R = B + FJ, \quad (1.2)$$

and the Schur complement of the split obeys the exact identity $A - B^*F^{-1}B = H_J - R^*F^{-1}R$. Choosing J as the Galerkin minimizer of the energy over 32 new Legendre modes makes the residual small, and the whole problem reduces to certifying inequalities between H_J , R and F on the entire old form domain, including the infinite-dimensional complement of the finite verification space.

1.2 Main results

Fix the verification space of $N = 256$ Legendre modes on $(0, a)$ and $M = 32$ on (a, L_q) , the exact rational continuation $J = J_{128,32}P_{128}$ of record R5, and the graph map $T = \begin{pmatrix} I & 0 \\ J & I \end{pmatrix}$ with $\tau = \frac{1}{2}(q + \sqrt{q^2 + 4})$ for $q \geq \|J\|$. All statements are in the working normalization.

Theorem 1.1 (First quarter-step). *On the entire old form domain, H_J is coercive and*

$$R^*F^{-1}R \preceq 0.78 H_J, \quad A - B^*F^{-1}B \succeq (1 - 0.78)H_J, \quad H_J \succeq 1.13 \times 10^{-7} A. \quad (1.3)$$

The central form satisfies the two-sided enclosure

$$2.99 \times 10^{-29} \leq \lambda_{\min}(Q_{0,L_q}) \leq 3.29 \times 10^{-29}, \quad (1.4)$$

so $Q_{0,L} \succeq 2.99 \times 10^{-29} I$ for every $0 < L \leq L_q$. Consequently, for $0 < L \leq L_q$ and $0 < \omega \leq 9 \times 10^{-16}$,

$$\|V_{\omega,L}\| \leq e^{-1.4 \times 10^{-29} \omega}, \quad D_{\omega,L} = I - V_{\omega,L}^* V_{\omega,L} \succeq (1 - e^{-2.9 \times 10^{-29} \omega})I. \quad (1.5)$$

The lower bound in (1.4) is certified twice: directly, by the absolute Schur test of Section 4, and through the graph-transformed form, where the relative test with an inserted floor certifies $M_\theta \succeq 2.90 \times 10^{-29} I$ at $\theta = 0.9$ and Lemma 3.2 gives $Q_{0,L_q} \succeq 2.90 \times 10^{-29} \tau^{-2} I$. The upper bound is a ball Rayleigh quotient of an explicit rational vector. The enclosure is tight to within ten percent, and the certified floor at L_q exceeds the earlier global certificate at the rational ceiling 1.98 (10^{-31} , record R6) by more than two orders of magnitude.

Theorem 1.2 (Second quarter-step). *Let $L_2 = L_q + h = \frac{1}{2} \log 56$ and split Q_{0,L_2} at $x = L_q$ with old form $A_2 = Q_{0,L_q}$, new form $F_2 = Q_{0,h}^\gamma$, cross block B_2 . With the three-interval verification space of $256 + 96 + 96$ Legendre modes and its exact rational Galerkin continuation J_2 (record R18), on the entire old form domain*

$$1.69 \times 10^{-30} \leq \lambda_{\min}(Q_{0,L_2}) \leq 1.89 \times 10^{-30}, \quad R_2^* F_2^{-1} R_2 \preceq 0.8 H_{J_2}, \quad H_{J_2} \succeq 5.25 \times 10^{-8} A_2, \quad (1.6)$$

and for $0 < L \leq L_2$, $0 < \omega \leq 2 \times 10^{-16}$, $\|V_{\omega,L}\| \leq e^{-8 \times 10^{-31} \omega}$ and $D_{\omega,L} \succeq (1 - e^{-1.6 \times 10^{-30} \omega}) I$. With the original $256 + 32 + 32$ space (records R11, R15) only $H_{J_2} \succeq 5.01 \times 10^{-8} A_2$ and the upper bound are certified; its full-operator and residual tests fail.

Section 6 explains the failure of the 32-mode build and how the 96-mode build was chosen: with every matrix, continuation and error budget unchanged and only the analytic complement floor replaced by a hypothetical value, the 32-mode tests pass once that floor reaches 0.55, whereas the certified three-interval floors are 0.411 (absolute) and 0.293 (residual). The shortfall is an artifact of the analytic estimate for three intervals with two adjacent Carleman couplings, not of the operator or of the verification dimension; Table 1 projected 0.768 and 0.665 for 96-mode slabs, and the build realized exactly these floors and passed.

Theorem 1.3 (Conditional continuation). *Suppose that at every step j of a schedule $L_{j+1} = L_j + h_j$ with $\sum_j h_j = \infty$ a bounded continuation J_j and a residual factor $\theta_j < 1$ are given for which the graph-scaled form M_{θ_j} of Section 3 satisfies $M_{\theta_j} \succeq m'_j I$ with $0 < m'_j \leq f_j$, where $F_j \succeq f_j I$. Then $Q_{0,L_{j+1}} \succeq m'_j \tau_j^{-2} I$, and the shift schedule $\omega_j = \min\{\omega_{j-1}/2, \frac{1}{2}, \sqrt{m_{j+1}/(2C_{L_{j+1}})}\}$ gives $\|V_{\omega_j, L_{j+1}}\| < 1$, $L_j \rightarrow \infty$ and $\omega_j \downarrow 0$.*

The hypotheses of Theorem 1.3 are verified for $j = 0$ and $j = 1$. Version 0.2 of this draft carried the floor through the scalar comparison $H_J \succeq \mu A$; Section 7 explains why that bookkeeping loses about six orders of magnitude per quarter-step against the true decay of the lowest eigenvalue, and why the direct floor in M_θ replaces it.

1.3 The mechanism, in one paragraph

The cross output of an old polynomial ϕ onto the new slab carries an endpoint logarithm at the join whose coefficient is $\frac{1}{2}[\sqrt{h/a} \phi(1) - (J\phi)(0)]$ in unit coordinates. A continuation that matches the join cancels this singularity, and the residual $R = B + FJ$ then has an output whose Legendre coefficients on the slab decay fast. This is why the graph-transformed form leaks much less into the new complement than the untransformed one, why the relative test passes with a verification space on which the absolute comparison of cross norm with old energy would fail by more than four orders of magnitude, and why the 32-mode Galerkin continuation reproduces the lowest eigenvalue of the full 288-mode head to four digits at both steps. Any later variant of the method, with local gamma profiles or exponential modes on the slab, must preserve this property.

1.4 Limits of the method

Three quantitative limits are established below. First, the analytic floor of a partition into a long interval and k appended short slabs decreases roughly linearly in k and becomes negative after a handful of slabs (Table 2), so a recursion that appends a fixed slab per step must periodically merge its intervals into one global verification space or enlarge the slabs. Second, the joint complement floor requires the gamma tail constant $H_N - \gamma - \log(\pi L)$ to exceed the arithmetic norm bound ρ_L , and high-frequency test functions that nearly resonate with every active delay show that ρ_L cannot be much improved, so the verification dimension grows like Le^{ρ_L} : about a thousand modes at $L = 3$ and tens of thousands at $L = 4$ (Table 3). Third, the lowest eigenvalue of the compressed form falls by a factor of about twenty per quarter-step at these depths, from about 6.6×10^{-28} at $\log 7$ to 3.3×10^{-29} at L_q and 1.9×10^{-30} at L_2 ; the working precision needed to resolve it grows only linearly in depth, and is not the barrier.

1.5 What this paper does not show

Nothing here is a statement about zeros of ζ . The certificates are finite computations with explicit error bounds inside a working normalization whose independent audit is pending. The conditional theorem is bookkeeping for data that exist at one step. The cumulative-storage identities of Appendix A, which motivated the project and gave version 0.2 its title, are exact but are not used by any theorem here; every certificate is a zero-shift statement combined with the inherited perturbation bound. Section 7 also records, from a Fourier-side computation, what finite-horizon positivity at a fixed horizon can and cannot see about zeros.

1.6 Organization

Section 2 recalls the framework. Section 3 contains the exact algebra: the residual identity, the graph-scaled form, and the direct-floor lemma. Section 4 describes the finite reductions and their analytic complement bounds. Sections 5 and 6 report the two quarter-steps. Section 7 discusses the all-depth problem, the recursion invariant, and the dimension estimate. Section 8 lists the records. Appendix A collects the generator and cumulative-storage material, Appendix B the full-output storage construction, and Appendix C the dependency map.

2 Conventions and the finite-horizon framework

Let $\mathcal{H}_L = L^2(0, L; \mathbb{C})$, with inner product linear in its second argument. Set $R_L f(x) = f(L - x)$ and $T_d f(x) = \mathbf{1}_{x > d} f(x - d)$. The total support length is L , the centered support is $(-L/2, L/2)$, and the multiplicative cutoff is $e^{L/2}$; a delay at $d = L$ has zero action. All norms and adjoints use unweighted Lebesgue measure.

Write $\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$ and $\mathcal{F}(p) = \xi(\frac{1}{2} + p)$. For $0 < \omega \leq 1/2$ the causal transfer has far-right Laplace symbol $K_\omega(p) = \mathcal{F}(p - \omega)/\mathcal{F}(p + \omega)$ and finite-horizon realization

$$V_{\omega, L} = \left(\sum_{\log n < L} b_\omega(n) T_{\log n} \right) V_{\omega, L}^\gamma, \quad b_\omega(n) = n^{\omega-1/2} \prod_{p|n} (1 - p^{-2\omega}), \quad b_\omega(1) = 1,$$

where V^γ uses the quotient of $\gamma(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)$ and has a locally integrable impulse of order $t^{\omega-1}$ at zero. The transfer contains mixed composite delays; the generator contains only prime powers.

For the centered zero extension f_c use $\widehat{f}_c(\tau) = \int f_c(x) e^{-i\tau x} dx$, and put $C(f) = \int f_c(x) \cosh(x/2) dx$, $S(f) = \int f_c(x) \sinh(x/2) dx$. The central form is

$$Q_{0,L}[f] = \frac{1}{2\pi} \int_{\mathbb{R}} (\operatorname{Re} \psi(\tfrac{1}{4} + \tfrac{i\tau}{2}) - \log \pi) |\widehat{f}_c(\tau)|^2 d\tau + 2|C(f)|^2 - 2|S(f)|^2 - \sum_{\log n < L} \frac{\Lambda(n)}{\sqrt{n}} \langle f, (T_{\log n} + T_{\log n}^*) f \rangle, \quad (2.1)$$

with closed form domain $\mathcal{D}_{\log,L} = \{f : \int \log(2 + |\tau|) |\widehat{f}_c(\tau)|^2 d\tau < \infty\}$, on which polynomials form a core. Smooth bounded multipliers and interval indicators preserve this domain, so a spatial split gives the direct sum of the two local form domains with a bounded cross block; the boundedness of the singular part of the cross block is the Hilbert inequality, with norm at most $\pi/2$. The off-diagonal gamma kernel is

$$q(t) = e^{t/2} - \frac{e^{-5t/2}}{1 - e^{-2t}} = -\frac{G_0(t)}{2t}, \quad G_0(t) = e^{t/2} \frac{t}{\sinh t} - 4t \cosh(t/2) = 1 - \frac{7}{2}t - \frac{1}{24}t^2 - \dots, \quad (2.2)$$

and the arithmetic part at shift s is $-\sum \Lambda(n) n^{-1/2} \cosh(s \log n) (T_{\log n} + T_{\log n}^*)$. The evolution identity and perturbation estimate

$$\frac{d}{ds} \|V_{s,L} f\|^2 = -2Q_{s,L}[V_{s,L} f], \quad \|Q_{s,L} - Q_{0,L}\| \leq C_L s^2, \quad (2.3)$$

with $V_{0,L} = I$ as a strong limit, have the core and limiting interpretation of [1]; the difference of the two forms is bounded although each is unbounded. In the working normalization the explicit formula gives $Q_{0,L}[f] = 2 \sum_{\gamma > 0} |\widehat{f}_c(\gamma)|^2$ over the ordinates of the nontrivial zeros, a fact checked numerically in the review cycle of [1] and used in Section 7 only as a diagnostic.

3 Spatial splitting, continuation and the direct floor

The algebra of this section applies to closed semibounded forms A, F with $F \succeq f_0 I > 0$, a bounded cross block B , and a bounded J into the operator domain of F with FJ bounded; finite-rank polynomial continuations satisfy this, because F applied to a polynomial on the slab is a polynomial plus endpoint logarithms, which is square integrable.

3.1 The cross block and the new-slab floor

At $a = \log 7$ and $0 < t < h$ the complete cross block is

$$(Bf)(t) = \int_0^a q(a+t-y) f(y) dy - \sum_{n \in \{2,3,4,5,7\}} \frac{\Lambda(n)}{\sqrt{n}} f(a+t-\log n), \quad (3.1)$$

in which the $n = 7$ term is exactly $-\log 7 f(t)/\sqrt{7}$. The new-slab form $F = Q_{0,h}^\gamma$ has the ground-state representation, in unit coordinates $v \in (0, 1)$ and with $\ell_h = \gamma + \log(2\pi h)$,

$$F[g] = \int_0^1 (F1)(v) |g(v)|^2 dv + \frac{1}{4} \int_0^1 \int_0^1 \frac{G_0(h|u-v|)}{|u-v|} |g(u) - g(v)|^2 du dv, \\ (F1)(v) = -\ell_h - \frac{1}{2} \log(v(1-v)) - \frac{1}{2} \sum_{j \geq 1} \frac{g_j h^j}{j} \{v^j + (1-v)^j\},$$

where $G_0(t) = 1 + \sum g_j t^j$. Since $G_0 > 0$ on $[0, h]$, $v(1-v) \leq \frac{1}{4}$ and $g_1 = -\frac{7}{2}$, the Cauchy remainder of the profile gives

$$F \succeq f_0 I, \quad f_0 \geq -\ell_h + \log 2 + \frac{7}{4}h - \frac{1}{2} \sum_{j=2}^d \frac{|g_j| h^j}{j} - \frac{128(h/3)^{d+1}}{(d+1)(1-h/3)} > 1.73617237627376657. \quad (3.2)$$

3.2 The residual identity

Lemma 3.1 (Residual identity and transported energy). *With H_J, R as in (1.2), on the old form domain*

$$S := A - B^* F^{-1} B = H_J - R^* F^{-1} R. \quad (3.3)$$

If H_J is coercive and $R^ F^{-1} R \preceq \theta H_J$ for some $0 \leq \theta < 1$, then $S \succeq (1 - \theta)H_J$ and, for all old and new form-domain inputs f, g ,*

$$Q[f, g] \geq (1 - \sqrt{\theta})\{H_J[f] + F[g - Jf]\}.$$

Proof. Expanding $R^* F^{-1} R = (B + FJ)^* F^{-1} (B + FJ)$ and cancelling the J terms gives (3.3). With $g = Jf + v$ the complete form is $H_J[f] + 2\operatorname{Re}\langle v, Rf \rangle + F[v]$; the residual bound gives $|\langle v, Rf \rangle| \leq \sqrt{\theta} H_J[f]^{1/2} F[v]^{1/2}$, and $2ab \leq a^2 + b^2$ finishes. $\square$

3.3 The graph-scaled form and the direct floor

Let $\mathcal{T} = \begin{pmatrix} I & 0 \\ J & I \end{pmatrix}$, so that $\mathcal{T}^* Q \mathcal{T} = \begin{pmatrix} H_J & R^* \\ R & F \end{pmatrix}$, and for $0 < \theta < 1$ put $\lambda = \theta^{-1/2}$ and

$$M_\theta = \begin{pmatrix} H_J & \lambda R^* \\ \lambda R & F \end{pmatrix}. \quad (3.4)$$

Coercivity of M_θ proves $H_J \succ 0$ and $R^* F^{-1} R \preceq \theta H_J$ by its own Schur complement; this is how the residual factor is certified. The following observation, absent from version 0.2, shows that the same test also carries the floor.

Lemma 3.2 (Direct floor). *If $M_\theta \succeq mI$ with $0 < m \leq f_0$, then $\mathcal{T}^* Q \mathcal{T} \succeq mI$ and $Q \succeq m\tau^{-2}I$, where $\tau = \frac{1}{2}(q + \sqrt{q^2 + 4})$ and $q \geq \|J\|$.*

Proof. Restricting $M_\theta \succeq m$ to inputs $(f, 0)$ and $(0, v)$ gives $H_J \succeq m$ and $F \succeq m$. Since $\lambda \geq 1$,

$$\mathcal{T}^* Q \mathcal{T} = \frac{1}{\lambda} M_\theta + \left(1 - \frac{1}{\lambda}\right) \begin{pmatrix} H_J & 0 \\ 0 & F \end{pmatrix} \succeq \left(\frac{m}{\lambda} + \left(1 - \frac{1}{\lambda}\right)m\right)I = mI.$$

The graph map has norm at most τ , the largest singular value of $\begin{pmatrix} 1 & 0 \\ q & 1 \end{pmatrix}$, and $\mathcal{T}$ is invertible, so $Q[u] = (\mathcal{T}^* Q \mathcal{T})[\mathcal{T}^{-1}u] \geq m\|\mathcal{T}^{-1}u\|^2 \geq m\tau^{-2}\|u\|^2$. $\square$

Remark 3.3. Lemma 3.1 loses the factor $1 - \sqrt{\theta}$ and needs a separate floor for H_J ; Lemma 3.2 loses only τ^{-2} , which is 0.392 at the first step, and needs nothing but the finite test that certifies $M_\theta \succeq m$. It also makes the comparison $H_J \succeq \mu A$ unnecessary for carrying a floor between steps; see Section 7.

3.4 The endpoint logarithm

In unit new coordinates the cross output of a unit old polynomial ϕ has the singular part

$$-\frac{\sqrt{r}}{2} \left[\phi(1+rv) \log \frac{v+a/h}{v} - H_\phi(1+rv) \right], \quad r = h/a, \quad H_\phi(z) = \int_0^1 \frac{\phi(z) - \phi(u)}{z-u} du,$$

and F applied to a new polynomial g has the singular part $-\frac{1}{2}g(0) \log v - \frac{1}{2}g(1) \log(1-v)$. The coefficient of $\log v$ in the output of Q on the pair $(\phi, J\phi)$ is therefore $\frac{1}{2}[\sqrt{h/a} \phi(1) - (J\phi)(0)]$. The Galerkin continuation, which minimizes H_J , drives this coefficient toward zero because a mismatched join costs energy of order its square in the logarithmic form. The residual output $R\phi$ is then a polynomial plus logarithms with small coefficients, and its Legendre coefficients on the slab decay fast. Section 5 records the consequences: the leakage of the graph-transformed form into the new complement is certified small, and the lowest eigenvalue of H_J agrees with that of the full head to four digits.

4 Finite certification

Take the verification projection $P = P_o \oplus P_n$ onto N old and M new Legendre modes, $K = I - P$, and write $H = PQP$, $W = KQP$, and $G = (QP)^*(QP)$ for the compressed head, the leakage, and the full-output Gram. Piecewise polynomials lie in the operator domain, so $E := W^*W = G - H^2$. All outputs are integrated on the full intervals; no output is projected. Model matrices $\tilde{H}, \tilde{G}$ carry the analytic profile truncation, and every test subtracts an explicit error budget before its interval LDL factorization.

4.1 Analytic complement floors

Arithmetic. Let $\alpha_n = \Lambda(n)/\sqrt{n}$ and $T = \sum_{\log n < L} \alpha_n (T_{\log n} + T_{\log n}^*)$. If a bounded positive weight w , bounded away from zero, satisfies $Tw \leq \rho w$, then $\|T\| \leq \rho$, by $2|f(x)f(y)| \leq |f(x)|^2 w(y)/w(x) + |f(y)|^2 w(x)/w(y)$ applied to the translated pairings. A 512-cell piecewise constant weight with exact rational values, verified on every open interval cut by the shifted discontinuities (2,152 intervals at L_q), certifies

$$\rho < 1.949177212012646 \quad (L = L_q), \quad \rho_2 < 2.02694013535790 \quad (L = L_2). \quad (4.1)$$

Gamma tails. On an interval of length ℓ , the pure gamma form restricted to the complement of k Legendre modes satisfies

$$Q_{0,\ell}^\gamma|_{P_k^\perp} \succeq g_{\ell,k} I, \quad g_{\ell,k} = H_k - \gamma - \log(\pi\ell) - \frac{K_\ell \ell}{2\sqrt{k(k+1)}}, \quad K_\ell \geq |w_0(0)| + \int_0^\ell |w'_0(t)| dt, \quad (4.2)$$

with $H_k = \sum_{j \leq k} 1/j$ and $w_0(t) = (G_0(t) - 1)/t$. The first three terms are the degree-zero logarithmic fractional-integration bound of [1]; the last is the smooth part, estimated through the primitive of a tail input, which vanishes at both endpoints and obeys $\|\int_0^\ell f\| \leq \ell \|f\|/(2\sqrt{k(k+1)})$. The variation hypothesis on K_ℓ is the correct one: for causal convolution $S_{w_0} = (w_0(0)I + S_{w'_0})I_1$, and Young's inequality then needs $|w_0(0)| + \|w'_0\|_{L^1}$. Version 0.1 stated a supremum hypothesis; the implemented constant $K_\ell = \sum_{j \leq d} |g_j| \ell^{j-1} + \frac{256}{3}(\ell/3)^d/(1 - \ell/3)$ bounds the variation, so no numerical value changed.

Cross couplings. Between adjacent intervals the singular kernel $-1/(2|x-y|)$ has norm at most $\pi/2$ (Hilbert's inequality); between intervals separated by a gap d it has Hilbert–Schmidt norm at most $\frac{1}{2}\sqrt{\log \frac{(d+\ell_i)(d+\ell_j)}{d(d+\ell_i+\ell_j)}}$. The smooth part, by integration by parts on either tail, is at most

$$b_{ij}^{\text{sm}} = \frac{\sqrt{\ell_i \ell_j}}{4} \min \left\{ \frac{\ell_i}{\sqrt{N_i(N_i+1)}}, \frac{\ell_j}{\sqrt{N_j(N_j+1)}} \right\} \sup_{[0,L]} |w'_0|.$$

Joint floor. Form the symmetric comparison matrix with diagonal g_{ℓ_i, N_i} and off-diagonal entries $-(b_{ij}^{\text{sing}} + b_{ij}^{\text{sm}})$, subtract ρI , and bound its least eigenvalue from below; this bounds KQK on the joint complement. For two intervals the bound is explicit,

$$\beta = \frac{1}{2}(g_{a,N} + g_{h,M} - \sqrt{(g_{a,N} - g_{h,M})^2 + 4b_\gamma^2}) - \rho, \quad b_\gamma = \frac{\pi}{2} + b^{\text{sm}}, \quad (4.3)$$

and at $(N, M) = (256, 32)$ the enclosed values are $g_{a,256} = 3.7145\dots$, $g_{h,32} = 5.7345\dots$, $b^{\text{sm}} = 0.00129$, $\beta = 0.90675\dots$. For the graph-scaled form M_θ the old–new cross couplings are multiplied by λ , and the arithmetic cross part, whose old source windows $(a - \log n, a + h - \log n)$ are pairwise disjoint because the gaps between the active $\log n$ exceed $0.22 \gg h$, is bounded by $b_{\text{ar}} = (\sum_n \alpha_n^2)^{1/2} = 1.3497$; the modified floor is

$$\beta_\theta = \frac{1}{2}(g_{a,N} + g_{h,M} - \sqrt{(g_{a,N} - g_{h,M})^2 + 4\lambda^2 b_\gamma^2}) - \rho - (\lambda - 1)b_{\text{ar}}, \quad (4.4)$$

which equals $0.76166\dots$ at $\theta = 0.9$.

4.2 The three guarded tests

Each test eliminates the complement by a Schur argument: if $KQK \succeq \beta I$ and $0 \leq m < \beta$, then $(\beta - m)(H - mI) - E \succ 0$ implies $Q \succeq mI$, because $(KQK - m)^{-1} \preceq (\beta - m)^{-1}I$.

Absolute test. With a model head error η and model output error δ , a valid leakage error is $e = 2\delta\sqrt{\text{tr } \tilde{G}} + \delta^2 + 2\eta\|\tilde{H}\| + \eta^2$, and the guarded test is

$$(\beta - m)(\tilde{H} - mI) - \tilde{E} - [(\beta - m)\eta + e]I \succ 0. \quad (4.5)$$

The profile remainder is $\eta_d(L) = 256(L/3)^{d+1}/((d+1)(1-L/3))$ for $L < 3$; the exterior-log expansion adds its own remainder. Passing at m certifies $Q \succeq mI$; the largest passing three-significant-figure m is found by bisection and recorded.

Relative test. Retain the old and new output Grams G_o, G_n separately, let H_o, H_n be the corresponding row blocks of H , and with the finite matrix T of $\mathcal{T}$ set $E_o = T^*(G_o - H_o^* H_o)T$, $E_n = T^*(G_n - H_n^* H_n)T$, $D_o = \text{diag}(I_N, \lambda I_M)$, $D_n = \text{diag}(\lambda I_N, I_M)$. The leakage Gram of M_θ is exactly

$$E_\theta = D_o E_o D_o + D_n E_n D_n, \quad (4.6)$$

because the old-tail output of M_θ on a head input (f, v) is $K_o \Pi_o Q \mathcal{T}(f, \lambda v)$, using $K_o J^* = 0$, while the new-tail output is $K_n \Pi_n Q \mathcal{T}(\lambda f, v)$. Let H_θ be $T^* H T$ with its spatial off-diagonal blocks scaled by λ . The guarded test with an inserted floor $m < \beta_\theta$ is

$$(\beta_\theta - m)(H_\theta - mI) - E_\theta - [(\beta_\theta - m) 2\lambda\eta\|T\|_F^2 + e(\|TD_o\|_F^2 + \|TD_n\|_F^2)]I \succ 0; \quad (4.7)$$

passing certifies $M_\theta \succeq mI$ on the whole space, hence the residual factor θ , the coercivity $H_J \succeq m$, and by Lemma 3.2 the floor $Q \succeq m\tau^{-2}I$. The head-scaling error uses $\|X_\theta\| \leq \lambda\|X\|$; the factor 2 is spare.

Comparison test. For $0 < \mu < 1$ put $M_\mu = H_J - \mu A$. Its old-head compression is $T_o^* H T_o - \mu A_N$ with $T_o = \begin{pmatrix} I_N \\ J \end{pmatrix}$; since $JK_o = 0$ its old-tail leakage Gram is exactly $U_\mu^* E_o U_\mu$ with $U_\mu = \begin{pmatrix} (1-\mu)I_N \\ J \end{pmatrix}$ and $E_o = G_o - H_o^* H_o$; and its tail compression is $(1-\mu)K_o A K_o \succeq (1-\mu)\beta_A I$ with $\beta_A = g_{a,N} - \rho$. The guarded test

$$(1-\mu)\beta_A(T_o^* H T_o - \mu A_N) - U_\mu^* E_o U_\mu - [(1-\mu)\beta_A\eta(\|T_o\|_F^2 + \mu) + e\|U_\mu\|_F^2]I \succ 0 \quad (4.8)$$

proves $H_J \succeq \mu A$ on the full old form domain.

Upper bounds. For a rational vector v in the head space, $\lambda_{\min}(Q) \leq (v^* \tilde{H} v) / \|v\|^2 + \eta$; the vector is the rounded lowest eigenvector of $\tilde{H}$ and the quotient is evaluated in ball arithmetic. This is the same device as in [1].

4.3 Sufficient, not necessary

A failed guarded test records a failure to certify, never operator negativity: the complement floors are bounds, the leakage is charged at its full norm, and the pivot at which an LDL fails is not a localization of anything. Conversely, a passing test with an inserted floor m is exactly as strong as m ; the recorded validators of version 0.2 inserted 10^{-33} and 10^{-36} and thereby understated what the matrices support by more than four orders of magnitude, which is why every floor below is the largest three-figure value passing a bisection.

5 The first quarter-step

5.1 Verification space, continuation and archives

The verification space has $N = 256$ old and $M = 32$ new modes, profile degree 320, exterior-log degree 100, and the matrices were built in 6144-bit ball arithmetic (record R7; about 21 minutes). Its full-output error is below 4.953×10^{-58} and its leakage error below 1.024×10^{-55} . The continuation is the Galerkin minimizer $-F_{32}^{-1} B_{32}$ on the first 128 old modes, rounded to exact decimal rationals (record R5), extended by zero to the remaining old modes; the rounding error is part of the measured residual. $\|J\|_F = 0.97098$, so $\tau^2 \leq 2.5508$.

5.2 Certified inequalities

Proposition 5.1. *In the working normalization, with the data above:*

- (i) (R14, absolute) the test (4.5) passes at $m = 2.99 \times 10^{-29}$ and fails at $m = 3.00 \times 10^{-29}$; hence $Q_{0,L_q} \succeq 2.99 \times 10^{-29} I$.
- (ii) (R14, relative) the test (4.7) passes at $\theta = 0.9$ with $m = 2.90 \times 10^{-29}$, at $\theta = 0.8$ with $m = 2.67 \times 10^{-29}$, and with $m = 0$ at $\theta = 0.78$ but not at $\theta = 0.76$; hence $R^* F^{-1} R \preceq 0.78 H_J$, $H_J \succeq 2.90 \times 10^{-29} I$, and, by Lemma 3.2, $Q_{0,L_q} \succeq 2.90 \times 10^{-29} \tau^{-2} I$.
- (iii) (R14, comparison) the test (4.8) passes at $\mu = 1.13 \times 10^{-7}$ and fails at the next three-figure value; hence $H_J \succeq 1.13 \times 10^{-7} A$ and, with (ii), $A - B^* F^{-1} B \succeq (1 - 0.78) 1.13 \times 10^{-7} A$.
- (iv) (R16, upper bound) the ball Rayleigh quotient of the rounded lowest eigenvector gives $\lambda_{\min}(Q_{0,L_q}) \leq 3.29 \times 10^{-29}$.

Items (i)–(iii) are re-evaluations at 2048 bits of the stored 6144-bit balls through the archive-based replay `replay_floor.py`, which verifies both archive hashes and the builder sources before use; the

earlier records R7, R8 and R10 with their fixed thresholds are reproduced by it to every printed digit. The compressed lowest eigenvalue of the 288-mode head is 3.28×10^{-29} (midpoint diagnostic), so the certified floor captures more than ninety percent of what the verification space can show.

5.3 Sharpness, and the ladder

The two-sided enclosure (1.4) continues the enclosures of [1]. Where the lowest eigenvalue of the 128-mode compression at $\log 7$ is 6.81×10^{-28} and its certified floor 1.37×10^{-28} , the 256-mode compression gives 6.56×10^{-28} , the 288-mode compression at L_q gives 3.28×10^{-29} , and the 320- and 448-mode compressions at L_2 both give 1.88×10^{-30} (certified two-sided in Section 6.4): a decrease by a factor of about twenty per quarter-step of length 0.0334, that is, a local rate near e^{-88} per unit depth. Two facts follow. The shift intervals that (2.3) yields shrink accordingly, so any depth schedule must accept rapidly vanishing shifts, which is what the diagonal argument requires anyway; and the working precision needed to resolve floors grows only linearly in depth, about forty digits per unit of L , which 6144-bit arithmetic supplies far beyond $L = 3$.

5.4 Positive-shift consequences

From $Q_{0,L_q} \succeq 2.99 \times 10^{-29} I$ and compression, the same floor holds at every $L \leq L_q$. The perturbation constant of (2.3) evaluated at L_q is $C_{L_q} < 16.536113848659$, so

$$C_{L_q}(9 \times 10^{-16})^2 < 1.34 \times 10^{-29} < \frac{1}{2} 2.99 \times 10^{-29},$$

and $Q_{s,L} \succeq 1.49 \times 10^{-29} I$ on the rectangle $0 \leq s \leq 9 \times 10^{-16}$, $L \leq L_q$. Integrating (2.3) gives (1.5). This replaces the interval $\omega \leq 5 \times 10^{-18}$ of version 0.2, which was derived from the inserted floor 10^{-33} , and it exceeds the interval $\omega \leq 5 \times 10^{-17}$ of the global certificate at 1.98 (R6, floor 10^{-31}). Together with the interval through 4×10^{-15} at $\log 7$ from [1], the certified region now covers the staircase $(\log 7, 4 \times 10^{-15}) \rightarrow (\log 7, 9 \times 10^{-16}) \rightarrow (L_q, 9 \times 10^{-16})$ with the spatial certificate alone. If $D_{\omega,L} \succeq dI$ then the cumulative relative coupling of Appendix A satisfies $c_D(\omega)^2 \leq 1 - d$, so $c_D(\omega)^2 \leq e^{-2.9 \times 10^{-29} \omega} < 1$ throughout the shift interval.

6 The second quarter-step

6.1 The three-interval construction

The construction keeps the first spatial head and appends a slab of length h , so the partition is $(\ell_0, \ell_1, \ell_2) = (\log 7, h, h)$ with $L_2 = \frac{1}{2} \log 56 = 2.01267584536757 \dots$ and retained dimensions $(256, 32, 32)$; the old verification space is the 288-dimensional first-step space, not a new global polynomial space. On each output interval the truncated gamma output of every retained input is $p(v) + \sum_c p_c(v) \log |v - c|$ with all logarithmic centres outside the open interval, arithmetic outputs are polynomials on exact translated windows whose endpoints are rational combinations of prime logarithms compared symbolically, and products are integrated without projection through polynomial-logarithm moments (dilogarithms for two distinct centres, $\frac{1}{2} \log^2$ for coincident ones). The first-step head and old/old Gram blocks are inherited from the hash-verified R7 archive; the mismatch between the inherited exterior-log series and the new exact-log representation is charged to the Gram error. The new archive (record R11, about 764 seconds, 297 MB) has $\eta < 5.494 \times 10^{-56}$, Gram error below 2.376×10^{-53} and leakage error below 3.202×10^{-53} . Its continuation J_2 is the Galerkin minimizer on all 288 old modes, rounded to 80 decimal digits, with $\|J_2\|_F = 1.0479$.

The three-interval complement floor is the least eigenvalue of the 3×3 comparison matrix with diagonal $(g_{a,256}, g_{h,32}, g_{h,32}) = (3.7145, 5.7345, 5.7345)$, adjacent couplings $\pi/2 + b^{\text{sm}}$, the separated coupling $0.4112 + b^{\text{sm}}$, minus $\rho_2 = 2.0269$: it equals $0.41090\dots$, with Perron-type eigenvector $(1, 0.69, 0.45)$. Its $\theta = 0.9$ modification is $0.29312\dots$. The old two-interval floor used by the comparison test is $0.82993\dots$

6.2 The 32-mode build: what is and is not certified

Proposition 6.1. *In the working normalization: (i) (R11, R15) the comparison test passes at $\mu = 5.01 \times 10^{-8}$ and fails at the next three-figure value, so $H_{J_2} \succeq 5.01 \times 10^{-8} A_2$ on the entire old form domain; (ii) (R16) $\lambda_{\min}(Q_{0,L_2}) \leq 1.89 \times 10^{-30}$; (iii) (R11, R12) the absolute test fails at $m = 10^{-36}$ (pivot 182), the residual test fails at $\theta = 0.9, 0.95, 0.99$ (pivots 55, 155, 174), and twenty interval-weighted variants of the residual test at $\theta = 0.99$ and 0.9999 fail.*

The failures in (iii) are reproduced at 2048 and 4096 bits, where the second run re-evaluates the same stored balls and confirms that no working-precision effect is involved.

6.3 Diagnosis

Replace the certified complement floor by a hypothetical value and leave everything else unchanged: the head, the three output Grams, the continuation, and the error budgets. This is a diagnostic of the finite matrices, recorded as such (R16), and certifies nothing.

test at L_2	certified floor	hypothetical floor	outcome
absolute, $m = 10^{-36}$	0.4109	0.50 / 0.55	fails / passes
absolute, bisected m	—	0.65	largest passing 1.64×10^{-30}
residual, $\theta = 0.9$	0.2931	0.50 / 0.55	fails / passes
residual, $\theta = 0.8$	—	0.65	passes

So the second-step operator and its 320-mode verification space would certify $\theta = 0.8$ and a floor within fifteen percent of the rigorous upper bound 1.89×10^{-30} if the analytic complement floor were 0.65, and both tests already pass at 0.55; the certified floors are 0.41 and 0.29. The shortfall has a transparent source. In the comparison matrix the long interval carries the small diagonal $g_{a,256} - \rho_2 = 1.69$, both adjacent couplings are the Carleman constant $\pi/2$, and the middle slab is coupled on both sides. Raising the slab dimension raises $g_{h,M}$ like H_M ; raising the old dimension raises $g_{a,N}$ like H_N . Table 1 evaluates the comparison matrix for candidate verification spaces; ninety-six modes on each slab supply both floors with margin, and Section 6.4 reports that build: floor 1.69×10^{-30} , $\theta = 0.8$, shift interval 2×10^{-16} .

Two refinements do not help. Separate tail floors per interval (record R12) redistribute the Perron floor rather than increase it: with the Perron weights all three floors equal the joint floor, and every other weight lowers one of them. Replacing the uniform ρ_2 by blockwise arithmetic norms (the first-step ρ on the long interval, b_{ar} across, zero on the slabs) is worse, because the triangle inequality across blocks loses what the global weight captures; the resulting floor is negative.

6.4 The 96-mode build closes the step

The first step was rebuilt with 96 modes on the slab (record R18; the same builder as R7 with **-new 96**, reusing the old-depth cache; 41 minutes at 6144 bits in the recorded environment), giving a two-interval complement floor of 1.109 and passing its absolute test. The second step was then built

Table 1: Projected complement floors at L_2 for candidate verification spaces (constants from the records; the uniform ρ_2 is subtracted). The tests need about 0.55. The 256 + 96 + 96 build (Section 6.4) realized 0.768 and 0.665.

old + slab + slab modes	joint floor	$\theta = 0.9$ floor
256 + 32 + 32 (recorded)	0.411	0.293
256 + 48 + 48	0.562	0.450
256 + 64 + 64	0.654	0.547
256 + 96 + 96	0.768	0.665
256 + 128 + 128	0.837	0.737
384 + 32 + 32	0.637	0.513
384 + 64 + 64	0.924	0.810

from this seed with 96 modes on the new slab as well (85 minutes; the 633 MB archive is external). Its three-interval complement floor is $0.76828\dots$ and the $\theta = 0.9$ modification $0.6654\dots$, matching Table 1 to the displayed digits; $\eta < 5.494 \times 10^{-56}$, Gram error below 3.236×10^{-53} , leakage error below 4.447×10^{-53} . The Galerkin continuation on the 352 old modes has $\|J_2\|_F = 1.0503$, so $\tau^2 \leq 2.7379$; its rounded record is about 3 MB and is kept beside the archive.

Proposition 6.2. *In the working normalization, with the 256 + 96 + 96 data: (i) the absolute test passes at $m = 1.69 \times 10^{-30}$ and fails at 1.70×10^{-30} ; (ii) the relative test passes at $\theta = 0.9$ with $m = 1.65 \times 10^{-30}$ (fails at 1.66×10^{-30}) and at $\theta = 0.8$ with $m = 0$; (iii) the comparison passes at $\mu = 5.25 \times 10^{-8}$ and fails at 5.26×10^{-8} ; (iv) the ball Rayleigh quotient of the rounded lowest eigenvector gives $\lambda_{\min}(Q_{0,L_2}) \leq 1.89 \times 10^{-30}$ (quotient 1.884216×10^{-30} , against 1.884228×10^{-30} from the 320-mode head).*

Hence Theorem 1.2: the certified floor is within twelve percent of the upper bound, $H_{J_2} \succeq 1.65 \times 10^{-30} I$, and by Lemma 3.2 the graph route alone gives $Q_{0,L_2} \succeq 1.65 \times 10^{-30} \tau^{-2} I \succeq 6.0 \times 10^{-31} I$. The perturbation constant at L_2 is $C_{L_2} < 17.4886$, so $C_{L_2} (2 \times 10^{-16})^2 < 7.0 \times 10^{-31} < \frac{1}{2} 1.69 \times 10^{-30}$, which gives the shift statements. The lowest eigenvalue of the 448-mode head agrees with that of the 320-mode head to five digits, so the 32-mode slabs had already resolved the near-null vector; the additional modes were needed only for the analytic complement floor, exactly as the diagnosis said.

6.5 Appended slabs degrade the floor

The same comparison matrix, with k slabs of M modes appended behind the 256-mode head, has the least eigenvalues of Table 2. Each appended slab adds an adjacent $\pi/2$ coupling to a path whose head diagonal stays at $g_{a,256} - \rho$, and the floor decreases roughly linearly in k . A recursion that appends one slab per step therefore cannot continue indefinitely with a fixed slab dimension; it must periodically merge the old intervals into a single interval with a fresh global basis, or grow the slab dimension with every step. Section 7 takes this up.

6.6 Beyond the radius-three representation

The current driver expands the gamma profile about zero and therefore enforces $L < 3$; this is a representation restriction, not a theorem. The exact identity $q(t) = 2 \cosh(t/2) - \sum_{k \geq 0} e^{-(1/2+2k)t}$ suggests a local representation: on any interval $e^{x/2}, e^{-x/2}$ span the finite-rank hyperbolic term, and for intervals with gap $d > 0$ the remainder after K exponential terms has norm at most $\sqrt{\ell_i \ell_j} e^{-(1/2+2K)d} / (1 - e^{-2d})$, so local Taylor profiles need only cover each diagonal and adjacent

Table 2: Least eigenvalue of the comparison matrix for a 256-mode head followed by k slabs of M modes each, with ρ_2 subtracted.

M	$k = 2$	$k = 3$	$k = 5$	$k = 9$
32	0.41	0.13	-0.26	-0.74
64	0.65	0.45	0.16	-0.23
128	0.84	0.69	0.48	0.19

pair of total length below three. This has not been implemented; it removes the $L < 3$ restriction and nothing else.

7 Towards an all-depth argument

7.1 The conditional theorem

Theorem 7.1 (Conditional continuation scheme). *Let $L_{j+1} = L_j + h_j$ with $h_j > 0$ and $\sum_j h_j = \infty$, and $Q_{0,L_0} \succeq m_0 I > 0$. Suppose that at every step the spatial form has the closed block realization $Q_{0,L_{j+1}} = \begin{pmatrix} A_j & B_j^* \\ B_j & F_j \end{pmatrix}$ with $A_j = Q_{0,L_j}$, $F_j \succeq f_j I > 0$ and bounded cross block; that J_j is bounded into the operator domain of F_j with $F_j J_j$ bounded; and that for some $\theta_j < 1$ the graph-scaled form M_{θ_j} built from $H_j = A_j + B_j^* J_j + J_j^* B_j + J_j^* F_j J_j$ and $R_j = B_j + F_j J_j$ satisfies $M_{\theta_j} \succeq m'_j I$ with $0 < m'_j \leq f_j$. Put $q_j \geq \|J_j\|$, $\tau_j = \frac{1}{2}(q_j + \sqrt{q_j^2 + 4})$ and $m_{j+1} = m'_j \tau_j^{-2}$. Then $Q_{0,L_j} \succeq m_j I > 0$ for every j . If moreover $\|Q_{s,L_j} - Q_{0,L_j}\| \leq C_{L_j} s^2$ for $0 \leq s \leq \frac{1}{2}$, choose $0 < \omega_0 \leq \min\{\frac{1}{2}, \sqrt{m_0/(2C_{L_0})}\}$ and $\omega_j = \min\{\omega_{j-1}/2, \frac{1}{2}, \sqrt{m_j/(2C_{L_j})}\}$; then $\omega_j \downarrow 0$, $L_j \rightarrow \infty$, and $\|V_{\omega_j, L_j}\| \leq e^{-m_j \omega_j/2} < 1$.*

Proof. Lemma 3.2 gives the floor at each step. For $0 \leq s \leq \omega_j$ the perturbation bound gives $Q_{s,L_j} \succeq m_j I/2$, and integrating (2.3) with the strong zero-shift limit gives the contraction. The halving condition forces $\omega_j \rightarrow 0$ and the divergent increments force $L_j \rightarrow \infty$. $\square$

The hypotheses hold for $j = 0$ with $(\theta_0, m'_0) = (0.9, 2.90 \times 10^{-29})$ or $(0.8, 2.67 \times 10^{-29})$, and for $j = 1$ with $(\theta_1, m'_1) = (0.9, 1.65 \times 10^{-30})$ on the 96-mode space. The theorem is bookkeeping; the mathematics is entirely in producing $M_{\theta_j} \succeq m'_j$ indefinitely, and in preventing a schedule from accumulating at finite depth.

7.2 Why the scalar comparison was dropped

Version 0.2 carried the floor through $H_j \succeq \mu_j A_j$ and the recurrence $m_{j+1} = (1 - \sqrt{\theta_j}) \min\{\mu_j m_j, f_j\} \tau_j^{-2}$. The comparison threshold is sharp—the recorded validators fail at the next three-figure value above 1.13×10^{-7} and 5.01×10^{-8} , and the generalized eigenvalue $\min H_j/A$ on the verification space is 1.22×10^{-7} at the first step and 6.0×10^{-8} at the second—but it measures the worst direction of H_j/A , which is not the weak direction of A . The lowest eigenvalue actually falls by a factor of about twenty per quarter-step, while the recurrence charges about 2×10^{-9} per step. After three steps the scalar floor would lie twenty-five orders of magnitude below the truth and the shift schedule would be useless. The comparison remains a clean statement about the retained fraction of old energy, $A - B^* F^{-1} B \succeq (1 - \theta)\mu A$, and a sensitive diagnostic of the continuation; it should not carry the floor.

Table 3: Dimension required for a positive complement floor, $N \approx \pi L e^\gamma e^{\rho_L^+}$, and the dimension at which the leakage tests are expected to pass (scaling the factor of about four observed at L_q , where 128 modes failed and 256 passed).

L	e^L	ρ_L^+	N for a positive floor	N for the tests
1.98	7.2	1.70	61	256 (observed)
2.5	12.2	2.80	230	about 10^3
3.0	20.1	4.09	10^3	$3-4 \times 10^3$
3.5	33.1	5.65	5.5×10^3	$1.5-2 \times 10^4$
4.0	54.6	7.54	4.2×10^4	above 10^5

7.3 Two invariants for a recursion

Table 2 rules out appending fixed slabs indefinitely. The two-interval invariant treats $(0, L_j)$ at every step as one interval with a fresh global Legendre basis of N_j modes, so the complement floor is the first-step value as long as $g_{L_j, N_j} - \rho_{L_j}$ stays near 1.8; its cost is the old/old Gram at every step, which scales like N_j^2 in 6144-bit moment products (21 minutes at $N = 256$). A hybrid appends slabs while the comparison floor stays above what the tests need, about 0.55, and then merges; with 96-mode slabs this happens every three or four quarter-steps. In either case the state carried between steps should be the head, the full-output Grams, the exact rational continuation, the certified m'_j , and the compressed lowest eigenvalue as the sharpness diagnostic.

7.4 The verification dimension

The joint floor requires, roughly, $H_N - \gamma - \log(\pi L) > \rho_L$. The arithmetic bound cannot be much improved on the complement: functions $\phi(x) \cos(\omega x)$ with $\omega \log n$ near $2\pi\mathbb{Z}$ for every active n are orthogonal to any fixed polynomial space up to a negligible correction and realize essentially the positive Rayleigh quotient of T , which is at least that of the constant function,

$$\rho_L^+ = \frac{2}{L} \sum_{\log n < L} \frac{\Lambda(n)}{\sqrt{n}} (L - \log n),$$

and the certified ρ is within fifteen percent of ρ^+ at $L \approx 2$. On such a function the central form is about $(\log \omega - c - \rho_L) \|k\|^2$, so the tail is positive only above frequency $e^{\rho_L + c}$ and the head must reach it: N grows like $L e^{\rho_L}$.

Dense 6144-bit Grams at $N = 3000$ occupy about 7 GB each and cost a few days on one core; $L = 3$ is therefore a workstation project, $L = 3.5$ is not feasible with this code, and $L = 4$ is not feasible with this method. Since precision is not the barrier, the polynomial-complement method as implemented has a practical horizon near $L = 3$, that is, arithmetic support up to $n = 20$. Going further needs a different idea for the complement.

7.5 What finite-horizon positivity at fixed L can see

In the working normalization $Q_{0,L}[f] = 2 \sum_{\gamma > 0} |\widehat{f}_c(\gamma)|^2$. For a hypothetical off-line pair $\frac{1}{2} \pm \delta + it$, together with its conjugates, the same explicit formula contributes

$$\Delta(\delta, t) = 2 \operatorname{Re}[\widehat{f}_c(t + i\delta) \overline{\widehat{f}_c(t - i\delta)}] = 2(|\langle f, \cosh(\delta X) e^{itX} \rangle|^2 - |\langle f, \sinh(\delta X) e^{itX} \rangle|^2),$$

a difference of two squares whose negative part is at most $L \sinh^2(\delta L/2) \|f\|^2$, of order $(\delta L)^2 L \|f\|^2/4$ when δL is small.

Record R17 evaluates the zero sum for the lowest eigenvector f_0 of the 288-mode head at L_q (midpoints, mpmath at 60 digits, zeros from mpmath). The partial sums over the first K zeros reach 76%, 92%, 96.1% and 97.8% of $\lambda_0 = 3.284 \times 10^{-29}$ at $K = 100, 400, 1000, 2000$ (heights 237, 680, 1419, 2515); the remainder scales as T^{-1} in the cut-off height to within a percent of itself, which is the Fourier tail generated by the endpoint values $f_0(0) = f_0(L_q) = 1.8 \times 10^{-14}$, and extrapolates to 100% within the accuracy of the fit; f_0 is continuous across the join to 10^{-15} , as the endpoint-logarithm argument of Section 3.4 predicts. The transform $\widehat{f_0}$ is of order one below $\tau = 10$ and of order 10^{-3} between the first zeros, but it vanishes at the zeros themselves: $|\widehat{f_0}(\gamma_k)|$ is 6×10^{-28} , 2×10^{-26} , 10^{-25} , 2×10^{-24} at $k = 1, 2, 3, 4$ and still below 10^{-18} at $k = 13$ (height 59). The residual energy λ_0 is carried by zeros between heights 85 and 130, each contributing $1\text{--}3 \times 10^{-30}$, where $|\widehat{f_0}|$ has settled onto its endpoint-generated plateau of about 10^{-15} . In other words, the near-null vector of Q_{0,L_q} is a function of exponential type $L_q/2$ whose transform has been tuned to vanish at the low zeros, and $\lambda_{\min}(Q_{0,L})$ is the extremal value of this interpolation problem; its decay with L measures how many zeros a type- $L/2$ function can annihilate, which is why the compressed eigenvalue ladder falls so fast at these depths.

Two consequences follow. Displacing the k -th zero off the line by δ changes $Q_{0,L_q}[f_0]$ by $\Delta \approx 2(|\widehat{f_0}(\gamma_k)|^2 - \delta^2|\widehat{f_0}'(\gamma_k)|^2)$, which equals $-\lambda_0$ at $\delta_k^* \approx 10^{-12}$, 4×10^{-11} , 3×10^{-10} , 7×10^{-9} for $k = 1, \dots, 4$, at 10^{-3} for $k = 12$ (height 56), and at 0.1 for $k = 17$ (height 70). Read through the explicit formula, the certificate $Q_{0,L_q} \succeq 2.99 \times 10^{-29}I$ therefore pins the first few zeros to the critical line to many digits and says nothing about zeros above height about $70 \approx e^{2L_q}$: this is the classical relation between the support of a test function and the height it resolves. A pair placed where $\widehat{f_0}$ is not tuned to vanish contributes positively for every $\delta \leq 0.45$, the cosh term dominating. Finite-horizon positivity at depths 2 to 3 is thus compatible with, and could only be contradicted by zeros already excluded by, classical knowledge; a horizon comparable to $\log T$ is needed to see height T , and the milestones $L = 2, 3$ carry no information about zeros. This is not an objection to the programme, whose stated goal is an analytic all-depth argument, but it fixes what the numerical milestones mean. The agreement of the zero sum with λ_0 is, on the other hand, a Fourier-side check of the spatial head's normalization at the 10^{-29} level, far beyond the relative 6×10^{-10} check of [1] on a smooth test; completing it with a rigorous tail bound from the endpoint values is listed among the open tasks.

7.6 Open problems

The immediate tasks are: the author's normalization audit; an independent reconstruction of the spatial matrices and review of the analytic complement bounds; the two-interval second step for comparison and the choice of recursion invariant; stepwise estimates that prevent a schedule from accumulating at finite depth; and, beyond $L \approx 3$, a complement estimate that does not pay the arithmetic norm on high frequencies.

8 Records and reproducibility

The record identifiers are used throughout. Paths for R1–R9 are relative to `archive/drafts/v0.1_2026-09-10/numerics/history/`; the others are relative to the current `numerics/`. Exact file names, hashes, parameters and scopes are in `CLAIMS.json`; the four external matrix archives and their dual hashes are described in `ARCHIVES.md`.

ID	Claim and scope	Principal record
R1–R4	Generator, storage and whole-step diagnostics (Appendix A)	critical-path-20260910/, cumulative-path-20260910/, log7-log8-extension-20260910/
R5	All-new residual on 128 old modes, $\theta = 0.004$; the rational continuation	quarter-step-20260910/ residual_128_32.json
R6	Separate global floor 10^{-31} at 1.98	quarter-step-20260910/ reference_256.json
R7	Spatial full-operator floor 10^{-33} at L_q (superseded by R14)	quarter-step-closure-20260910/ closure_256_32.json
R8	Direct all-old residual factor 0.9 (superseded by R14)	quarter-step-closure-20260910/ relative_256_32.json
R9	Positive-shift consequences of R7 (superseded by Section 5)	quarter-step-closure-20260910/ path_corollary.json
R10	First-step comparison $\mu = 10^{-7}$ (superseded by R14)	records/metric-comparison-2048.json, -4096.json
R11	Second-step build; comparison $\mu = 10^{-8}$; failed absolute and residual tests	records/second-quarter-build.json, second-quarter-2048.json, -4096.json
R12	Separate-tail residual searches, unsuccessful	records/second-quarter-weighted-*.json
R13	Implementation consistency checks	records/spatial-consistency.json
R14	First-step bisections: absolute floor, relative floors at $\theta = 0.9, 0.8$, residual grid, comparison threshold	records/floor-bisection-first-2048.json
R15	Second-step bisection of the comparison threshold; recorded failures	records/floor-bisection-second-2048.json
R16	Ball Rayleigh upper bounds at L_q and L_2 ; hypothetical-floor diagnostics (not certificates)	records/rayleigh-upper-*.json, records/oracle-second-*.json
R18	First step rebuilt with 96 slab modes; second step $256 + 96 + 96$: absolute floor, relative floor, residual factor, comparison, upper bound	records/closure_256_96.json, records/second-quarter-96-build.json, records/ floor-bisection-second96-2048.json, records/ rayleigh-upper-second96-2048.json
R17	Fourier-side zero sums and zero-displacement sensitivity for the lowest eigenvector at L_q (diagnostic)	fourier_side/zero_sum_check_K400.json, -K2000.json, fourier_side/zero_sensitivity_K20.json

The recorded environment is Python 3.10–3.11, python-flint 0.9.0, FLINT 3.6.0, mpmath 1.3–1.4; assertions must remain enabled. The archive-based replays read the matrices through `numerics/recursion/stream_io.py`, a streaming loader that recomputes the canonical content hash while parsing and runs in a few gigabytes of memory; a corrupt or mismatched archive fails closed. The replay entry point `numerics/recursion/replay_floor.py` accepts chosen thresholds or brackets for bisection and writes the trace of every guarded test it runs; `build_step_seed.py` and `build_from_seed_v2.py` extend a first-step archive of any slab dimension, and `numerics/tools/` holds the continuation and upper-bound helpers used for R18. The recorded environment for R18 is a cloud container (Python 3.11, python-flint 0.9.0, FLINT 3.6.0); its two archives must be regenerated locally from the recorded commands, and are recognized by their content hashes. Independent small-example checks compare the moment assembly with quadrature (R13); a second implementation of the validators, written for the review of version 0.2, reproduced every recorded sign outcome and pivot diagnostic from the stored matrices. No second independent reconstruction

of the 256-plus-32 or 256-plus-32-plus-32 matrices has been made.

A Generator geometry and cumulative storage

This appendix collects the exact identities and diagnostics that motivated the spatial method. None is used by the theorems of the main text.

A.1 The hyperbolic identity and local extension

Put $X_L f(x) = (x - L/2)f(x)$ and $Q = Q_{0,L}$. Shift multiplies the off-diagonal gamma kernel and every arithmetic pairing by $\cosh(\omega(x - y))$, so on the common form domain

$$Q_{\omega,L}[f] = Q[\cosh(\omega X_L)f] - Q[\sinh(\omega X_L)f]. \quad (\text{A.1})$$

If $Q \succeq mI > 0$ then $Q_{\omega,L} \succeq 0$ if and only if $\|\tanh(\omega X_L)\|_{Q \rightarrow Q} \leq 1$, and with $\kappa_L = \|X_L\|_{Q \rightarrow Q}$, for $0 \leq \omega \kappa_L \leq \pi/4$ and $\omega \leq \frac{1}{2}$,

$$Q_{\omega,L} \succeq \cos(2\omega \kappa_L)Q, \quad \|V_{\omega,L}\| \leq \exp\left[-\frac{m}{2\kappa_L} \sin(2\kappa_L \omega)\right],$$

by absolute Taylor majorants ($\tanh(x) = -i \tan(ix)$, $\text{sech}(x) = \sec(ix)$). A computed finite-input κ_L is a lower bound, not the upper bound needed. For a positive block form with $c = \|F^{-1/2}BA^{-1/2}\| < 1$ one has $(1 - c)(A \oplus F) \preceq Q_{0,L+h} \preceq (1 + c)(A \oplus F)$, and an elementary local extension follows from $\|B\| \leq b_R$ and $F \succeq [-\gamma - \log(\pi h) - W_R h]I$ when $m_L[-\gamma - \log(\pi h) - W_R h] > b_R^2$; at the log 7 floor this can require $h = e^{-10^{31}}$. The lowest generator direction bends down initially: for $Qe = me$, $\frac{1}{2}\langle e, [X_L, [X_L, Q]]e \rangle = m\|X_L e\|^2 - Q[X_L e] \leq 0$, and at $L = \log 7$ an exact even polynomial of degree 126 gives $-2.999 \times 10^{-27} < Q_{10^{-11},L}[f]/\|f\|^2 < -2.998 \times 10^{-27}$ and $\kappa_L > 2.32 \times 10^{11}$ (R1). This is a negative instantaneous generator witness, not a negative central form or a noncontractivity witness.

A.2 Cumulative storage

The object needed for contraction is $D_{\omega,L} = I - V^*V = 2 \int_0^\omega Q_{s,L}[V_{s,L} \cdot] ds$. The same polynomial has positive storage, $8.717034818878968 \times 10^{-38} < D_{10^{-11}, \log 7}[f]/\|f\|^2 < 8.717034818878970 \times 10^{-38}$ (R2): negative generator energy and positive accumulated storage coexist on one input. At a spatial split, causality gives $V = \begin{pmatrix} X & 0 \\ Y & Z \end{pmatrix}$; with $E = I - X^*X$ and $F_{\text{out}} = I - ZZ^*$ strict contractions, elimination of the new input in $I - V^*V$ gives $S_D = E - Y^*F_{\text{out}}^{-1}Y$ and the full transfer is a strict contraction if and only if $c_D = \|F_{\text{out}}^{-1/2}Y E^{-1/2}\| < 1$.

Proposition A.1 (Cayley storage). *For $\omega > 0$ let $\mathcal{Z} = (I - V)(I + V)^{-1}$ and $P_{\omega,L} = \frac{2}{\omega} \text{Re } \mathcal{Z}$. Then $I + V$ is invertible, $P_{\omega,L}$ commutes with reflection, $D_{\omega,L} = \frac{\omega}{2}(I + V^*)P_{\omega,L}(I + V)$, and if the diagonal storage blocks of $P_{\omega,L+h} = \begin{pmatrix} A_\omega & B_\omega^* \\ B_\omega & F_\omega \end{pmatrix}$ are coercive then $\|F_\omega^{-1/2}B_\omega A_\omega^{-1/2}\| = c_D$.*

Proof. Finite-horizon Volterra convolution with an L^1 kernel is quasinilpotent, so -1 is outside the spectrum of V ; the identity for D is direct multiplication; real causality gives $V^* = R_L V R_L$ and hence reflection symmetry. With $R_X = (I + X)^{-1}$ and $R_Z = (I + Z)^{-1}$, block inversion gives $B_\omega = -\frac{2}{\omega} R_Z Y R_X$, $A_\omega = \frac{2}{\omega} R_X^* E R_X$, $F_\omega = \frac{2}{\omega} R_Z F_{\text{out}} R_Z^*$; the operators $U = \sqrt{2/\omega} F_\omega^{-1/2} R_Z F_{\text{out}}^{1/2}$ and $W = \sqrt{2/\omega} E^{1/2} R_X A_\omega^{-1/2}$ are unitary, and the normalized Cayley cross block equals $-U(F_{\text{out}}^{-1/2} Y E^{-1/2})W$. $\square$

The causal symbol is $a_{\text{cum},\omega}(p) = \frac{2}{\omega} [\mathcal{F}(p+\omega) - \mathcal{F}(p-\omega)] / [\mathcal{F}(p+\omega) + \mathcal{F}(p-\omega)]$, with the formal even expansion $a_0 + \omega^2(a_0''/6 - a_0^3/12) + O(\omega^4)$, $a_0 = 2\mathcal{F}'/\mathcal{F}$, on a core; $P_\omega = \frac{2}{\omega}I + \text{compact}$ is bounded while Q_0 is not, so no uniform tail estimate as $\omega \downarrow 0$ follows. For a finite step changing both depth and shift, with reference shift s and candidate t , the exact coercivity test is $I + H_F \succ 0$ and $I + H_A - (\mathcal{K} + H_B)^*(I + H_F)^{-1}(\mathcal{K} + H_B) \succ 0$, where $\mathcal{K} = F_s^{-1/2}B_sA_s^{-1/2}$ and H_A, H_B, H_F are the normalized changes of the blocks. This retains the signs of storage changes and is the robust formulation if a positive-shift certificate is ever attempted; none is attempted here.

A.3 Why the whole step is difficult

At $a = \log 7$ and $h_{\max} = \log(8/7)$, finite diagnostics with 128 old and 32 new modes put the normalized gamma and arithmetic pieces of the cross block separately near 2.1554×10^{13} while their sum has norm near one, and an exact rational test pair proves the unrestricted lower bound $c_{\text{central,full}}^2 > 1 - 1.84 \times 10^{-17}$ on the coupling (R4). The finite-input slack $1 - c_{128,32}^2$ is 1.27×10^{-7} , 1.78×10^{-11} and 1.84×10^{-17} for steps of one quarter, one half and the whole of $\log(8/7)$; this motivated the quarter-step. A uniform short-slab storage bound, $\|V_{\omega,h}^\gamma\| \leq e^{-0.06\omega}$ for $0 < h \leq h_{\max}$ and $0 < \omega \leq \frac{1}{2}$, follows from the positivity of the shifted profile $G_\omega^V > 0.4935$ on $[0, h_{\max}]$ and the inequality $-\log \Gamma(1+\omega)/\omega \leq \gamma - (\pi^2/4 - 2)\omega$ on $(0, \frac{1}{2}]$ (R4; the constant $\pi^2/4 - 2$ is a valid but not sharp replacement for the leading term $\zeta(2)/2$ of the Weierstrass product).

B Full-output storage construction

The cumulative calculations use the gamma profile

$$g(t) = e^{t/2} \left(\frac{\sinh t}{t} \right)^{\omega-1}, \quad y_c(t) = \omega \int_0^1 v^{\omega-1} e^{ct(1-v)} g(tv) dv, \quad G_\omega^V(t) = g(t) - 4t\{\alpha y_\alpha(t) + \beta y_{-\beta}(t)\},$$

with $\alpha = \frac{1}{2} - \omega$, $\beta = \frac{1}{2} + \omega$, and the stable recurrence $y_{c,0} = 1$, $y_{c,j} = (\omega g_j + c y_{c,j-1})/(\omega + j)$. On $|z| = 3$, $|z/\sinh z| \leq 3/\sin 3 < 24$ and $|G_\omega^V(z)| < 32768$ for $0 < \omega \leq \frac{1}{2}$, so degree d gives the gamma operator error $\delta_\gamma \leq \frac{(2\pi L)^\omega}{\Gamma(\omega)} \frac{32768(L/3)^{d+1}}{(d+1+\omega)(1-L/3)}$ and the full error $\delta = (\sum_{\log n < L} b_\omega(n))\delta_\gamma$. Delayed polynomial outputs are integrated with the moments $J_k(d_0, \ell) = \int_0^\ell t^{k+\omega} (t+d_0)^\omega dt$, initialized by an enclosed hypergeometric value and continued by $J_{k+1} = [\ell^{k+\omega+1}(\ell+d_0)^{\omega+1} - d_0(k+\omega+1)J_k]/(k+2\omega+2)$. If $\tilde{G}$ is the model Gram on orthonormal retained inputs, the exact defect differs by at most $2\delta\sqrt{\text{tr } \tilde{G}} + \delta^2$ in operator norm on that input space; this is subtracted before testing positivity, and it is why “finite inputs, full output” cannot be enlarged to omitted inputs without a further argument.

C A compact map of the logical dependencies

The framework (2.1)–(2.3) supports two separate full-depth calculations at L_q . The global method of [1] at 1.98 gives R6 and its shift interval. The spatial method uses the old-depth matrices at $\log 7$, the arithmetic bound (4.1), the tail floors (4.2)–(4.3), and the full-output Grams; the absolute test gives the floor of Theorem 1.1, the relative test gives the residual factor and, through Lemma 3.2, an independent floor, and the comparison test gives $H_J \succeq \mu A$. Neither spatial test depends on R6; the relative test does not depend on the sign of the absolute test; R5 supplies the rational continuation but its factor 0.004 is not a premise. The upper bounds are ball Rayleigh quotients and depend on nothing but the head. At L_2 the comparison test and the upper bound are certified; the floor and the residual factor are not, and the hypothetical-floor runs of Section 6.3 are diagnostics with no

logical weight. The conditional theorem uses Lemma 3.2 and the framework only. Appendix A is motivational.

Acknowledgment of computational assistance

This draft and its numerical implementation were prepared with substantial assistance from language models, under the author’s direction. OpenAI models drafted versions 0.1 and 0.2 and their research packets; Anthropic’s Claude reviewed version 0.2 with an independent re-implementation of the validators, found the understated floors and the second-step diagnosis, wrote the archive-based replay and streaming loader, and drafted version 0.3. The record distinguishes analytic proofs, validated arithmetic, and exploratory diagnostics. No result has been human-verified end to end or formally verified.

References

- [1] E. Baker, *Finite-horizon Weil coercivity and shifted-zeta contraction*, working draft v0.4, 10 September 2026, `papers/weil-depth/` of the repository <https://github.com/ebbaker/shifted-zeta-positivity>. The v0.3 source used for the constants of this paper is preserved at `archive/drafts/v0.1_2026-09-10/reference/weil-depth-v0.3/`.
- [2] *Depth-shift research packets*, 10 September 2026: critical path; cumulative storage; log 7–log 8 extension; quarter-step residual; quarter-step closure. Preserved under `archive/drafts/v0.1_2026-09-10/numerics/history/`.
- [3] Review of version 0.2 and note on improvements and next steps, 10 September 2026, `archive/reviews/`.