

Positive factorizations and supersymmetric extensions of finite-interval Weil forms

Working draft prepared for Edward Baker

Version 0.1 11 September 2026

Abstract

We investigate whether positive auxiliary systems can explain the sign of finite-interval Weil forms arising from the shifted Riemann zeta function. The objective is an independently specified square-root or bulk construction, followed by an exact arithmetic identification. We give an explicit nonlocal factorization of the prime-free form, uniform for total interval length $L \leq 1/4$ and shift $0 \leq \omega \leq 1/2$, with lower bound greater than 0.099. The gamma kinetic multiplier admits an exact realization by a positive tower of local auxiliary fields and a closed graded supercharge. We also prove restrictions on proposed completions. Scalar gauge changes cannot remove the off-diagonal sign obstruction. More generally, positive two-point square factors fail already at $L = 2/3$, as established by a comparison-form test with an exact rational certificate. Comparison positivity survives elimination of internal nodes in finite pairwise networks. A coherent multicomponent toy model escapes this restriction. We identify additional boundary and first-prime matching constraints, and show that no closable norm factor of the full Weil form exists on ordinary whole-line L^2 with all smooth compactly supported inputs in its domain. These results delimit a supersymmetric research program; they supply neither an all-depth positivity theorem nor a proof of the Riemann hypothesis.

Draft status. This manuscript consolidates two exploratory investigations. Proofs are included, and one sign argument has a reproducible exact rational checker. There has been no independent specialist review or complete priority audit. The graded constructions are linear operator models, not a constructed interacting superspace field theory. The small-window result does not extend the stronger finite-depth certificates already recorded in the project.

Keywords. Weil positivity; auxiliary fields; supersymmetric quantum mechanics; nonlocal forms; Schur complements; comparison matrices.

Contents

1	Motivation and the question of mechanism	3
1.1	From higher-dimensional extensions to positive boundary systems	3
1.2	Why a supersymmetric extension is a reasonable test	3
1.3	Results and limits of the present paper	4
2	The forms, domains, and graded convention	4
3	An explicit positive factor on a small window	5
3.1	The full-output jump identity	5
3.2	The positive range and shift-uniform estimate	6
3.3	Relation to the shift-energy identity	7
4	Two restrictions on direct scalar factors	7
5	A positive auxiliary-field realization of the gamma kinetic term	8
6	A comparison-form obstruction to pairwise factors	9
6.1	The class of factors and its necessary condition	9
6.2	A counterexample at total length $2/3$	10
6.3	Eliminating hidden nodes in finite networks	11
7	Coherent couplings and two completion constraints	11
7.1	A positive graded model outside the pairwise class	11
7.2	The pole term and a growing Green function	12
7.3	The first prime forces joint diagonal accounting	12
8	The input space of a global norm identity	13
9	What a useful supersymmetric completion must establish	14
9.1	The boundary object must be specified	14
9.2	Two concrete milestones	14
9.3	What should be retained if this route stops	14
A	Exact rational certificate for the comparison test	15
B	Verification, provenance, and remaining review	16

1 Motivation and the question of mechanism

1.1 From higher-dimensional extensions to positive boundary systems

The research program motivating this paper began with an attempt to extend the completed xi function into higher-dimensional settings and constrain its zeros by rotational symmetry. Investigations using slice-regular and related extensions repeatedly encountered the same difficulty: the proposed extension encoded information already present in the original complex function. An additional coordinate did not, by itself, create an additional constraint on the zeros. This is the history reported by the researcher, not a general impossibility theorem about higher-dimensional approaches.

The search then moved toward physical models, including gauge theory and ideas inspired by bulk–boundary correspondences. The enduring question was whether an arithmetic boundary observable could be derived from an independently positive system. Here a “bulk” may be a linear field system, an auxiliary channel space, or a network. The term does not assert an AdS geometry or a conformal field theory. A sufficiently explicit positive linear model would already be useful if its boundary response were exactly the required zeta object.

The shifted zeta program supplies concrete targets for this question: causal transfers, finite-interval generators, contraction defects, and string or spectral representations. Its existing finite-depth work proves restricted inequalities by analytic estimates and certificates [1, 2]. Those results provide benchmarks for an explanatory construction. They do not establish positivity at every depth, and a sequence of finite certificates is not itself a rule forcing the next certificate to succeed.

The intended direction of explanation is

specified arithmetic data $\longrightarrow$ independently positive system $\longrightarrow$ exact boundary identity.

The global quantifiers needed for the Riemann hypothesis (RH) come only after these steps and require a separate limiting or compatibility theorem. Reconstructing a positive system from a measure already assumed positive does not supply the missing direction of implication.

1.2 Why a supersymmetric extension is a reasonable test

Supersymmetric quantum mechanics suggests beginning with a defining operator A and a positive inner product. Its adjoint then yields a nonnegative Hamiltonian A^*A , or a graded Hamiltonian with two partner blocks. The substantive work is to define A independently and identify the resulting energy or observable with the arithmetic target.

Signed terms in a displayed operator do not rule out such a mechanism. For example, on a compactly supported smooth core, a real smooth function W gives

$$(\partial_x + W)^*(\partial_x + W) = -\partial_x^2 + W^2 - W'. \quad (1.1)$$

The potential need not be pointwise positive, but the terms are constrained by a common square. On an interval the same statement also requires the correct domain and boundary terms. The present problem is to find an analogous constraint relating the gamma, pole, and prime contributions.

The positive component-space interpretation of fermionic variables is standard in supersymmetric quantum mechanics [5]. A formal Berezin integral is not a positive measure, and a supertrace is a difference of ordinary traces. Boson–fermion cancellation therefore does not establish positivity of an arbitrary reduced observable. Likewise, nilpotence of a BRST differential does not establish positivity on its physical quotient. A BRST construction would require a specific gauge redundancy and a separate argument about the induced inner product. No such gauge system is assumed here.

1.3 Results and limits of the present paper

Theorem 3.1 constructs a positive factor on a small prime-free window. Theorem 5.1 realizes the gamma kinetic term through local auxiliary fields; the reduction produces the correct nonlocal boundary order. Theorem 6.3 excludes a larger class of pairwise factors, and Proposition 6.4 explains why adding internal nodes to a finite pairwise network does not automatically help. Section 7 exhibits a coherent positive model outside that class and identifies two elementary completion failures. Theorem 8.1 addresses the input space for a global factor.

These statements have different scopes. A small-interval construction does not prove all-depth positivity. A restriction on a class of factors does not prove negativity of the Weil form. A coherent toy model establishes possibility within a larger class, not an arithmetic selection principle. The proposed continuation in Section 9 is kept separate from the proved statements.

There is substantial prior literature on Weil forms and Hilbert-space realizations. In particular, Suzuki gives norm-identity criteria related to RH [6] and a screw-function treatment of finite-interval operators and small-window arguments [7]. We do not claim that the possibility of a norm formulation, a graded block construction, or a small-window Dirichlet argument is new. The application-specific formulas and restrictions below are recorded with proofs; their priority requires further review.

2 The forms, domains, and graded convention

Let $I_L = (-L/2, L/2)$, where L is the *total* interval length. Extend $f \in L^2(I_L)$ by zero to $\tilde{f}$ on $\mathbb{R}$, and use

$$\hat{f}(\tau) = \int_{I_L} f(x) e^{-i\tau x} dx, \quad \mathcal{D}_{\log, L} = \left\{ f : \int_{\mathbb{R}} \log(2 + |\tau|) |\hat{f}(\tau)|^2 d\tau < \infty \right\}. \quad (2.1)$$

We use inner products conjugate-linear in the first variable. Write

$$C(f) = \int_{I_L} f(x) \cosh(x/2) dx, \quad S(f) = \int_{I_L} f(x) \sinh(x/2) dx. \quad (2.2)$$

For $a > 0$, define the truncated translation $(T_a f)(x) = \tilde{f}(x - a)$ for $x \in I_L$.

The central finite-interval form, in the normalization of [2], is

$$\begin{aligned} Q_{0,L}[f] = & \frac{1}{2\pi} \int_{\mathbb{R}} \left(\operatorname{Re} \psi\left(\frac{1}{4} + \frac{i\tau}{2}\right) - \log \pi \right) |\hat{f}(\tau)|^2 d\tau \\ & + 2|C(f)|^2 - 2|S(f)|^2 - \sum_{\log n < L} \frac{\Lambda(n)}{\sqrt{n}} \langle f, (T_{\log n} + T_{\log n}^*) f \rangle. \end{aligned} \quad (2.3)$$

Here $\psi = \Gamma'/\Gamma$, and Λ is the von Mangoldt function. The prime-free form $Q_{0,L}^\gamma$ omits the last sum and agrees with $Q_{0,L}$ for $L \leq \log 2$. The Fourier multiplier differs from $\log(2 + |\tau|)$ by a bounded function [3]; the other terms are bounded on each fixed interval. Thus these forms are closed and semibounded on $\mathcal{D}_{\log, L}$, regardless of their sign. Smooth compactly supported functions form a core: first shrink the support by dilation, then mollify; the weighted Fourier norm converges in each step.

Set

$$w_0 = \psi(1/4) - \log \pi = -\gamma_E - \pi/2 - 3 \log 2 - \log \pi, \quad (2.4)$$

$$j(t) = \frac{e^{-t/2}}{1 - e^{-2t}}, \quad R(t) = 2 \cosh(t/2) - j(t), \quad t > 0. \quad (2.5)$$

The shifted gamma form can be defined without an endpoint ambiguity by

$$Q_{\omega,L}^\gamma[f] = Q_{0,L}^\gamma[f] + \int_{I_L^2} (\cosh(\omega|x-y|) - 1) R(|x-y|) \overline{f(x)} f(y) dx dy, \quad (2.6)$$

for $0 \leq \omega \leq 1/2$. This is the bounded perturbation identity proved for the finite-horizon shift generator in [1]. The kernel is continuous across the diagonal after assigning its limiting value there. In particular, (2.6) includes the distributional correction required at $\omega = 1/2$ in a Fourier-plus-residue description.

Lemma 2.1 (Graded realization). *Let $A : \text{Dom } A \subset \mathcal{H}_0 \rightarrow \mathcal{H}_1$ be closed and densely defined between positive Hilbert spaces. Then*

$$\mathcal{S} = \begin{pmatrix} 0 & A^* \\ A & 0 \end{pmatrix}, \quad \text{Dom } \mathcal{S} = \text{Dom } A \oplus \text{Dom } A^*, \quad (2.7)$$

is self-adjoint and

$$\mathcal{S}^2 = \text{diag}(A^*A, AA^*) \geq 0. \quad (2.8)$$

Equivalently, $q(f, g) = (0, Af)$, with domain $\text{Dom } A \oplus \mathcal{H}_1$, is nilpotent and $\{q, q^*\} = \mathcal{S}^2$ on the natural operator domain.

Proof. Computing the adjoint of the off-diagonal block gives the same block and domain because $A^{**} = A$. Squaring on its natural domain gives (2.8). The nilpotent formulation is the same calculation. $\square$

This lemma organizes an independently constructed factor. Choosing $A = Q^{1/2}$ for an operator whose sign is unknown would assume the premise we seek to establish.

3 An explicit positive factor on a small window

3.1 The full-output jump identity

The digamma difference integral [3, §5.9] gives

$$\text{Re } \psi(1/4 + i\tau/2) - \log \pi = w_0 + 2 \int_0^\infty j(t)(1 - \cos \tau t) dt. \quad (3.1)$$

If U_t denotes translation on $L^2(\mathbb{R})$, Plancherel implies

$$Q_{0,L}^\gamma[f] = w_0 \|f\|^2 + \int_0^\infty j(t) \|\tilde{f} - U_t \tilde{f}\|_{L^2(\mathbb{R})}^2 dt + 2|C(f)|^2 - 2|S(f)|^2. \quad (3.2)$$

The norm includes output outside I_L . Dropping that output would change the boundary contribution and invalidate the subsequent factorization.

Define

$$F(a) = \int_a^\infty j(t) dt = \text{artanh}(e^{-a/2}) + \arctan(e^{-a/2}), \quad a > 0, \quad (3.3)$$

and put $h(t) = -R(t)$. Splitting the jump integral into interior and exterior pairs, then expanding the pole kernel, gives

$$Q_{0,L}^\gamma[f] = \frac{1}{2} \int_{I_L^2} h(|x-y|) |f(x) - f(y)|^2 dx dy + \int_{I_L} \kappa_{0,L}(x) |f(x)|^2 dx, \quad (3.4)$$

$$\kappa_{0,L}(x) = w_0 + F(x + L/2) + F(L/2 - x) + 8 \sinh(L/4) \cosh(x/2). \quad (3.5)$$

The endpoint logarithms in F are part of the form. They are integrable against smooth functions. Identity (3.4) remains valid when its weights lose positivity.

3.2 The positive range and shift-uniform estimate

Theorem 3.1 (Small-window factor). *For $0 < L \leq 1/4$ and $0 \leq \omega \leq 1/2$, define*

$$h_\omega(t) = \cosh(\omega t)h(t), \quad (3.6)$$

$$\kappa_{\omega,L}(x) = \kappa_{0,L}(x) - \int_{I_L} (\cosh(\omega|x-y|) - 1)h(|x-y|)dy. \quad (3.7)$$

Both weights are nonnegative, and

$$\kappa_{\omega,L}(x) \geq c_* := \log(4/\pi) - \gamma_E + \frac{7}{16} - \frac{e^{1/8} \cosh(1/8)}{512} > 0.099. \quad (3.8)$$

The closed map from $L^2(I_L)$ to $L^2(I_L^2) \oplus L^2(I_L)$ given by

$$\mathcal{A}_{\omega,L}f = \left(\sqrt{h_\omega(|x-y|)/2} [f(x) - f(y)], \sqrt{\kappa_{\omega,L}(x)}f(x) \right) \quad (3.9)$$

has domain $\mathcal{D}_{\log,L}$ and satisfies

$$Q_{\omega,L}^\gamma[f] = \|\mathcal{A}_{\omega,L}f\|^2 \geq c_* \|f\|^2. \quad (3.10)$$

Proof. The sign of $R(t)$ is that of $q^3 - q - 1$ with $q = e^t$. Its unique positive sign-change point is

$$t_0 = \log \rho = 0.2811995743 \dots, \quad \rho^3 = \rho + 1. \quad (3.11)$$

In particular, $h(t) > 0$ for $0 < t \leq 1/4$.

Since j decreases, F is convex. Both the sum of the two F terms in (3.5) and its hyperbolic-cosine term attain their minimum at $x = 0$. Using elementary hyperbolic identities, this minimum is

$$-\gamma_E - \log(8\pi) + \log \coth(L/8) - 2 \arctan(\tanh(L/8)) + 8 \sinh(L/4). \quad (3.12)$$

The inequalities $\coth u \geq 1/u$, $\arctan(\tanh u) \leq u$, and $\sinh u \geq u$ give

$$\kappa_{0,L}(x) \geq -\gamma_E - \log(\pi L) + 7L/4 \geq \log(4/\pi) - \gamma_E + 7/16 =: c_0, \quad (3.13)$$

where the last inequality uses monotonic decrease of the preceding expression for $L \leq 1/4$.

Expanding the difference squares shows that changing h to h_ω produces the off-diagonal perturbation in (2.6); the subtraction in (3.7) cancels the extra diagonal. Moreover,

$$0 \leq h(t) \leq j(t) = \frac{e^{t/2}}{2 \sinh t} \leq \frac{e^{1/8}}{2t}, \quad \cosh(\omega t) - 1 \leq \frac{t^2}{8} \cosh(1/8).$$

Since $\int_{I_L} |x-y|dy \leq L^2/2$, the subtracted potential is at most $e^{1/8} \cosh(1/8)/512$. This proves (3.8).

The norm identity follows first on the smooth core from (3.4). The corresponding graph norm is equivalent to the closed positive form norm on $\mathcal{D}_{\log,L}$. Closing the map on that core gives a closed operator with this domain and preserves the identity. The sign estimate precedes this closure argument; no unknown spectral square root is used. $\square$

3.3 Relation to the shift-energy identity

On the smooth core, the finite-horizon transfer satisfies $\partial_\omega V_{\omega,L} = -G_{\omega,L} V_{\omega,L}$, with symmetric generator form $Q_{\omega,L}^\gamma$ in the prime-free region. Integration of the energy derivative, followed by the form limit, gives

$$\|f\|^2 - \|V_{\omega,L} f\|^2 = 2 \int_0^\omega \|\mathcal{A}_{s,L} V_{s,L} f\|^2 ds, \quad (3.14)$$

in the domain and limiting sense of [1]. Equation (3.14) is a consequence of that transfer identity and the new explicit factor, not a separately constructed bulk duality. Its range is only a subset of the already certified prime-free slab.

4 Two restrictions on direct scalar factors

Proposition 4.1 (Scalar conductances and scalar gauges). *For $L > t_0$, the gamma form cannot be written as nonnegative scalar conductances multiplying $|f(x) - f(y)|^2$, plus an arbitrary multiplication form. A positive scalar rescaling, or a single position-dependent phase change of the input, does not remove this obstruction.*

Proof. A nonnegative difference conductance has nonpositive off-diagonal kernel, whereas $R(|x - y|)$ is positive on an open set when $L > t_0$. A positive rescaling preserves these signs. For a phase change, choose $t_0/2 < d < \min(t_0, L/2)$ and three interior points with consecutive separation d . The signs of the three kernel entries are $-, -, +$. Their product is positive and is invariant under a scalar phase gauge. Three negative real entries would have negative product. The argument can be made on small separated neighborhoods, so it is an obstruction to an almost-everywhere kernel identity, not merely to values at three points. $\square$

Independent edge phases are more general than a scalar gauge. They are addressed in Section 6; the preceding argument alone does not exclude them.

Proposition 4.2 (Local first-order boundary factors). *No finite-component map $Af = a(x)f' + b(x)f$, with locally bounded smooth coefficients and the ordinary positive component L^2 norm, has $\|Af\|^2 = Q_{0,L}[f]$ on all $C_c^\infty(I_L)$.*

Proof. For fixed $\chi \in C_c^\infty(I_L)$, put $f_N = \chi e^{iNx}$. The digamma asymptotic and the bounded remaining terms imply

$$Q_{0,L}[f_N] = \log N \|\chi\|^2 + O(1). \quad (4.1)$$

For the multiplier term, translate the Fourier variable by N and use rapid decay of $\hat{\chi}$ outside $|\tau - N| \leq N/2$. If a is nonzero, choose χ where $\|a\chi\| > 0$; then $\|Af_N\|^2 = N^2 \|a\chi\|^2 + O(N)$, a contradiction. If a vanishes identically, the left side remains bounded in N , also a contradiction. The same argument applies to a finite vector of rows. $\square$

This proposition concerns the square root acting directly on the original boundary coordinate. Eliminating local auxiliary variables can change the order of the reduced operator, as the next construction demonstrates.

5 A positive auxiliary-field realization of the gamma kinetic term

Define $a_k = 2k + 1/2$ and

$$B(s) = \sum_{k=0}^{\infty} \frac{2}{a_k} \frac{s}{a_k^2 + s}, \quad s \geq 0. \quad (5.1)$$

The geometric expansion $j(t) = \sum_{k \geq 0} e^{-a_k t}$ and the digamma difference integral give

$$B(\tau^2) = \operatorname{Re} \psi(1/4 + i\tau/2) - \psi(1/4). \quad (5.2)$$

Consequently the nonnegative kinetic form is

$$K[f] = \frac{1}{2\pi} \int_{\mathbb{R}} B(\tau^2) |\hat{f}(\tau)|^2 d\tau. \quad (5.3)$$

Theorem 5.1 (Positive local tower). *For $a > 0$, let*

$$\mathcal{E}_a[f, u] = \frac{2}{a} \int_{\mathbb{R}} (|f - u|^2 + a^{-2} |u'|^2) dx, \quad f \in L^2(\mathbb{R}), \quad u \in H^1(\mathbb{R}). \quad (5.4)$$

Then, with extended values allowed,

$$K[f] = \inf_{(u_k)} \sum_{k=0}^{\infty} \mathcal{E}_{a_k}[f, u_k]. \quad (5.5)$$

For $K[f] < \infty$, the infimum is attained by

$$\hat{u}_k(\tau) = \frac{a_k^2}{a_k^2 + \tau^2} \hat{f}(\tau). \quad (5.6)$$

The unreduced energy is the norm square of a closed densely defined map between positive component Hilbert spaces and therefore has the graded realization of Lemma 2.1.

Proof. At a single frequency, completion of the square gives

$$\begin{aligned} |f - u|^2 + \frac{\tau^2}{a^2} |u|^2 &= \frac{\tau^2}{a^2 + \tau^2} |f|^2 \\ &\quad + \left(1 + \frac{\tau^2}{a^2}\right) \left|u - \frac{a^2}{a^2 + \tau^2} f\right|^2. \end{aligned} \quad (5.7)$$

Plancherel gives the minimum of (5.4). Summing and using monotone convergence proves (5.5). Each multiplier in (5.6) maps L^2 into H^1 , and the minimum has finite total energy whenever $K[f]$ is finite.

For a precise infinite-component space, set

$$\mathcal{H}_B = L^2(\mathbb{R}) \oplus \ell^2(L^2(\mathbb{R}); 2/a_k^3), \quad \mathcal{H}_F = \ell^2(L^2(\mathbb{R}) \oplus L^2(\mathbb{R})), \quad (5.8)$$

and define

$$A(f, (u_k)) = \left(\sqrt{2/a_k} (f - u_k), \sqrt{2/a_k^3} u'_k \right)_{k \geq 0}. \quad (5.9)$$

Give A its maximal domain: all $u_k \in H^1$ for which the displayed sequence belongs to $\mathcal{H}_F$. It is closed. Indeed, convergence in $\mathcal{H}_B$ gives convergence of every component in L^2 , and convergence of

A gives convergence of every weak derivative; closedness of weak differentiation identifies the limit and its residuals.

The domain is dense. Choose smooth f , let finitely many smooth u_k be independent, and set $u_k = f$ in the tail. The tail derivative energy is finite since $\sum a_k^{-3} < \infty$. Such vectors are dense in the weighted product space by truncation and approximation. Finally, the minimizers satisfy $\|u_k\| \leq \|f\|$, so their auxiliary Hilbert norm is finite; their derivative and difference energies sum to $K[f]$. Lemma 2.1 now applies to (5.9). $\square$

The fields u_k live on the whole line even when f is supported in I_L . Truncating them at the input endpoints changes the minimization. The square-root map uses only first derivatives and multiplication in the unreduced coordinate. Its reduced boundary multiplier is logarithmic, which explains precisely why the direct order argument of Section 4 does not apply.

Corollary 5.2 (An extra-coordinate extension). *The function B in (5.1) is a complete Bernstein function. Hence $B(-\partial_x^2)$ has a positive extension realization of the type established by Kwaśnicki and Mucha, allowing a measure-valued coefficient.*

Proof. The representation

$$B(s) = \int_0^\infty \frac{s}{s+r} \nu(dr), \quad \nu = \sum_{k \geq 0} \frac{2}{a_k} \delta_{a_k^2},$$

has a positive measure satisfying $\int (1+r)^{-1} \nu(dr) < \infty$. This is the complete Bernstein representation, and the extension conclusion follows from [4], including its general measure-coefficient theorem. $\square$

The extension coefficient is not computed here. This corollary concerns the explicitly positive function B , not the shifted zeta Weyl function. Moreover, the exact gamma form still reads

$$Q_{0,L}^\gamma[f] = K[\tilde{f}] + w_0 \|f\|^2 + 2|C(f)|^2 - 2|S(f)|^2. \quad (5.10)$$

The negative normalization and odd pole contribution remain. They cannot be removed by changing the zero of energy without changing the target. The positive tower is an ingredient for a completion, not the completion. Related positive diffusive gamma expansions already appear in [1]; the point here is the explicit variational and graded realization at the generator level.

6 A comparison-form obstruction to pairwise factors

6.1 The class of factors and its necessary condition

The scalar gauge obstruction does not control arbitrary phases attached to edges. A stronger test uses the diagonal contributions forced by each positive two-point square.

Definition 6.1 (Positive pairwise representation). A positive pairwise representation on I is a sum or integral of terms

$$\|u(x, y)f(x) + v(x, y)f(y)\|^2, \quad x < y, \quad (6.1)$$

plus a nonnegative multiplication form. The coefficient vectors may have arbitrary positive Hilbert component spaces. After grouping channels for each pair, their Gram coefficients are locally integrable away from $x = y$. The nonnegative integral is finite on the smooth test core and equals the proposed form there. Negative renormalizing counterterms are not permitted as additional energy terms in this definition.

For a fixed pair the Gram matrix has diagonal entries $\alpha = \|u\|^2$, $\beta = \|v\|^2$, and off-diagonal entry $c = \langle u, v \rangle$. Cauchy–Schwarz gives $|c|^2 \leq \alpha\beta$. Replacing c by $-|c|$ therefore preserves positivity of this two-by-two matrix. Different endpoint magnitudes and independent edge phases are all allowed in Definition 6.1.

Lemma 6.2 (Necessary comparison positivity). *If $Q_{0,L}^\gamma$ has a positive pairwise representation, then*

$$Q_{\text{cmp},L}[f] := Q_{0,L}^\gamma[f] - 2 \int_{I_L^2} R_+ (|x - y|) \overline{f(x)} f(y) dx dy \quad (6.2)$$

is nonnegative on nonnegative real smooth supported inputs. Here $R_+ = \max(R, 0)$.

Proof. Polarization with disjointly supported tests identifies the grouped off-diagonal coefficient with $R(|x - y|)$ almost everywhere away from the diagonal. For nonnegative real inputs, replacing that coefficient by $-|R|$ subtracts exactly the second term in (6.2). Each grouped pair remains positive by the preceding Gram-matrix argument, and the nonnegative multiplication term is unchanged. The subtraction has a bounded kernel, so the argument does not split the singular diagonal into separately divergent integrals. $\square$

6.2 A counterexample at total length $2/3$

Theorem 6.3 (Failure of positive pairwise factors). *The form $Q_{0,2/3}^\gamma$ has no representation of Definition 6.1. In particular, independent edge phases, unequal endpoint weights, and arbitrary separate vector channels for each pair cannot provide such a factorization.*

Proof. Take $f = L^{-1/2} \mathbf{1}_{I_L}$. It belongs to the logarithmic form domain. The jump identity and cancellation of the exponential tail give

$$Q_{0,L}^\gamma[f] = w_0 + \frac{2}{L} \sum_{k=0}^{\infty} \frac{1 - e^{-a_k L}}{a_k^2} + \frac{32}{L} \sinh^2(L/4), \quad (6.3)$$

while

$$Q_{\text{cmp},L}[f] = Q_{0,L}^\gamma[f] - \frac{4}{L} \int_0^L (L - t) R_+(t) dt. \quad (6.4)$$

For $L = 2/3$ and $b = 1/3$, the right side is at most

$$\mathcal{U} := Q_{0,2/3}^\gamma[f] - 6 \int_{1/3}^{2/3} (2/3 - t) R(t) dt. \quad (6.5)$$

The exact rational enclosure established in Appendix A proves $\mathcal{U} < -1/25$.

Choose nonnegative smooth interior cutoffs converging to the constant. Each endpoint error is a scaled fixed profile of width ε ; changing variables in its weighted Fourier norm gives an upper bound $O(\varepsilon(1 + \log(1/\varepsilon)))$. Thus the cutoffs converge in the logarithmic form norm. The bounded-kernel term in (6.2) also converges. A sufficiently close smooth cutoff therefore has negative comparison form, contradicting Lemma 6.2. $\square$

For orientation, separate quadratures give

$$Q_{0,2/3}^\gamma[f] \simeq 0.17402032645, \quad Q_{\text{cmp},2/3}[f] \simeq -0.06189510026.$$

These decimal values are diagnostics; the proof uses the weaker exact bound on $\mathcal{U}$. Since $2/3 < \log 2$, the obstruction occurs before any prime is active. It concerns a class of positive representations, not negativity of the gamma form itself.

6.3 Eliminating hidden nodes in finite networks

For a Hermitian matrix G , let $M(G)$ have the same diagonal and off-diagonal entries $-|G_{ij}|$.

Proposition 6.4 (Comparison positivity survives a Schur reduction). *If $M(G) \succeq 0$, then eliminating internal nodes by positive-pivot Schur complements preserves comparison positivity. In particular, every finite positive network assembled from two-point squares and nonnegative one-point terms has a boundary Schur complement S with $M(S) \succeq 0$.*

Proof. Eliminate one node with diagonal $d > 0$. The resulting entries are $S_{ij} = G_{ij} - G_{i0}G_{0j}/d$. The diagonal of $M(S)$ equals that of the Schur complement N of $M(G)$, and, for $i \neq j$,

$$-|S_{ij}| \geq -|G_{ij}| - |G_{i0}||G_{0j}|/d = N_{ij}. \quad (6.6)$$

Both $M(S)$ and N have nonpositive off-diagonal entries. For every complex vector z , writing $|z|$ componentwise gives

$$z^* M(S) z \geq |z|^T M(S) |z| \geq |z|^T N |z| \geq 0. \quad (6.7)$$

The last step follows from positivity of the Schur complement of $M(G)$. A zero diagonal in a positive semidefinite comparison matrix has a zero row and column, so such nodes decouple. Iterate the argument. Finally, the grouped two-by-two Gram argument proves $M(G) \succeq 0$ for a positive pairwise network. $\square$

The proposition is finite dimensional. It passes to a limit if the comparison forms themselves converge on the relevant tests. No claim is made here that every singular continuum extension admits that passage. The theorem does establish that adding arbitrarily many nodes to a finite scalar magnetic spring network does not, by itself, evade the comparison restriction.

7 Coherent couplings and two completion constraints

7.1 A positive graded model outside the pairwise class

Consider three complex boundary amplitudes and one auxiliary amplitude:

$$\mathcal{E}(f, u) = |f_1 + f_2 - u|^2 + |f_3 + u|^2. \quad (7.1)$$

Completion of the square gives

$$\inf_u \mathcal{E}(f, u) = \frac{1}{2} |f_1 + f_2 + f_3|^2, \quad u = \frac{1}{2} (f_1 + f_2 - f_3). \quad (7.2)$$

The boundary matrix is half the all-ones matrix; its eigenvalues are $0, 0, 3/2$. Its comparison matrix has eigenvalues $-1/2, 1, 1$. Thus a positive bulk can produce a boundary form outside the pairwise comparison class.

The independently specified factor is

$$A = \begin{pmatrix} 1 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \end{pmatrix}. \quad (7.3)$$

Lemma 2.1 supplies its graded realization. The first row mixes three scalar amplitudes before taking the norm; it is not a two-endpoint spring. Dividing it into independent pairwise energies would discard the cross terms responsible for the example.

This is an architectural example, not a zeta model. Its positivity is already present before the grading is introduced. A useful supersymmetric completion of the arithmetic form would need a principle that forces the analogous coherent couplings and their normalization.

7.2 The pole term and a growing Green function

The identity

$$R(t) = e^{t/2} - \sum_{k \geq 1} e^{-(2k+1/2)t} \quad (7.4)$$

suggests replacing one decaying massive propagator by a growing one. The following elementary computation identifies the boundary cost of this particular attempt.

Proposition 7.1 (Growing propagator boundary defect). *For $a > 0$, the kernel*

$$G_a(x, y) = -\frac{1}{2a} e^{a|x-y|}, \quad 0 < x, y < L, \quad (7.5)$$

is the Green function of $-\partial_x^2 + a^2$ with boundary conditions $u'(0) = -au(0)$ and $u'(L) = au(L)$. Its quadratic form is not nonnegative.

Proof. The derivative jump at $x = y$ and the two endpoint derivatives verify the Green equation and Robin conditions. The homogeneous equation has no nonzero solution satisfying both conditions, so the inverse is well defined. Integration by parts gives the form on $H^1(0, L)$,

$$\begin{aligned} \mathcal{E}_R[u] &= \int_0^L (|u'|^2 + a^2|u|^2) dx - a|u(0)|^2 - a|u(L)|^2 \\ &= \|u' - au\|^2 - 2a|u(0)|^2. \end{aligned} \quad (7.6)$$

The form-domain test $u(x) = e^{ax}$ gives $\mathcal{E}_R[u] = -2a < 0$. $\square$

This calculation excludes treating the replacement channel as independently positive. It does not exclude stabilization through a larger coherent system. It also illustrates why a formal differential square without its boundary domain is insufficient.

7.3 The first prime forces joint diagonal accounting

Translate the interval to $(0, L)$ and assume $\log 2 < L \leq \log 3$. Let $a = \log 2$, $c = \log 2/\sqrt{2}$, and $T = T_a$. Since $L < 2a$, $T^2 = 0$. Its initial and final projections are

$$P = T^*T = \mathbf{1}_{(0, L-a)}, \quad F = TT^* = \mathbf{1}_{(a, L)}. \quad (7.7)$$

The first-prime correction is $-c(T + T^*)$. On each paired component it has eigenvalues $\pm c$, so it cannot be a separate positive additive channel.

The explicit edge factor $B_p = \sqrt{c}(P - T^*)$ satisfies

$$B_p^* B_p = c(P + F - T - T^*). \quad (7.8)$$

Hence the exact identity is

$$Q_{0,L}[f] = \|B_p f\|^2 + \left(Q_{0,L}^\gamma[f] - c\langle f, (P + F)f \rangle \right). \quad (7.9)$$

Positivity of the remainder is not proved. At positive shift the arithmetic coefficient becomes $c \cosh(\omega \log 2)$; the same accounting applies to that correction.

There is a complementary restriction on an auxiliary-field approach. For a quadratic block with a fixed bare boundary block A_0 and positive invertible auxiliary block C_0 , minimization gives

$$\begin{pmatrix} A_0 & B_0^* \\ B_0 & C_0 \end{pmatrix} \longmapsto A_0 - B_0^* C_0^{-1} B_0. \quad (7.10)$$

The change is negative semidefinite, whereas the first-prime correction is indefinite. Thus adding an auxiliary Gaussian field while leaving the bare boundary block fixed cannot reproduce that correction either. Equations (7.9) and (7.10) require a successful model to change the joint couplings and diagonal terms together.

8 The input space of a global norm identity

Let Q_W denote the full central Weil form on $C_c^\infty(\mathbb{R})$ with no fixed support bound. The following restriction does not concern the finite-interval closed forms.

Theorem 8.1 (No closable factor on ordinary whole-line L^2). *There is no closable linear map $A : L^2(\mathbb{R}) \supset \text{Dom } A \rightarrow \mathcal{K}$ to a positive Hilbert space, with $C_c^\infty(\mathbb{R}) \subset \text{Dom } A$, such that*

$$\|Af\|_{\mathcal{K}}^2 = Q_W[f] \quad \text{for every } f \in C_c^\infty(\mathbb{R}). \quad (8.1)$$

Proof. Such an identity implies Weil positivity and hence RH; we use the standard Weil criterion as presented in [6, 7]. Under RH the explicit formula becomes

$$Q_W[f] = \sum_{\gamma} m_{\gamma} |\hat{f}(\gamma)|^2, \quad (8.2)$$

where distinct zero ordinates are indexed by γ and m_{γ} is their multiplicity. Fix an ordinate γ_0 and choose $\phi \in C_c^\infty(\mathbb{R})$ with $\hat{\phi}(0) = 1$. Set

$$f_N(x) = N^{-1} e^{i\gamma_0 x} \phi(x/N). \quad (8.3)$$

Then $\|f_N\|^2 = N^{-1} \|\phi\|^2 \rightarrow 0$, while $\hat{f}_N(\gamma) = \hat{\phi}(N(\gamma - \gamma_0))$. These sampling vectors converge in $\ell^2(m_{\gamma})$ to the vector equal to one at γ_0 and zero elsewhere. To justify norm convergence, use isolation of γ_0 , rapid Fourier decay, and the zero-counting estimate $N_{\zeta}(T) = O(T \log T)$ to obtain a summable dominating sequence, uniform for $N \geq 1$.

Consequently $Q_W[f_N - f_M] \rightarrow 0$, but $Q_W[f_N] \rightarrow m_{\gamma_0} > 0$. The assumed norm identity makes Af_N a Cauchy sequence with nonzero limiting norm while $f_N \rightarrow 0$. This contradicts closability. $\square$

The argument is an unconditional impossibility proof: assuming the factor first supplies RH and then yields a contradiction. It neither assumes RH as a result of this investigation nor challenges the possibility of norm identities on other input spaces. Related measure-theoretic closability questions occur in Wiener–Hopf form theory [8]; the proof here is direct for the present sampling measure.

Corollary 8.2. *A bounded embedding $J : L^2(\mathbb{R}) \rightarrow \mathcal{H}_B$ followed by a closed supercharge component S cannot give such a global factor if $C_c^\infty(\mathbb{R}) \subset \text{Dom}(SJ)$.*

Proof. The composition SJ is closed: if $f_n \rightarrow f$ and $SJf_n \rightarrow g$, then $Jf_n \rightarrow Jf$, and closedness of S applies. Theorem 8.1 therefore excludes the claimed norm identity. $\square$

There is no contradiction with the kinetic tower, which realizes a different form with a continuous Fourier density. There is also no contradiction with finite-interval factors: the supports of f_N grow without bound. Test-function topologies, other independently defined positive spaces, and unbounded boundary observables remain possible. Adding a fixed finite number of ordinary Sobolev derivatives to the whole-line norm does not remove this particular sequence: its derivative norms still tend to zero.

9 What a useful supersymmetric completion must establish

9.1 The boundary object must be specified

Three possible constructions need different proofs. A direct factor requires $Q[f] = \|Af\|^2$ on its specified input space. A minimized bulk energy requires an identity with an infimum over auxiliary fields, including the correct endpoint conditions. A spectral response requires identification of a resolvent or covariance with the zeta function of interest. Positivity of a Hamiltonian does not identify these objects with one another.

For example, $H \geq 0$ and a Hilbert-space vector b imply

$$\langle b, (H - z)^{-1}b \rangle = \int_0^\infty \frac{d\mu_b(\lambda)}{\lambda - z}, \quad d\mu_b \geq 0.$$

This is a useful model of a positive response, but an actual boundary trace can be unbounded and can require a different formulation. To use such a mechanism for zeta, the exact response and normalization must be derived from independently specified dynamics.

The shifted transfer family has the form

$$\Theta_\omega(z) = \frac{\xi(1/2 - \omega - iz)}{\xi(1/2 + \omega - iz)}, \quad \xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s). \quad (9.1)$$

Increasing support length adds logarithmic arithmetic delays; changing ω changes the shifted family. Neither operation is automatically the extra coordinate of an extension problem. The RG and bulk-boundary language remains motivation until a theorem identifies the corresponding dynamics and observables.

9.2 Two concrete milestones

The first milestone is a coherent positive model reproducing the complete prime-free form throughout the known positive slab, with the fixed normalization and pole terms in (5.10). The kinetic tower is a controlled ingredient, and the comparison theorem excludes a broad family of simple completions. A proposed model should be tested against both its off-diagonal kernel and its diagonal terms.

The second milestone is the first prime. The joint model must derive the indefinite correction and the diagonal accounting in (7.9); appending an independent positive square or using (7.10) with the original boundary block fixed is insufficient. Beyond the first prime, overlapping delays impose further compatibility conditions not investigated here.

An arithmetic or geometric rule forcing coherent mixing would be substantive progress. A Ward identity could contribute if the model, domain, and invariant positive inner product were independently defined and the identity forced the required equality. At present, no such rule has been found. A fitted positive matrix, a spectral square root of the target, or an inverse realization conditioned on positivity does not provide this missing explanation.

9.3 What should be retained if this route stops

The earlier higher-dimensional investigations warn that additional coordinates can reproduce existing information without strengthening it. The present calculations refine that lesson. Auxiliary dynamics can change the mathematical class of a boundary operator, as demonstrated by the gamma tower. Coherent mixing can change the class of positive boundary couplings, as demonstrated by (7.1). Both are real possibilities, but neither selects the arithmetic data needed for RH.

The obstructions should be used at their stated scope. A failure of pairwise squares motivates testing coherent amplitudes; it does not establish a successful coherent completion. A domain obstruction requires a different boundary topology; merely renaming the original L^2 space as superspace does not change it. These are design constraints for future explorations rather than evidence that a global proof is near.

A Exact rational certificate for the comparison test

The sign used in Theorem 6.3 is verified with rational interval arithmetic in the companion file `check_round2.py`. This appendix states the finite calculation and all truncation bounds.

Let $L = 2/3$, $b = 1/3$, $N = 32$, and $a_k = 2k + 1/2$. Define

$$J(\lambda) = \int_b^L (L - t)e^{-\lambda t} dt = \frac{e^{-\lambda b}[\lambda(L - b) - 1] + e^{-\lambda L}}{\lambda^2}. \quad (\text{A.1})$$

Using (7.4),

$$\int_b^L (L - t)R(t)dt = J(-1/2) - \sum_{k=1}^{\infty} J(a_k). \quad (\text{A.2})$$

The required infinite sums are bounded by

$$\frac{1}{2a_N} \leq \sum_{k=N}^{\infty} a_k^{-2} \leq \frac{1}{2a_N} + a_N^{-2}, \quad (\text{A.3})$$

$$0 \leq \sum_{k=N}^{\infty} \frac{e^{-a_k L}}{a_k^2} \leq \frac{e^{-a_N L}}{a_N^2(1 - e^{-2L})}, \quad (\text{A.4})$$

$$0 \leq \sum_{k=N}^{\infty} J(a_k) \leq \frac{(L - b)e^{-a_N b}}{a_N(1 - e^{-2b})}. \quad (\text{A.5})$$

Equation (A.3) is the integral bound for a decreasing inverse-square function. Equation (A.4) follows by bounding all denominators below by a_N^2 and summing the geometric numerator. For (A.5), bound $L - t \leq L - b$, extend the exponential integral to infinity, and then sum its geometric majorant.

For rational $x \geq 0$, the checker encloses e^x by its Taylor polynomial through degree $m = 160$. If $x < m + 2$, the remaining positive tail is at most

$$\frac{x^{m+1}}{(m+1)!} \frac{1}{1 - x/(m+2)}. \quad (\text{A.6})$$

Negative arguments are handled by reciprocal intervals. All arguments in this calculation obey the required bound. For rational $x \geq 1$, put $z = (x - 1)/(x + 1)$. Then

$$\log x = 2 \sum_{k=0}^{n-1} \frac{z^{2k+1}}{2k+1} + r_n, \quad 0 \leq r_n \leq \frac{2z^{2n+1}}{(2n+1)(1 - z^2)}, \quad (\text{A.7})$$

with $n = 48$. The alternating arctangent series, with 24 terms, and Machin's formula $\pi = 16 \arctan(1/5) - 4 \arctan(1/239)$ give rational endpoints for π . The logarithm endpoints are then evaluated by monotonicity. Finally, the elementary integral-test bounds

$$H_{128} - \log 128 - 1/128 < \gamma_E < H_{128} - \log 128, \quad \log 128 = 7 \log 2, \quad (\text{A.8})$$

are sufficient; no high-precision tabulated value of γ_E is used.

Combining these intervals with (6.3) and (6.5) produces an enclosure for $\mathcal{U}$ with approximate endpoints

$$[-0.0494803686403, -0.0409467581007]. \quad (\text{A.9})$$

The actual comparison in the checker is the exact rational assertion

$$\mathcal{U}_{\text{upper}} < -1/25. \quad (\text{A.10})$$

The displayed decimals are for interpretation only. All additions, multiplications, divisions, and interval endpoint comparisons in this certificate use exact fractions. This is a reproducible computational proof of a sign in the stated normalization, subject to review of the short checker and the elementary bounds above.

B Verification, provenance, and remaining review

The supplied `checks/` directory contains the two original diagnostic programs and their recorded outputs. They use Python and NumPy and require no zeta-zero data. The second program's rational certificate uses only standard-library fraction arithmetic. Its other calculations, and all calculations in the first program, are separate floating-point diagnostics.

The first program checks the digamma/jump normalization, the full-output difference identity on complex polynomial inputs at two quadrature orders, and the first-prime projection algebra. The second checks the mass expansion at six frequencies against an independent digamma evaluation, the auxiliary minimization, the coherent model and its graded algebra, and the constant-input integrals at two quadrature orders. The observed identity discrepancies are consistent with the stated numerical truncations and floating-point precision. These diagnostics do not certify an extension of the small-window theorem.

The normalization was taken from the project manuscripts [1, 2] at repository commit `c6eb19cc47b972b410b077be5af4e86a112afd95`. The researcher's previously reported Fourier checks were accepted as context; their large archive outputs were not required for the present derivations. The data archive is named `szp-archive`. No source repository, synced project document, or archive file was modified in preparing this draft.

The introductory history is based on the researcher's account and the project brief. The mathematical results consolidate two exploratory notes dated 11 September 2026. AI assistance was used in their derivation, checking, and exposition, and in preparing this manuscript. No external author was contacted and no independent review is implied.

Before this could be treated as a submission draft, the priority review should compare the small-window argument with existing Dirichlet-form methods and the pairwise restriction with established comparison-matrix and factor-width results. The domains of the infinite tower and the continuum pairwise definition also merit specialist review. The present paper makes no general continuum Schur-limit theorem and no claim to have constructed an interacting superspace theory.

One formula-convention question concerning negative time in a published norm-map construction was recorded during the first investigation. It is not needed for any proof here and is retained in the companion research notes for further checking, rather than presented as an established correction to that work.

References

- [1] E. Baker, *Archimedean first-slab positivity for shifted zeta canonical systems*, project working manuscript, 2026. Repository source: `papers/first-slab-positivity/first_slab_positivity.tex`. The source file, rather than a public publication, is the cited object.
- [2] E. Baker, *Finite-horizon Weil coercivity and shifted-zeta contraction*, working draft, version 0.4, 10 September 2026. Repository source: `papers/weil-depth/manuscript/finite_horizon_weil.tex`.
- [3] NIST Digital Library of Mathematical Functions, Chapter 5, especially §§5.7, 5.9, and 5.11. <https://dlmf.nist.gov/5>.
- [4] M. Kwaśnicki and J. Mucha, *Extension technique for complete Bernstein functions of the Laplace operator*, Journal of Evolution Equations (2018). [doi:10.1007/s00028-018-0444-4](https://doi.org/10.1007/s00028-018-0444-4). [arXiv:1707.02475](https://arxiv.org/abs/1707.02475).
- [5] A. Singleton, *Superconformal quantum mechanics and the exterior algebra*, 2014. [arXiv:1403.4933](https://arxiv.org/abs/1403.4933).
- [6] M. Suzuki, *On the Hilbert space derived from the Weil distribution*, 2025. [doi:10.4153/S0008414X25101739](https://doi.org/10.4153/S0008414X25101739). [arXiv:2301.00421v3](https://arxiv.org/abs/2301.00421v3).
- [7] M. Suzuki, *Weil’s quadratic form via the screw function*, 2026. [arXiv:2606.09096v1](https://arxiv.org/abs/2606.09096v1). This is the version consulted during the exploratory investigation; a later revision exists and remains to be compared in the priority audit.
- [8] D. R. Yafaev, *On semibounded Wiener–Hopf operators*, 2016. [arXiv:1606.01361](https://arxiv.org/abs/1606.01361).