

Finite response matrices for Weil positivity

Draft for author review

13 September 2026

Abstract

We derive an explicit boundary representation and controlled finite response for the localized Weil form. A digamma resolvent expansion and a Neumann comparison give an unconditional positive lower bound on a high-frequency sector. Its exact elimination leaves a finite Hermitian matrix, retaining every coupling to the low modes. The boundary correction is bounded and noncompact, but has convergent tails on finite input spaces. These estimates yield lower and upper response enclosures and a matrix conjecture equivalent to the Riemann hypothesis, also expressible with arbitrarily small positive tolerances. The unresolved conjecture is positivity of these matrices on an unbounded sequence of intervals. The equivalence and the analytic reduction do not establish that positivity. No priority is claimed for the general finite-codimension or Schur-complement principles.

1 Boundary response and the matrix conjecture

The localized gamma operator differs from its diagonal Neumann comparison by an explicit boundary correction. Controlling that correction on finite input spaces gives certified bounds for the complete low/high response, and isolates all possible negativity in a finite matrix. These are the analytic results underlying the formulation developed from [6]; the arithmetic sign of the matrix remains open.

The ingredients below define a matrix S_n for every integer $n \geq 1$ without using zeta zeros or assuming their location. Its dimension is a prescribed integer $N(n)$ selected by the terminating certification procedure in Section 2.

Conjecture 1.1 (Finite response). For every positive integer n , the matrix S_n defined in (4.1) is positive semidefinite.

In energy language, allowing the positive high modes to respond to a low-mode input never removes more than the low input's original energy. Theorem 4.1 proves that this assertion is equivalent to RH.

Localized Weil positivity, finite-codimension positivity and variational formulations have substantial antecedents; see [1, Section 1]. Connes–Consani–Moscovici give a localized form core, explicit matrices and finite restrictions, whose smallest eigenvalues recover the localized lower bound [2, Proposition 3.4 and Sections 4–5]. Here the finite matrix retains the exact response of the positive infinite sector, with enclosures for that response. The particular boundary and remainder formulas are the focus; the localization and Schur principles are standard. No general priority claim is made.

2 The arithmetic operator and an unconditional cutoff

Use $X_L = L^2(0, L)$, zero extension E_L , and $\widehat{F}(\tau) = \int_{\mathbb{R}} F(x) e^{-i\tau x} dx$. Inner products are antilinear in the first slot. Set

$$a_k = 2k + \frac{1}{2}, \quad b(s) = \sum_{k=0}^{\infty} \frac{2}{a_k} \frac{s}{a_k^2 + s} = \operatorname{Re} \psi\left(\frac{1}{4} + i\sqrt{s}/2\right) - \psi\left(\frac{1}{4}\right), \quad (2.1)$$

$$w_0 = \psi\left(\frac{1}{4}\right) - \log \pi, \quad \mathcal{D}_L = \left\{ f \in X_L : \int_{\mathbb{R}} \log(2 + |\tau|) |\widehat{E_L f}(\tau)|^2 d\tau < \infty \right\}. \quad (2.2)$$

Let T_L be the nonnegative operator associated with the closed form

$$t_L[f] = \frac{1}{2\pi} \int_{\mathbb{R}} b(\tau^2) |\widehat{E_L f}(\tau)|^2 d\tau.$$

For $d > 0$, $U_d F(x) = F(x - d)$ and $V_d = E_L^* U_d E_L$. Put $c_L(x) = \cosh((x - L/2)/2)$ and $s_L(x) = \sinh((x - L/2)/2)$. The bounded remainder and the full operator are

$$R_L = w_0 I + 2c_L c_L^* - 2s_L s_L^* - \sum_{\substack{p \text{ prime, } m \geq 1 \\ m \log p < L}} (\log p) p^{-m/2} (V_{m \log p} + V_{m \log p}^*), \quad (2.3)$$

$$W_L = T_L + R_L, \quad Q_L[f] = t_L[f] + \langle f, R_L f \rangle. \quad (2.4)$$

Here $vv^* f = v \langle v, f \rangle$. The sum includes every active prime power. All sums in R_L are finite. In the additive normalization of the Weil functional in [1, Section 1.1], Q_L is exactly the Weil form of $E_L f$. The normalization and its spatial-to-Fourier bridge are detailed in [7, Sections 2–3]. The constant, both poles and all prime powers are essential.

The digamma expansion [3] gives $b(\tau^2) = \log |\tau| - \log 2 - \psi(1/4) + O(\tau^{-2})$. Consequently $\mathcal{D}_L$ is the form domain and $C_c^\infty(0, L)$ is a form core. The support contraction followed by mollification proves the core assertion in this logarithmic Fourier norm [7, Section 3]. Bounded perturbation then defines the self-adjoint, semibounded W_L on $\operatorname{Dom} T_L$.

Let H_N and H_D denote the Neumann and Dirichlet interval Laplacians. Minimizing

$$\frac{2}{a} \left(\|E_L f - u\|_{L^2(\mathbb{R})}^2 + a^{-2} \|u'\|_{L^2(\mathbb{R})}^2 \right)$$

over $u \in H^1(\mathbb{R})$ gives $(2/a)s/(a^2 + s)$ in Fourier space. Restricting u to the interval lowers this minimum; allowing only zero extensions of $H_0^1(0, L)$ competitors raises it. Summation proves the closed-form bracket

$$b(H_N) \leq T_L \leq b(H_D). \quad (2.5)$$

The positive resolvent representation is a concrete case of the extension framework in [4]. The comparison does not assign Neumann boundary conditions to the actual whole-line fields.

Write

$$e_j(x) = \nu_j \cos(j\pi x/L), \quad \nu_0 = L^{-1/2}, \quad \nu_j = \sqrt{2/L} \ (j > 0), \quad b_j = b((j\pi/L)^2).$$

Let $P_h = \sum_{j=0}^{h-1} e_j e_j^*$. For $q \in (0, 1)$ define

$$r_h(q) = \frac{q}{1-q} \left(2 - q^{\lfloor (h-1)/2 \rfloor} - q^{\lceil (h-1)/2 \rceil} \right),$$

and specify

$$\beta_L = -w_0 + 2 \sinh(L/2) - L + \sum_{\log p < L} (\log p) r_{\lceil L/\log p \rceil} (p^{-1/2}). \quad (2.6)$$

The prime bound follows by decomposing translation by $\log p$ into finite chains and bounding each symmetric correlation matrix by its largest row sum r_h : successive row sums differ by $q^i - q^{h-i}$, so the maximum occurs centrally. The odd pole costs at most $2\|s_L\|^2 = 2 \sinh(L/2) - L$; the even pole is nonnegative. Thus

$$W_L \geq b(H_N) - \beta_L I. \quad (2.7)$$

Fix $\delta_0 = 1/16$. For any $L > 0$, choose a cutoff satisfying

$$b_{N(L)} > \beta_L + \delta_0, \quad P = P_{N(L)}, \quad Q = I - P. \quad (2.8)$$

For the integer family use this certified convention. Fix deterministic rational enclosure routines as in [7, Section 4]. At stage $r = 1, 2, \dots$, enclose β_n and b_j , $1 \leq j \leq r$, in intervals of width at most 2^{-r} . Stop at the first stage with $\underline{b}_{j,r} > \overline{\beta}_{n,r} + \delta_0$, choosing the least certified j at that stage as $N(n)$. The search terminates because $b_j \rightarrow \infty$; it never waits to resolve an equality at a smaller index. Fixing the routines fixes the family. Different admissible choices can change its matrices, but all the equivalences below hold for each choice.

Let H be the self-adjoint operator associated with the high form Q_L restricted to $Q\mathcal{D}_L$. Then $H \geq \delta_0 I$ unconditionally, so H^{-1} is well defined with norm at most δ_0^{-1} . No invariance of QX_L under W_L is assumed.

3 An explicit boundary operator

The following representation makes that coupling accessible. For $a > 0$ set

$$u_a(x) = e^{-ax}, \quad v_a(x) = e^{-a(L-x)}, \quad r_a = e^{-aL}, \quad K_a = \frac{u_a u_a^* + v_a v_a^* + r_a(u_a v_a^* + v_a u_a^*)}{1 - r_a^2}.$$

Theorem 3.1 (Boundary representation). *The positive finite sums $K^{(J)} = \sum_{k < J} K_{a_k}$ converge strongly to a bounded positive operator K_L satisfying*

$$T_L = b(H_N) + K_L, \quad \text{Dom } T_L = \text{Dom } b(H_N), \quad \mathcal{D}_L = \text{Dom } b(H_N)^{1/2}. \quad (3.1)$$

Finite cosine sums are an operator core. The essential norm is $\|K_L\|_{\text{ess}} = \pi/2$, so finite mass sums do not converge in operator norm. Nevertheless, for the first h cosine modes,

$$\|(K_L - K^{(J)})P_h\| \leq \eta_{h,J} := \sqrt{\frac{2(2h-1)}{L}} (a_J^{-1/2} + a_J^{-3/2}) \rightarrow 0. \quad (3.2)$$

Proof. The compressed free Green function is $e^{-a|x-y|}/(2a)$; the Neumann Green function is

$$\frac{\cosh(a \min(x, y)) \cosh(a(L - \max(x, y)))}{a \sinh(aL)}.$$

Their difference multiplied by $2a$ is K_a . Subtracting the corresponding resolvent energies proves the finite-tower version of (3.1). The coefficient matrix of K_a is positive.

Let $n(t) = e^{-t/2}/(1 - e^{-2t})$. Summing the kernels gives $n(x+y) + n(2L-x-y)$ plus a bounded kernel. Since $n(t) = (2t)^{-1} + O_L(1)$ on $(0, 2L]$, this is one half of each endpoint Carleman kernel,

modulo a bounded-kernel, hence compact, operator. The weighted Schur test bounds the Carleman operator by π . The partial towers have the same uniform kernel bound, so monotone convergence yields a bounded strong limit and the asserted domains.

Splitting the interval in half makes the two endpoint operators a direct sum modulo compact operators. Each has essential norm π before the factor $1/2$: normalized functions proportional to $t^{-1/2}$ on $[1, R]$, then dilated toward zero, have limiting Rayleigh quotient π and tend weakly to zero. This also proves the lower essential-norm bound; see the Carleman background in [5].

Direct integration, with $\omega_j = j\pi/L$, gives

$$K_a e_j = \frac{\nu_j a}{a^2 + \omega_j^2} (u_a + (-1)^j v_a). \quad (3.3)$$

Its norm is at most $\sqrt{2}\nu_j a^{-3/2}$. Sum the decreasing series using $\sum_{k \geq J} a_k^{-3/2} \leq a_J^{-3/2} + a_J^{-1/2}$, then sum squared column bounds over $j < h$ to obtain (3.2). $\square$

In particular all low basis vectors lie in $\text{Dom } W_L$. The boundary entries vanish for opposite parities; for equal parities they are

$$\langle e_i, K_L e_j \rangle = 2\nu_i \nu_j \sum_{k \geq 0} \frac{a_k^2 (1 - (-1)^j e^{-a_k L})}{(a_k^2 + \omega_i^2)(a_k^2 + \omega_j^2)}. \quad (3.4)$$

The remaining entries are elementary integrals of the rank-two pole kernel and of translated cosines on their overlap intervals. These formulas specify the operator using gamma and prime data, independently of RH. For a fixed finite compression the entry series has a sharper $O(J^{-1})$ mass tail; the full-column tail (3.2) remains necessary for residual norms [7, Section 6].

4 Exact elimination and the equivalence theorem

For each L , use the prescribed P, Q and define

$$A = PW_L P, \quad B = QW_L P, \quad H = QW_L Q, \quad \text{Dom } H = Q \text{Dom } b(H_N).$$

Here A is an $N(L)$ by $N(L)$ Hermitian matrix, $B : PX_L \rightarrow QX_L$ is bounded, and H is the operator associated with the high form above; the displayed compression and domain follow from (3.1). Define the finite matrix

$$S_L = A - B^* H^{-1} B, \quad S_n := S_L|_{L=n}. \quad (4.1)$$

Its entries involve an infinite-dimensional inverse. It is a finite matrix with explicitly specified, approximable entries, not a formula involving only finitely many elementary operations.

Theorem 4.1 (Exact RH equivalence). *For each $L > 0$, $Q_L \geq 0$ on $\mathcal{D}_L$ if and only if $S_L \succeq 0$. Consequently Conjecture 1.1 is equivalent to RH.*

Proof. Write $f = p + h$, $p = Pf$, $h = Qf$. Completing the high square gives the identity of closed forms

$$Q_L[p + h] = \|H^{1/2}(h + H^{-1}Bp)\|^2 + \langle p, S_L p \rangle. \quad (4.2)$$

Every term is defined: $p \in \text{Dom } W_L$, $h \in \text{Dom } H^{1/2}$, and $H^{-1}Bp \in \text{Dom } H$. If $S_L \succeq 0$, this proves positivity. Conversely, choose $h = -H^{-1}Bp$ to recover exactly $\langle p, S_L p \rangle$.

RH implies Weil positivity on smooth compactly supported functions. The form core extends it to $\mathcal{D}_L$, so every $S_n \succeq 0$. For the converse, every compact support can be translated into $(0, n)$ for

some integer n . The Weil form is translation invariant: its gamma multiplier and prime correlations are invariant, and the pole kernel is $2 \cosh((x - y)/2)$. Zero extension to a larger interval preserves all these terms. Thus positivity of all S_n implies Weil positivity on all required test functions, and hence RH by the classical criterion [1]. $\square$

No uniform gap in n , common source on all intervals, or compatible limit of the matrices is required. The dimensions can grow without bound.

Reflection $\mathcal{J}f(x) = f(L - x)$ preserves the form and the cosine cutoff: $\mathcal{J}e_j = (-1)^j e_j$ and $\mathcal{J}V_d \mathcal{J} = V_d^*$. Thus H^{-1} and the response respect parity, and $S_L = S_L^{\text{even}} \oplus S_L^{\text{odd}}$. Both blocks must be nonnegative; their dimensions are $\lceil N/2 \rceil$ and $\lfloor N/2 \rfloor$. Trials respecting parity retain this split in the enclosures below [7, Sections 6–7].

5 Bounds on the unresolved matrix

Put $d_j = b_j - \beta_L$ for $j \geq N$, and let $\Lambda e_j = d_j e_j$ on QX_L . Then $H \geq \Lambda \geq \delta_0 I$. For any finite-input trial map $Y : PX_L \rightarrow \text{Dom } H$, set $C(Y) = (P - Y)^* W_L (P - Y)$ and $R_Y = B - HY$. Expansion gives the variational identity

$$S_L = C(Y) - R_Y^* H^{-1} R_Y, \quad 0 \preceq R_Y^* H^{-1} R_Y \preceq R_Y^* \Lambda^{-1} R_Y. \quad (5.1)$$

It explains why the energy of a trial is an upper bound until its remaining response is subtracted. For $K > N$, let $P_{N,K}$ project onto $e_N, \dots, e_{K-1}$ and define the Galerkin trial by the positive finite system

$$(P_{N,K} H P_{N,K}) Y_K = P_{N,K} B, \quad \text{ran } Y_K \subset P_{N,K} X_L.$$

Its residual $R_K = B - H Y_K$ is supported in cosine modes $j \geq K$.

Theorem 5.1 (Convergent enclosures). *The finite Hermitian matrices*

$$L_{L,K} = A - B^* Y_K - d_K^{-1} R_K^* R_K, \quad U_{L,K} = A - B^* Y_K \quad (5.2)$$

satisfy $L_{L,K} \preceq S_L \preceq U_{L,K}$ and converge to S_L in norm as $K \rightarrow \infty$. Their gap is exactly $d_K^{-1} R_K^* R_K$.

Proof. Galerkin orthogonality gives $Y_K^* H Y_K = B^* Y_K = Y_K^* B$ and $P_{N,K} R_K = 0$. Apply (5.1) and $\Lambda^{-1} \leq d_K^{-1} I$ on the remaining modes to obtain the bounds. Write $H = \Lambda + E$, where $E = Q(K_L + R_L + \beta_L I)Q$ is bounded. Coercivity gives $\|Y_K\| \leq \delta_0^{-1} \|B\|$, while $R_K = (I - P_K)(B - E Y_K)$. Consequently

$$\|U_{L,K} - L_{L,K}\| \leq \frac{\|B\|^2 (1 + \|E\|/\delta_0)^2}{d_K} \rightarrow 0.$$

Each enclosure's distance from S_L is at most this gap. Since $d_K = \log K + O_L(1)$, this is a crude $O_L(1/\log K)$ bound. $\square$

The column tail (3.2) controls finite mass approximations to B and $H Y_K$. The residual Gram must include their interference with the pole and prime columns. Positive eigenvalues of an upper enclosure alone are not a positivity certificate.

Corollary 5.2 (An equivalent tolerance formulation). *Conjecture 1.1, and hence RH, is equivalent to*

$$\forall n, m \in \mathbb{N} \exists K > N(n) : \quad L_{n,K} + 2^{-m} I \succ 0. \quad (5.3)$$

Proof. If $S_n \succeq 0$, choose K with $\|L_{n,K} - S_n\| < 2^{-m}$, using Theorem 5.1. Conversely $S_n \succeq L_{n,K} \succ -2^{-m}I$ for every m implies $S_n \succeq 0$. This avoids assuming that a semidefinite limit is detected by an unshifted, finite strict certificate. $\square$

The strict margin makes each true instance amenable to sufficiently accurate enclosures. Arbitrary finite trials and explicit column-error allowances are treated in [7, Section 10]; its Section 9 proves the stronger residual convergence needed there. An optional equivalent comparison of two positive matrices is given in its Section 8. No finite collection of successful intervals or tolerances establishes (5.3).

6 What makes the remaining conjecture arithmetic

The reduction works for many semibounded operators. For example $W = \begin{pmatrix} 1/2 & 1 \\ 1 & 1 \end{pmatrix}$ has a positive high block $H = 1$ and $W + I > 0$, but its low Schur complement is $-1/2$. A positive shifted source and a positive high sector alone therefore do not imply Conjecture 1.1.

There is a more relevant negative control. Define the Chebyshev function $\psi_{\text{Ch}}(x) = \sum_{p^m \leq x} \log p$ and $E(r) = \psi_{\text{Ch}}(e^r) - e^r + 1$. For $h_f(r) = \text{Re}\langle E_L f, U_r E_L f \rangle$, define the gamma form with its lowest mass removed by

$$t_{\geq 1, L}[f] = \frac{1}{2\pi} \int_{\mathbb{R}} \sum_{k \geq 1} \frac{2}{a_k} \frac{\tau^2}{a_k^2 + \tau^2} |\widehat{E_L f}(\tau)|^2 d\tau.$$

Cancellation of the smooth prime density against the poles and the lowest gamma mass gives

$$Q_L[f] = t_{\geq 1, L}[f] + (4 + w_0)\|f\|^2 - 2 \int_0^L e^{-r/2} h_f(r) dE(r). \quad (6.1)$$

The Stieltjes form holds on $\mathcal{D}_L$; derivatives of h_f are not required. If the prime measure is replaced by $e^r dr$, the last term vanishes. On the unit input $f(x) = \sin(\pi x/2)$ on $(0, 2)$,

$$Q_2^{\text{dens}}[f] < -1 + \frac{26}{125} \frac{(22/7)^2}{4} = -\frac{2979}{6125} < 0.$$

Here $w_0 < -5$ and $\sum_{k \geq 1} 2/a_k^3 \leq 26/125$; smooth approximation preserves negativity. The actual prime discrepancy carries information that the leading density and the positive response architecture cannot replace.

At $L = 1$, an explicit rational certificate permits the alternative cutoff $N = 4$ with $\delta_0 = 1/16$ [7, Section 13]. This certifies only the high sector, and does not evaluate the response matrix or establish its sign. Present bounds may require very large dimensions as L increases. No practical complexity estimate, new positivity interval, or zero computation is asserted here.

The boundary and response estimates isolate the arithmetic ordering that remains to be proved. Positive auxiliary constructions alone do not settle it; the exact prime contribution remains essential.

Preparation and verification status

This draft was prepared with substantial assistance from OpenAI's Codex in mathematical analysis, derivation checking, code and exposition. It is a draft for human mathematical review; completion of that review is not claimed. The companion [7] expands the analytic arguments and identifies the checks needed to support the equivalence. This revision incorporates the documented Codex review of 13 September 2026. The repository citation records the source revision used here. A version-specific archival DOI has not yet been supplied.

References

- [1] M. Suzuki, *Weil's quadratic form via the screw function*, 2026, [arXiv:2606.09096v2](#).
- [2] A. Connes, C. Consani and H. Moscovici, *Zeta Spectral Triples*, 2025, [arXiv:2511.22755v1](#).
- [3] NIST, *Digital Library of Mathematical Functions*, [Section 5.7](#), [equation \(5.7.6\)](#), [Section 5.11](#), [equation \(5.11.2\)](#), and [Section 5.4](#).
- [4] M. Kwaśnicki and J. Mucha, *Extension technique for complete Bernstein functions of the Laplace operator*, 2017, [arXiv:1707.02475](#).
- [5] D. R. Yafaev, *Spectral and scattering theory for perturbations of the Carleman operator*, 2012, [arXiv:1210.5709v2](#).
- [6] *Arithmetic ground-state geometry: Physical caps, arithmetic sources and analytical sign tests*, working manuscript in the [shifted-zeta-positivity](#) project, September 2026, [repository revision efb908a](#), especially Sections 10, 14 and 15.
- [7] *Detailed derivations for human review: Companion to the finite-response RH formulation*, accompanying derivation supplement, September 2026, [derivations.pdf](#).