RH-Conditional Approximation Estimates and the Remaining Simplicity Problem

Drafted for Edward Baker

21 October 2026

Research consolidation through Notes 17–18

Abstract

We consolidate analytic estimates obtained while investigating whether the Riemann hypothesis implies simplicity of the nontrivial zeros of the zeta function. The principal approximation argument establishes, within the research notes, a complete first-order limit for fixed Gevrey-tapered Möbius polynomials and an RH-conditional infimum constant equal to the sum of squared multiplicities. A completed-ratio energy has an explicit heat-scale transition governed by the same multiplicity excess. Exact finite Dirichlet filters give a sharp local resolution cost. A normalized heat target admits a complete first-order limit at bounded scaled time. We compute the sharp cost of transferring its profiles back to the original arithmetic problem over fixed finite filtered-Gevrey families, and show that specified exact heat completions still fail to retain an ordinary Dirichlet series at fixed positive time. These estimates do not exclude a multiple zero: the missing ingredient is an arithmetic upper estimate that contradicts its additional cost. This brief presents precise statements and proof mechanisms, with the full analytic payments referenced to the research notes.

1 Scope and the unresolved quantity

Unless stated otherwise, RH is an assumption, not a conclusion. Write $\rho = \frac{1}{2} + i\gamma$ for a distinct nontrivial zero, m_ρ for its multiplicity, and sum over both ordinate signs. The two masses are

$$K = \sum_{\rho \text{ distinct}} \frac{m_\rho^2}{|\rho|^2}, \qquad Z = \sum_{\rho \text{ distinct}} \frac{m_\rho}{|\rho|^2} = 2 + \gamma_E - \log(4\pi). \tag{1}$$

They are finite; the usual unit-window zero count also bounds the sum of squared multiplicities with this weight. Their difference is

$$K - Z = \sum_{\rho \text{ distinct}} \frac{m_\rho(m_\rho - 1)}{|\rho|^2}. \tag{2}$$

A double conjugate pair contributes $4/(\gamma^2 + 1/4)$. The approximation and heat estimates below retain this excess. Identifying it is a bookkeeping conclusion, not a new simplicity criterion or a proof that the excess vanishes.

The results inherited from boundary winding justify the local multiplicity calculation and finite-height continuation under RH. They do not furnish a strict surplus with the opposing sign. The present judgment is therefore that the investigation has produced analytic estimates, but does not have a nearly complete simplicity proof.

Status. The new analytic results are proved in the linked research notes and have undergone separate same-model internal derivations and adversarial checks. They have not received specialist external review or analytic formalization. In particular, the complete scaled-tail theorem below is a substantial internal premise, not a previously published theorem being cited under a new name. No novelty or priority claim is made.

2 Fixed Gevrey approximation and complete tails

For a Dirichlet polynomial D, define

$$E(D) = \frac{1}{2\pi}\int_{\mathbb{R}} \frac{|1-\zeta(\frac{1}{2}+it)D(\frac{1}{2}+it)|^2}{1/4+t^2}\,\mathrm{d}t, \qquad d_N^2 = \inf_{D(s)=\sum_{n\le N} a_n n^{-s}} E(D).$$

We also use $\|g\|_{\mathcal{H}}^2 = (2\pi)^{-1}\int_{\mathbb{R}} |g(\frac{1}{2}+iy)|^2(1/4+y^2)^{-1}\,\mathrm{d}y$, so $E(D) = \|1-\zeta D\|_{\mathcal{H}}^2$. Let W be real, smooth, supported in $[-1,1]$, with $W(0)=1$ and

$$\|W^{(j)}\|_1 \le A^{j+1}(j!)^{\sigma}, \qquad 1<\sigma<2.$$

Set $w = W|_{[0,1]}$, $L=\log N$, and

$$V_{N,w}(s) = \sum_{n\le N} \mu(n) n^{-s} w(\log n/L), \qquad p_{m,w}(u) = -\frac{(u^{m-1}w(u))^{(m)}}{(m-1)!}, \qquad C_m(w) = \|p_{m,w}\|_{L^2(0,1)}^2.$$

Theorem 1 (Complete fixed-taper estimate)**.** *Assume RH and fix W as above. If $E_{N,w}^{>T}$ denotes the part of $E(V_{N,w})$ at $|t|>T$, then for every $0<\eta<2-\sigma$,*

$$\sup_{N\ge 2} (\log N) E_{N,w}^{>T} \ll_{W,\eta} T^{\sigma-2+\eta}, \qquad T\ge 2, \tag{3}$$

$$\lim_{N\to\infty} (\log N) E(V_{N,w}) = \sum_{\rho \text{ distinct}} \frac{C_{m_\rho}(w)}{|\rho|^2} < \infty. \tag{4}$$

No simplicity, zero-gap lower bound, or inverse-derivative moment is assumed.

Proof mechanism and dependencies. The full proof is Note 16, Sections 2–6, 8 and 11. For $s=\frac{1}{2}+it$ and $H>0$, its exact starting identity is

$$V_{N,w}(s) = \int_{\mathbb{R}} \frac{K_H(v)}{\zeta(s+(H+iv)/L)}\,\mathrm{d}v, \quad K_H(v) = \frac{1}{2\pi}\int_{\mathbb{R}} W(u) e^{(H+iv)u}\,\mathrm{d}u.$$

Fourier inversion on an absolutely convergent Möbius line, followed by a shift within the RH zero-free half-plane, preserves every coefficient. Integration by parts against the whole complex exponential gives

$$|K_H(v)| \le C_W e^H e^{-c_W|v|^{1/\sigma}}, \qquad \int_{\mathbb{R}} K_H(v)\,\mathrm{d}v = 1,$$

with constants independent of $H>0$. The factor e^H is retained.

The primary height inputs are the RH argument bound of Carneiro–Chandee–Milinovich [2] and the subpower zeta bound of Chandee–Soundararajan [3]. In the low-height range, the proof keeps the whole nearby zero cluster, including repetitions. Its kernel moment has logarithm $(\sigma-1)M\log(M+2)+O_W(M+1)$ for a cluster of size M. This is paid using the half-coefficient

local counting allowance; no individual reciprocal derivative is estimated. A separate moving line handles the high-height range and its complete remote convolution.

Subtracting the constant term coherently gives decay in the actual distance to the nearest zero. Squaring and integrating that decay over nearest-zero cells costs $O(1/L)$ independently of the gaps. This proves (3). On a fixed isolated zero neighborhood, the complete error has profile $\int_0^1 p_{m,w}(u)e^{-iLyu}\,\mathrm{d}u + O_{W,\rho}(L^{-1})$, with constants allowed to depend on that neighborhood. Its unscaled norm is $C_m(w)/(L|\rho|^2) + O(\log L/L^2)$. The separately established tail bound then permits height exhaustion. Thus local asymptotics alone are not being used as a global upper bound.

A controlled moving family. For the almost-linear taper $w_\eta(u) = (1-u)\chi((1-u)/\eta)$, with a fixed Gevrey cutoff $\chi = 0$ on $(-\infty, \frac{1}{2}]$ and $\chi = 1$ on $[1,\infty)$, a separate proof gives

$$E(V_{N,w_\eta}) \le \frac{C\exp(C\eta^{-a})}{L}, \quad L^{-\theta_0} \le \eta < \tfrac{1}{4}, \quad a > \frac{1}{2-\sigma}, \quad 0 < \theta_0 < 1. \tag{5}$$

Here the parameters σ, θ_0, a are fixed (Note 16, Section 12). Consequently $\eta_N = (\log L)^{-\kappa}$, $0 < \kappa < 2-\sigma$, gives $E(V_{N,w_{\eta_N}}) \le L^{-1+o(1)}$. The exponential allowance is compatible with the multiple-zero lower costs. It supplies neither a moving first-order asymptotic nor a multiplicity-free upper constant.

3 The infimum constant and finite Dirichlet filters

Integration by parts gives $\int_0^1 u^j p_{m,w}(u)\,\mathrm{d}u = \delta_{j0}$ for $0 \le j < m$. Projection onto polynomials of degree below m therefore yields

$$C_m(w) = m^2 + \|p_{m,w} - \mathcal{K}_m(\,\cdot\,,0)\|_2^2 \ge m^2, \tag{6}$$

where $\mathcal{K}_m$ is their endpoint reproducing kernel on $[0,1]$. Flat Gevrey approximations to $(1-u)^m$ approach this lower cost. One common taper need not optimize different multiplicities at once.

Theorem 2 (Infimum consequence of Theorem 1)**.** *Assuming RH and the complete fixed-taper estimate,*

$$\lim_{N\to\infty} (\log N) d_N^2 = K. \tag{7}$$

Real coefficients suffice.

Proof outline. Burnol's lower bound [1] gives the lower limit K. For the upper bound, fix a baseline taper and a finite set F of zeros carrying all but a small part of its profile mass. Choose a fixed near-optimal taper separately at each target. Bounded finite Dirichlet multipliers P_ρ select these profiles:

$$D_N = V_{n,w_0} + \sum_{\rho\in F} P_\rho(V_{n,w_\rho} - V_{n,w_0}), \qquad n = \lfloor N/M \rfloor,$$

where all multipliers have length at most M. The multiplier coefficients of the error sum identically to one, so every cross term and moment constraint is retained. For fixed multipliers, the local profile expansion together with Theorem 1's scaled-tail bound gives the complete mixed limit. Its profile tail is bounded by a fixed multiple of the sum of the finitely many original profile tails.

One may interpolate exactly on any larger finite zero set G. For $0 < \theta < \pi$, $k = \lfloor \pi/\theta \rfloor$, the polynomial

$$B_\theta(z) = \frac{\sin\theta - \sin((k+1)\theta)z^k + \sin(k\theta)z^{k+1}}{\sin\theta - \sin((k+1)\theta) + \sin(k\theta)}$$

has nonnegative coefficients summing to one, equals one at 1, and vanishes at $e^{\pm i\theta}$. At $\theta = \pi$, use $(1+z)/2$. Substitute $z = p^{\rho-s}$, choosing $p = 2$ or 3 with nonzero relative phase, and multiply over the other nodes of G. Irrationality of $\log 2/\log 3$ guarantees such a choice. The products remain bounded by one on the critical line. Their complete exterior profile allowance is independent of their degrees and tends to zero as G exhausts the zeros.

Keep F, tapers and finite G fixed before taking $N \to \infty$. Then exhaust G, optimize the finite profile costs, and let the initial baseline-tail tolerance tend to zero. Conjugate pairing makes the final coefficients real. This proves the upper limit K; see Note 16, Sections 13–14. □

This conclusion concerns the unrestricted polynomial infimum, not the original linear Möbius polynomial. It removes a common-taper restriction but leaves the squared multiplicities in (1). It provides no effective threshold in N.

Proposition 3 (Sharp two-point isolation cost)**.** *Let P have Dirichlet length at most M, be bounded by one on $\Re s = \frac{1}{2}$, and satisfy $P(\rho) = 1$ and $|P(\rho + i\Delta)| \le \varepsilon$, $0 \le \varepsilon < 1$. Then*

$$|\Delta| \log M \ge 2 \arccos \varepsilon.$$

For exact cancellation at a small nonzero gap, the least log-length lies between $\pi/|\Delta|$ and $\pi/|\Delta| + \log 2$.

Proof. Centering the frequency interval gives an entire function of type at most $(\log M)/2$. Its real part starts at one, is bounded by one, and ends at most ε. The sharpened Bernstein inequality [5], integrated after arccos, gives the lower bound. The polynomial B_θ with $p = 2$ and $\theta = |\Delta| \log 2$ gives the upper bound. □

This prices root isolation only. Assigning the same taper to two roots need not isolate them, and the proposition is not a lower bound for every approximation scheme.

4 Completed-ratio energy and heat separation

Use the original theta source

$$H(t,x) = \int_0^\infty e^{tu^2} \Phi(u) \cos(xu)\, \mathrm{d}u, \qquad \Phi(u) = \sum_{n \ge 1} (2\pi^2 n^4 e^{9u} - 3\pi n^2 e^{5u}) e^{-\pi n^2 e^{4u}},$$

with $\xi(s) = \frac{1}{2} s(s-1) \pi^{-s/2} \Gamma(s/2) \zeta(s)$. The completion and heat normalization are

$$\mathcal{F}_t(z) = 8H(t, -2iz), \quad \mathcal{F}_0(z) = \xi(\tfrac{1}{2} + z), \quad \partial_t \mathcal{F}_t = \tfrac{1}{4} \partial_z^2 \mathcal{F}_t.$$

For $\alpha > 0$, define

$$J_t(\epsilon) = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{|1 - \mathcal{F}_t(iy)/\mathcal{F}_t(\epsilon + iy)|^2}{\alpha^2 + y^2}\, \mathrm{d}y, \qquad M_\alpha = \sum_\gamma \frac{m_\gamma}{\alpha^2 + \gamma^2}, \quad K_\alpha = \sum_\gamma \frac{m_\gamma^2}{\alpha^2 + \gamma^2}.$$

Under RH, for $t \ge 0$, the paired order-one product makes the ratio contractive. Its positive harmonic majorant has a known complete mass. Local profiles followed by exhaustion of that mass give

$$\lim_{\epsilon \downarrow 0} \frac{J_0(\epsilon)}{\epsilon} = \sum_\gamma \frac{c_{m_\gamma}}{\alpha^2 + \gamma^2}, \qquad c_m = m \left(1 - \frac{\binom{2m}{m}}{4^m} \right). \tag{8}$$

In particular $c_1 = \frac{1}{2}$, $c_2 = \frac{5}{4}$; no mass at infinity is discarded. The proof is Note 15, Sections 1–3.

For $t = v\epsilon^2$, the local heat expansion at an order-m zero gives Hermite centers $r\lambda_j$, where $r = \sqrt{v/2}$ and $\mathrm{He}_m(\lambda_j) = 0$. Define

$$P_{m,r}(y) = \prod_{j=1}^{m}(y - r\lambda_j), \qquad \mathcal{C}_m(r) = \frac{1}{2\pi}\int_{\mathbb{R}}\left|1 - \frac{P_{m,r}(y)}{P_{m,r}(y-i)}\right|^2 \mathrm{d}y.$$

The same complete-mass argument proves, at each fixed $v \geq 0$,

$$\lim_{\epsilon\downarrow 0}\frac{J_{v\epsilon^2}(\epsilon)}{\epsilon} = \mathcal{A}_\alpha(v) := \sum_{\gamma}\frac{\mathcal{C}_{m_\gamma}(\sqrt{v/2})}{\alpha^2+\gamma^2}.$$

The analytic transition is quantified by

$$\mathcal{C}_2(r) = 1 + \frac{1}{4(1+r^2)}, \tag{9}$$

$$0 \leq \mathcal{A}_\alpha(v) - \frac{M_\alpha}{2} \leq \frac{K_\alpha - M_\alpha}{4v}, \qquad v > 0, \tag{10}$$

$$\lim_{v\to\infty} v\left(\mathcal{A}_\alpha(v) - \frac{M_\alpha}{2}\right) = \frac{K_\alpha - M_\alpha}{4}. \tag{11}$$

The estimate follows from pairwise overlap integrals of the local contractive factors and $\sum_{i<j}(\lambda_i - \lambda_j)^{-2} = m(m-1)/8$. The weighted multiplicity-square sum dominates the passage to all zeros. At $\alpha = \frac{1}{2}$, the last coefficient is $(K - Z)/4$.

These are ordered limits: first $\epsilon \to 0$ at fixed v, then $v \to \infty$. They do not assert convergence of time derivatives or a joint limit with unbounded $v(\epsilon)$. There is also a global obstruction to using the original completion $A(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)$. For every fixed $0 < t \leq \frac{1}{2}$ and nonzero finite Dirichlet polynomial D, the original weighted error against $Z_t(s) = \mathcal{F}_t(s - \frac{1}{2})/A(s)$ is infinite, unconditionally. Polymath's two-carrier asymptotic [4], with finite-frequency integration by parts, gives a block lower bound $T^{-1}\exp[(t/8)\log^2(T/(2\pi))] \to \infty$. For $b \geq 0$, multiplying the weight at large heights by $e^{-b\log^2(|y|/(2\pi))}$ restores finiteness exactly when $b \geq t/8$ (Note 16, Section 16). This changes the norm; it leaves the compact heat calculation and completed-ratio energy intact.

5 A complete normalized heat limit

The divergence just described motivates removing the leading completion carrier while keeping the original polynomial and boundary weight. Put

$$a(s) = \frac{1}{2s} + \frac{1}{s-1} + \frac{1}{2}\operatorname{Log}\frac{s}{2\pi}, \qquad T_\tau(s) = e^{-\tau a(s)^2/4}\frac{\mathcal{F}_\tau(s-\frac{1}{2})}{A(s)}. \tag{12}$$

The principal logarithm is used on the critical line. For fixed $v \geq 0$, set

$$E_{N,w}^{[v]} = \|1 - T_{v/L^2}V_{N,w}\|_{\mathcal{H}}^2, \qquad f_m(u) = \frac{u^{m-1}w(u)}{(m-1)!},$$

and define the heat profile

$$h_{m,v,w} = -\sum_{k=0}^{\lfloor m/2\rfloor}\frac{m!}{k!(m-2k)!}\left(\frac{v}{4}\right)^k f_m^{(m-2k)}. \tag{13}$$

Here h denotes an approximation profile, not a physical derivative of H.

Theorem 4 (Normalized complete limit)**.** *Assume RH and fix the normalized Gevrey taper of Theorem 1. For every fixed $V < \infty$ and $0 < \eta < 2 - \sigma$,*

$$\sup_{N\geq 2}\sup_{0\leq v\leq V} L(E_{N,w}^{[v]})^{>T} \ll_{W,V,\eta} T^{\sigma-2+\eta}, \qquad T \geq 2, \tag{14}$$

$$\lim_{N\to\infty} LE_{N,w}^{[v]} = \sum_{\rho \text{ distinct}} \frac{\|h_{m_\rho,v,w}\|_2^2}{|\rho|^2} < \infty. \tag{15}$$

The limit is locally uniform in $v \geq 0$. The taper is fixed; no joint limit with unbounded $v = v_N$ is asserted.

Proof mechanism. Note 17, Sections 1–5, starts with the exact complete-source identity

$$\mathcal{F}_\tau(s - \tfrac{1}{2}) = \frac{1}{\sqrt{\pi}} \int_{\mathbb{R}} e^{-x^2} \xi(s + \sqrt{\tau}x)\, \mathrm{d}x.$$

Gaussian domination justifies both integrations before any approximation. Carrier cancellation and reflection give a subpower bound for T_τ uniformly over bounded $\tau \geq 0$. Combining this Gaussian integral with the Möbius convolution retains both remote tails and the entire nearby zero cluster. The same nearest-zero-cell argument as in Theorem 1 pays the scaled height tail. The compact heat polynomial at an order-m zero then gives (13); only after (14) is established is the compact calculation exhausted over all heights.

At $m = 1$ the profile is unchanged. At $m = 2$ it is $h_{2,v,w} = -(uw)'' - (v/2)uw$. In particular the squared local norm of $(T_{v/L^2} - \zeta)V_{N,w}$ at an exact double zero is

$$\frac{v^2\|uw\|_2^2}{4L|\rho|^2} + O_{v,w,\rho}(L^{-2});$$

the norm here is over a fixed isolated neighborhood of that zero. It occurs at the same first-order scale as the proposed heat gain.

Analytic and arithmetic limitations. For each fixed $t > 0$, Note 17, Sections 6–7, proves unconditionally, with $\ell = \log(\sigma/(2\pi))$ and real $\sigma \to +\infty$,

$$T_t(\sigma) = 1 + \frac{t^2\ell^2/64 + t/8}{\sigma} + O_t\left(\frac{(1 + |\ell|)^4}{\sigma^2}\right). \tag{16}$$

An ordinary Dirichlet series converging in a right half-plane has remainder $O(2^{-\sigma})$ after its constant term, so cannot represent this target, even if its initially assumed convergence is conditional. Separately, $s = 1$ becomes an essential singularity:

$$T_t(1 + z) = z^{-1}e^{-t/(4z^2)-tb_0/(2z)}U_t(z), \qquad b_0 = \frac{1 - \log(2\pi)}{2},$$

where U_t is holomorphic and $U_t(0) > 0$. A finite-order zero or pole subtraction cannot repair it. The essential singularity alone would not exclude a Dirichlet series farther right; that conclusion uses (16). Thus the finite critical-line norm does not restore the original Nyman–Beurling/Hardy hypotheses or authorize the original arithmetic lower bound for this changed target.

6 The sharp arithmetic repair of a heat profile

Heating the finite coefficients instead of the numerator is exact:

$$e^{\frac{v}{4L^2}\partial_s^2} V_{N,w} = V_{N,w_v}, \qquad w_v(u) = e^{vu^2/4} w(u). \tag{17}$$

This stays within the fixed Gevrey class, but its original-arithmetic profile is $q_{m,v,w} = -(e^{vu^2/4} f_m)^{(m)}$, not $h_{m,v,w}$. In fact

$$\partial_v (q_{m,v,w} - h_{m,v,w})|_{v=0} = -\tfrac{1}{4}\{u^2 f_m^{(m)} + 2mu f_m^{(m-1)}\} \neq 0.$$

The braces equal $2mu + O(u^2)$ near zero. Even a simple zero has this coefficient-transfer defect. The two complete tail theorems give the actual all-zero transfer limit by polarization; no infinite heat-evolved Euler series is used.

Let Π_m project onto the polynomials of degree below m, and abbreviate $K_m = \mathcal{K}_m(\,\cdot\,, 0)$. Every original arithmetic profile satisfies $\Pi_m p = K_m$, by its first m moments. The least squared correction to a specified heat profile is therefore

$$\delta_{m,v}(w)^2 = \|K_m - \Pi_m h_{m,v,w}\|_2^2, \qquad p_* = h_{m,v,w} + K_m - \Pi_m h_{m,v,w}. \tag{18}$$

This distance is sharp in the closure of real normalized flat Gevrey tapers, without a positivity or monotonicity constraint. To see this, solve $f^{(m)} = -p_*$ with the normalized initial jets. The moment conditions force $f^{(j)}(1) = 0$ for $j < m$; flattening f near one changes its mth derivative by $O(\eta^{1/2})$ in L^2. Orthogonality makes the excess squared correction $O(\eta)$. See Note 18, Sections 1–3.

The correction is explicit. If $b_j = \int_0^1 u^j (p_{m,w} - h_{m,v,w})\,\mathrm{d}u$ and $G_m = (1/(i+j+1))_{0\le i,j<m}$, then

$$\delta_{m,v}(w)^2 = b^* G_m^{-1} b \ge (m - \|h_{m,v,w}\|_2)_+^2.$$

For a double zero, with $I_j = \int_0^1 u^j (uw(u))\,\mathrm{d}u$,

$$\delta_{2,v}(w)^2 = \frac{v^2}{4}(4I_0^2 - 12 I_0 I_1 + 12 I_1^2). \tag{19}$$

A double heat profile with squared norm below two consequently requires a correction strictly greater than $(2-\sqrt{2})^2$ at that zero, before division by $|\rho|^2$.

Theorem 5 (Complete optimal transfer in the declared class)**.** *Assume RH and the complete estimates of Theorems 1 and 4. Fix v, w. Consider families*

$$D_N = \sum_{j=0}^{J} A_j(s) V_{\lfloor N/M \rfloor, w_j}(s), \qquad \sum_{j=0}^{J} A_j \equiv 1,$$

with fixed finite Dirichlet multipliers supported through M and fixed normalized Gevrey tapers. Put $q_\rho = \sum_j A_j(\rho) p_{m_\rho, w_j}$. Then

$$\mathcal{T}(D; v, w) := \lim_{N\to\infty} L\|\zeta D_N - T_{v/L^2} V_{N,w}\|_{\mathcal{H}}^2 = \sum_\rho \frac{\|q_\rho - h_{m_\rho,v,w}\|_2^2}{|\rho|^2}, \tag{20}$$

$$\inf_D \mathcal{T}(D; v, w) = \Delta_v(w) := \sum_\rho \frac{\delta_{m_\rho,v}(w)^2}{|\rho|^2} < \infty. \tag{21}$$

The infimum is over these fixed families after taking their N limits, not over arbitrary coefficient sequences depending on N.

Proof outline. The compact mixed profiles and the two complete tail estimates prove (20). Projection gives the lower bound in (21). Use the original taper as a baseline with finite total transfer cost, repair a finite set of profiles to within any tolerance, and assign them using the bounded two-prime filters. The complete majorant includes the baseline and every selected taper and is independent of filter degree. Dominated convergence pays the exterior profile mass. First take $N \to \infty$ for each fixed family, then increase filter degree and optimize tapers. This gives the matching upper bound (Note 18, Section 5). □

An exact example distinguishing correction from final cost. At $v = 2\pi^2$, take the order-two closure profile

$$f(u) = \frac{(1-u)\sin(\pi u)}{\pi}, \qquad w(u) = f(u)/u, \qquad h = -f'' - \pi^2 f = 2\cos(\pi u).$$

Its heat cost is $\|h\|_2^2 = 2$, whereas

$$\begin{aligned} \delta^2 &= 4 - \frac{48}{\pi^2} + \frac{192}{\pi^4}, \\ p_* &= 2\cos(\pi u) + 4 - \frac{24}{\pi^2} + \left(-6 + \frac{48}{\pi^2}\right) u, \\ \|p_*\|_2^2 &= 6 - \frac{192}{\pi^4} > 4. \end{aligned}$$

Fixed flat cutoffs realize these local limits, with $N \to \infty$ before the cutoff width decreases. The closest repair to h need not minimize the repaired profile's own norm. This example does not raise the original infimum above K; it quantifies the price of retaining this particular heat profile. No uniform full-height limit for a moving base cutoff is claimed.

7 Exact completion still changes the arithmetic

The algebraic error in (16) might suggest heating the completion exactly. Note 18, Sections 6–10, treats that alternative without assuming RH. Write $X_t(s) = \mathcal{F}_t(s - \frac{1}{2})$. For fixed $t > 0$ and $c > 1$, define the entire first-mode factor

$$B_t^c(s) = \frac{1}{\sqrt{\pi t}} \int_c^\infty e^{-(x-s)^2/t} A(x)\, \mathrm{d}x. \tag{22}$$

It is positive on the real axis and solves $\partial_t B_t^c = \frac{1}{4}\partial_s^2 B_t^c$. Splitting the Gaussian formula for X_t at c, and expanding the absolutely convergent Euler series only on $x \geq c$, gives

$$X_t(s) = Q_t^c(s) + \sum_{n \geq 1} n^{-s} e^{t(\log n)^2/4} B_t^c(s - \tfrac{t}{2}\log n). \tag{23}$$

The series converges absolutely and locally uniformly for every complex s; its displayed coefficients depend on s. The entire left contribution satisfies $Q_t^c(\sigma) = O_{t,c}(e^{-(\sigma-c)^2/t})$ for real $\sigma \geq c$. This retains the full negative-real-axis integral.

Proposition 6 (Fixed-time second-mode obstruction)**.** *For fixed $t > 0, c > 1$, as real $\sigma \to +\infty$,*

$$\frac{X_t(\sigma)}{B_t^c(\sigma)} - 1 = 2^{-\sigma} e^{t(\log 2)^2/4} \left(\frac{\sigma}{2\pi}\right)^{-t\log 2/4} \left[1 + O_{t,c}\left(\frac{\log\sigma}{\sigma}\right)\right]. \tag{24}$$

This quotient has no ordinary Dirichlet-series representation converging in a right half-plane, even conditionally.

Proof mechanism. A fixed-shift Laplace estimate for B_t^c gives the $n = 2$ term. Monotonicity on the real axis bounds the complete $n \geq 3$ tail by $O_{t,c}(3^{-\sigma}\sigma^{-t\log 3/4})$; the left term is Gaussian-small. A hypothetical Dirichlet series has expansion $d_1 + d_2 2^{-\sigma} + O(3^{-\sigma})$ sufficiently far right. Equation (24) forces $d_1 = 1, d_2 = 0$, but its nonzero second-mode remainder is not $O(3^{-\sigma})$. □

The same conclusion holds for the explicit contour prescriptions

$$B_{t,\eta}(s) = \frac{1}{\sqrt{\pi t}} \int_{\mathbb{R}+i\eta} e^{-(z-s)^2/t} A(z)\,\mathrm{d}z, \qquad \eta \in \mathbb{R} \setminus \{0\},$$

and their symmetric upper/lower average. Gamma reflection pays the negative horizontal ray. Deforming only the pole-free right part gives $B_{t,\eta}(\sigma) - B_t^c(\sigma) = O_{t,c,\eta}(e^{-(\sigma-c)^2/t})$, preserving (24). More generally the comparison applies to analytic normalizers differing from B_t^c by a Gaussian-small term on this ray. It is not an exclusion of arbitrary completions.

Wherever a heat-solution normalizer B is nonzero, its quotient obeys

$$\partial_t(X_t/B) = \tfrac{1}{4}\partial_s^2(X_t/B) + \tfrac{1}{2}(B'/B)\partial_s(X_t/B).$$

Thus exact division leaves a completion drift. No global zero-free property, critical-line norm theorem for these exact quotients, or uniform assertion for $t = t_N \to 0$ is established here. In particular, Theorem 4 is not being reassigned to these new normalizers.

8 Relation to BW09

The originating BW09 route seeks to rule out a hypothetical positive all-real threshold by forcing a later collision. Its current manuscript assumes an attained finite-height ordinary event at that threshold; coverage of every hypothetical positive threshold remains a separate unproved step. BW10 assumes RH and starts from a multiple zero at the specified time zero, allowing arbitrary finite multiplicity. It removes that unknown-time and attainment issue from its conditional problem. This is a structural distinction, not a proved ordering of difficulty.

The unchanged theta source, local Hermite splitting, root-wall cancellation and complete boundary-phase identities are shared. The ratio-energy analysis also applies relative to a known all-real base time T, with source $e^{Tu^2}\Phi(u)$, provided the all-real forward continuation and order-one product hypotheses are retained. The finite-selector estimates are unconditional filter lemmas. These tools sharpen the accounting of an existing event; they do not supply additional winding.

The strongest approximation estimates have a narrower scope. Theorem 1 uses RH for the original zeta function and its unchanged Möbius convolution. Importing that assumption into BW09's positive-threshold contradiction would assume the desired conclusion. All-realness of a positive-time heat deformation supplies neither this original Dirichlet arithmetic nor a substitute theorem. The normalized heat and sharp transfer theorems inherit this RH premise. In contrast, the fixed-positive-time completion obstructions in Sections 5 and 7 are unconditional and may inform either route, within their stated normalization classes.

BW09 still needs a complete theta estimate forcing an ordinary born pair's cap-phase decrease strictly beyond π, with wall continuation, a regular terminal cap, endpoint deficits and source remainders all controlled. Its fixed-index large-time cosine model produces the ordinary π, not a surplus. BW10 has not supplied that strict estimate or the threshold-event coverage step. See the BW09 manuscript, Sections on scope, root walls, surplus and information limits, and its phase-surplus progress register.

9 What is missing, and what to brainstorm next

The common bottleneck is now explicit. The complete approximation upper bound has coefficient K, and the heat transition has excess coefficient $(K - Z)/4$. Both are compatible with a multiple endpoint. A simplicity proof needs an additional estimate using the unchanged arithmetic of zeta that rules out this excess, or a different mechanism that excludes an original multiple zero. Sections 5–7 make the transfer issue quantitative: normalized numerator heat has a complete norm, but returning its profiles to the original arithmetic class costs $\Delta_v(w)$, and the tested exact completions retain a non-Dirichlet second mode. These results close specific unpaid transfers; they do not rule out every use of heat flow or every possible completion.

The tested absolute prime and Möbius-correlation payments do not supply that estimate. In the exact ratio-energy representation, signed off-diagonal terms cancel a diagonal of size $1/\epsilon$ to bounded size, with the residual energy determined at absolute precision $O(\epsilon)$; taking absolute values gives a divergent double payment (Note 15, Section 6). The prime-error linearization also has a first-order remainder at multiple zeros (Section 7). Neither remainder can be omitted at the scale where simplicity is being tested. Generic positive-source collision controls demonstrate the insufficiency of source positivity alone, while failing the ordinary Dirichlet-series premise; they are not counterexamples satisfying all of zeta's arithmetic properties.

The next discussion should focus on two questions:

1. Is there a specific signed arithmetic identity whose *complete* remainder can be bounded more sharply than these multiplicity-compatible estimates? A proposed bound must retain all zero and prime tails.
2. Is there a theta-specific sign or rigidity principle, absent from the screened collision controls, that excludes the multiple endpoint without merely rewriting the desired conclusion?

Before further expansion, the complete scaled-tail theorem deserves external scrutiny because several later conclusions depend on it. The present estimates warrant consolidation and a change of viewpoint, not a claim of proximity to a proof.

Supporting checks. The exact-algebra replay and bounded Lean check support selected identities and an inner-product tradeoff. The BW09 manuscript cites supporting numerics. These checks do not verify the analytic tail theorem or simplicity and are not proof premises here.

Acknowledgement and draft record. Drafted for Edward Baker with substantial OpenAI `gpt-6-astra` assistance. The original derivations and reviews recorded effort `ultra`; the 21 October manuscript integration of Notes 17–18 and associated discussion recorded effort `max`. Derivations and reviews are internal LLM work. Detailed proofs and corrections are in Note 12, Note 15, Note 16, Note 17, Note 18, and Review 16, Review 17 and Review 18. `../DRAFT_HISTORY.md` indexes milestones of this canonical draft.

References

[1] J.-F. Burnol, *A lower bound in an approximation problem involving the zeros of the Riemann zeta function*, Theorem 1.3. arXiv:math/0103058.

[2] E. Carneiro, V. Chandee and M. B. Milinovich, *Bounding $S(t)$ and $S_1(t)$ on the Riemann hypothesis*, Theorem 2. arXiv:1309.1526.

[3] V. Chandee and K. Soundararajan, *Bounding $|\zeta(1/2 + it)|$ on the Riemann hypothesis*, Theorem 1.1. arXiv:0908.2008.

[4] D. H. J. Polymath, *Effective approximation of heat flow evolution of the Riemann ξ function, and a new upper bound for the de Bruijn–Newman constant*, Proposition 9.1(i). arXiv:1904.12438v2.

[5] R. M. Redheffer, *On a theorem of Plancherel and Pólya*, Pacific J. Math. **3** (1953), 823–835, Section 5, Theorem 11. Primary text.