

1 Model

Stochastic Block Model (SBM)

- Choose the community proportions $\pi \sim \text{Dir}(\gamma)$, where $\pi \in \mathbb{R}^C$ with C latent communities
- For each representative u
 1. Choose a community membership assignment $M_u \stackrel{\text{iid}}{\sim} \text{Cat}(\pi)$
- For each pair of communities $k, l \in \{1, \dots, C\}$, draw coexpression rate $P_{kl} \stackrel{\text{iid}}{\sim} \text{Gamma}(\lambda_0, \lambda_1)$
- For each pair of representatives $u, v \in \{1, \dots, U\}$, draw $R_{uv} \mid P, M_u = k, M_v = l \sim \text{Poisson}(P_{kl})$

Ideal Point Model (IPM)

- For each document d
 1. Choose a discrimination $a_d \sim \mathcal{N}(\eta_a, \sigma_d^2)$
 2. Choose a difficulty $b_d \sim \mathcal{N}(\eta_b, \sigma_d^2)$
- For each representative u
 1. Choose a position $x_u \mid \nu \sim \mathcal{N}(\nu, \sigma_x^2)$
- Draw representative u 's vote on document d as $V_{ud} \mid x_u, a_d, b_d \sim \text{Bern}(\sigma(a_d \cdot (x_u - b_d)))$

Latent Dirichlet Allocation (LDA)

- Draw a topic $\varphi_k \stackrel{\text{iid}}{\sim} \text{Dir}(\beta)$, $\varphi_k \in \mathbb{R}^V$ as a distribution over words, for each $k \in \{1, \dots, K\}$
- For each document, draw the topic proportions $\theta_d \stackrel{\text{iid}}{\sim} \text{Dir}(\alpha)$, where $\theta_d \in \mathbb{R}^K$
- For each document $d \in \{1, \dots, D\}$ and each word $n \in \{1, \dots, N_d\}$ in the document
 1. Choose a topic $z_{dn} \mid \theta_d \stackrel{\text{iid}}{\sim} \text{Mult}(\theta_d)$
 2. Choose a word $W_{dn} \mid z_{dn} = k, \varphi_k \stackrel{\text{iid}}{\sim} \text{Mult}(\varphi_k)$

1.1 Frankenstein Model

The ideal point model (IPM) is useful to us as a baseline model for the roll call voting data (V_{ud}) for a couple of reasons. For one, using it alone we can attempt to predict missing votes, a problem of interest in political science. Another problem of more qualitative interest is analyzing and interpreting the factors a_d, b_d specific to a document and those x_u specific to the representative. All are assumed to reside in some latent space \mathbb{R}^S and so depending on how we set up the model, we might be able to interpret quantities like x_u as u 's *political stance* or *ideological position* or $x_u - b_d$ as representative u 's propensity for the bill/document d . There are a number of problems we cannot address in IPM. A major problem is predicting on heldout documents (the 'cold start'), which is a potentially useful performance measure. Similarly if we have relatively junior representatives, they may not have had enough votes for the inferred x_u to represent something (1) meaningful / interpretable or (2) reliable. We want to incorporate more information to inform the choices of a_d, b_d and x_u .

Ideal Point Allocator (IPA)

- Run the generative processes for SBM and LDA as described above. Then,
- For each document d
 1. Calculate the empirical topic proportions $\bar{z}_d = \frac{1}{N_d} \sum_{i=1}^{N_d} z_d$ (a $K \times 1$ vector)
 2. Generate $S \times K$ matrices η_a, η_b with iid normal entries
 3. Choose a discrimination $a_d \sim \mathcal{N}(\eta_a' \bar{z}_d, \sigma_d^2)$
 4. Choose a difficulty $b_d \sim \mathcal{N}(\eta_b' \bar{z}_d, \sigma_d^2)$
- For each representative u
 1. Generate the community means $\nu_k \sim \mathcal{N}(\tau, \sigma_x^2)$
 2. Choose a position $x_u \mid M_u = k, \nu \sim \mathcal{N}(\nu_k, \sigma_x^2)$
- Draw representative u 's vote on document d as $V_{ud} \mid x_u, a_d, b_d \sim \text{Bern}(\sigma(a_d \cdot (x_u - b_d)))$




Figure 1: IPA graphical model

2 Variational Inference

2.1 For SBM.

After observing the symmetric matrix $R = (R_{uv})$, where R_{uv} is the number of caucuses that representatives u and v have in common, we see to find a distribution q over the latent community assignments $M = (M_u)$, the community coexpression rates $P = (P_{kl})$, and the community proportions $\pi = (\pi_k)$ which is close in relative entropy to the true posterior and lies in the factorized family $q(M)q(P)q(\pi)$. Each factor has free parameters described below and denoted with hats. The approximation q is equivalently scored by the ELBO objective \mathcal{L} , which we break down as:

$$\begin{aligned} \mathcal{L}(q) &= \mathbb{E}_q \left[\log \frac{p(R, M, P, \pi)}{q(M, P, \pi)} \right] \\ &= \underbrace{\mathbb{E}_q \left[\log p(R \mid M, P) + \log \frac{p(P)}{q(P)} \right]}_{\mathcal{L}_{\text{data}}} + \underbrace{\mathbb{E}_q [-\log q(M)]}_{\mathcal{L}_{\text{ent}}} + \underbrace{\mathbb{E}_q [\log p(M \mid \pi)]}_{\mathcal{L}_{\text{local}}} + \underbrace{\mathbb{E}_q \left[\log \frac{p(\pi)}{q(\pi)} \right]}_{\mathcal{L}_{\text{global}}} \end{aligned} \quad (1)$$

Variational Factors. To each u we associate variational parameters $\hat{r}_u = (\hat{r}_{uk})_{k=1}^C$, so

$$q(M) = \prod_{u=1}^U q(M_u \mid \hat{r}_u) = \prod_{u=1}^U \prod_{k=1}^C \hat{r}_{uk}^{\delta_k(M_u)}. \quad (2)$$

We define $q(\pi) \triangleq \text{Dir}(\hat{\gamma}_1, \dots, \hat{\gamma}_C)$ and $q(P) = \prod_{kl} q(P_{kl} \mid \hat{\lambda}_{kl})$ where $q(P_{kl} \mid \hat{\lambda}_{kl}) \triangleq \text{Gamma}(\hat{\lambda}_{0kl}, \hat{\lambda}_{1kl})$.

Computing the ELBO. Now we can write out the component terms of the ELBO more explicitly:

$$\begin{aligned} \mathcal{L}_{\text{data}} &= \mathbb{E}_q \left[\log p(R \mid M, P) + \log \frac{p(P)}{q(P)} \right] = \sum_{kl} \mathbb{E}_q \left[\sum_{u,v} \delta_k(M_u) \delta_l(M_v) \log p(R_{uv} \mid P_{kl}) + \log \frac{p(P_{kl})}{q(P_{kl})} \right] \\ &= - \sum_{u,v} \log R_{uv}! + \sum_{k,l} \left(\lambda_0 \log \lambda_1 - \hat{\lambda}_{0kl} \log \hat{\lambda}_{1kl} - \log \frac{\Gamma(\lambda_0)}{\Gamma(\hat{\lambda}_{0kl})} \right) + \sum_{k,l} \mathcal{L}_{kl}(R) \\ \mathcal{L}_{\text{ent}} &= \mathbb{E}_q [-\log q(M)] = - \sum_{u,k} \mathbb{E}_q [\delta_k(M_u) \log \hat{r}_{uk}] = - \sum_{u,k} \hat{r}_{uk} \log \hat{r}_{uk} \\ \mathcal{L}_{\text{local}} &= \mathbb{E}_q [\log p(M \mid \pi)] = \sum_{u,k} \mathbb{E}_q [\delta_k(M_u) \log \pi_k] = \sum_k N_k \mathbb{E}_q [\log \pi_k] \\ \mathcal{L}_{\text{global}} &= \mathbb{E}_q \left[\log \frac{p(\pi)}{q(\pi)} \right] = \log \Gamma(C\gamma) - C \log \Gamma(\gamma) - \log \Gamma \left(\sum_k \hat{\gamma}_k \right) + \sum_k \{ \log \Gamma(\hat{\gamma}_k) + (\gamma - \hat{\gamma}_k) \mathbb{E}_q [\log \pi_k] \} \end{aligned} \quad (3)$$

where $N_k = \sum_u \hat{r}_{uk}$, $S_{uk} = \sum_v \hat{r}_{vk} R_{uv}$, $N_{kl} = \sum_{uv} \hat{r}_{uk} \hat{r}_{vl}$, $S_{kl} = \sum_{uv} \hat{r}_{uk} \hat{r}_{vl} R_{uv}$, and

$$\mathcal{L}_{kl}(R) = (S_{kl} + \lambda_0 - \hat{\lambda}_{0kl}) \mathbb{E}_q [\log P_{kl}] - (N_{kl} + \lambda_1 - \hat{\lambda}_{1kl}) \mathbb{E}_q [P_{kl}],$$

and the posterior expectations can also be computed explicitly as

$$\mathbb{E}_q [P_{kl}] = \frac{\hat{\lambda}_{0kl}}{\hat{\lambda}_{1kl}}, \quad \mathbb{E}_q [\log P_{kl}] = \psi(\hat{\lambda}_{0kl}) - \log \hat{\lambda}_{1kl}, \quad \mathbb{E}_q [\log \pi_k] = \psi(\hat{\gamma}_k) - \psi \left(\sum_l \hat{\gamma}_l \right)$$

CAVI Updates. The simplest approach to variational inference maximizes the ELBO \mathcal{L} via coordinate-ascent, i.e. choosing the best value of a variational parameter with all others fixed. Iteratively applying these updates, the variational approximation q improves at every step toward some local optimum. Conditional conjugacy yields closed form updates for the global variational parameters.

- **Global Update to $q(\pi)$.** We have $\hat{\gamma}_k = \gamma + N_k$.
- **Global Update to $q(P)$.** We have $\hat{\lambda}_{0kl} = \lambda_0 + S_{kl}$ and $\hat{\lambda}_{1kl} = \lambda_1 + N_{kl}$.
- **Local Update to $q(M)$.** First note

$$\frac{\partial \mathcal{L}_{kl}}{\partial \hat{r}_{uk}} = \begin{cases} 2S_{uk}\mathbb{E}_q[\log P_{kk}] - 2N_k\mathbb{E}_q[P_{kk}], & k = l \\ S_{ul}\mathbb{E}_q[\log P_{kl}] - N_l\mathbb{E}_q[P_{kl}] & k \neq l \end{cases}$$

We want to update \hat{r}_{uk} subject to the constraint that $\sum_k \hat{r}_{uk} = 1$, so augment \mathcal{L} with Lagrange multipliers $\tilde{\mathcal{L}} = \mathcal{L} + \sum_u \kappa_u (\sum_k \hat{r}_{uk} - 1)$.

$$\begin{aligned} 0 &= \frac{\partial \tilde{\mathcal{L}}}{\partial \hat{r}_{uk}} = -\log \hat{r}_{uk} - 1 + \mathbb{E}_q[\log \pi_k] + \frac{\partial \mathcal{L}_{kk}}{\partial \hat{r}_{uk}} + \sum_{l \neq k} \frac{\partial \mathcal{L}_{kl}}{\partial \hat{r}_{uk}} + \frac{\partial \mathcal{L}_{lk}}{\partial \hat{r}_{uk}} \\ &= -\log \hat{r}_{uk} - 1 + \mathbb{E}_q[\log \pi_k] + \sum_l S_{ul} (\mathbb{E}_q[\log P_{kl}] + \mathbb{E}_q[\log P_{lk}]) - N_l (\mathbb{E}_q[P_{kl}] + \mathbb{E}_q[P_{lk}]) \end{aligned}$$