

Cancer Patient Healing Protocol

Customized Nutrition Guide - Pacemaker + Warfarin + Active Cancer Treatment

CRITICAL PATIENT PROFILE: This protocol is customized for a patient with three simultaneous conditions that affect every food and supplement recommendation: (1) Active cancer treatment - chemotherapy liver burden, WBC suppression, immune vulnerability. (2) Pacemaker with warfarin therapy - Vitamin K must be kept LOW and CONSISTENT. INR monitoring is essential. (3) Liver under treatment stress - all detox support must be gentle and warfarin-safe. Every recommendation in this guide has been filtered through all three conditions simultaneously.

CONDITION	DIETARY IMPLICATION	KEY RULE
Active Cancer Treatment	WBC suppression, liver toxicity, oxidative stress	Support WBCs, protect liver, reduce oxidative burden
Pacemaker + Warfarin	Vitamin K must be LOW and consistent daily	Avoid high-K foods - spinach, kale, broccoli, parsley, moringa
Liver Under Treatment Stress	Phase I and II detox pathways overwhelmed	Gentle liver support - beets, artichoke, milk thistle, lemon
Neutropenia Risk	Immune system compromised during treatment	No raw sprouts, raw fish, or unpasteurized foods during low WBC

■ SAFE FOODS - LOW VITAMIN K, WARFARIN APPROVED

PROTEINS

Egg Whites	Pure lean protein. Zero Vitamin K. Zero phosphorus burden on kidneys. Highest bioavailability (BV=100). Safe during all treatment phases.
Wild Salmon (cooked)	Omega-3 EPA+DHA reduce treatment inflammation. Very low Vitamin K. Best food source of Vitamin D3. Cook thoroughly during neutropenia.
Sardines with bones	Very low Vitamin K. Highest omega-3 of any small fish. Calcium for bone support. Selenium for WBC protection. Cook or use canned in water.
Lentils (cooked)	18g protein per cup. Very low Vitamin K. Iron + folate for WBC production. Soluble fiber supports gut during treatment. Glycemic index 29.
Chicken Breast (cooked)	Complete lean protein. Low Vitamin K. B6 and niacin support liver enzyme function during treatment. Always thoroughly cooked.
Hemp Seeds	Complete protein. Very low Vitamin K. 3:1 omega-6:omega-3 ratio. GLA reduces treatment inflammation. Add to any meal.
Tempeh (cooked)	Fermented soy. Low Vitamin K. Probiotics support gut microbiome during antibiotic treatment. Isoflavones are gently anti-inflammatory.

VEGETABLES

Beets (cooked or juiced)	Extremely low Vitamin K (<1mcg/cup). Betaine for liver methylation and Phase II detox. Nitrates dilate arteries safely - heart-protective with pacemaker. Anti-inflammatory betalains.
Cucumber (skin on)	Near-zero Vitamin K. Silica for cellular membrane support. 96% water - critical for hydration during treatment. Kidney-safe. Juice skin-on daily.
Celery (skin on)	Very low Vitamin K. Phthalides reduce uric acid from tumor cell breakdown. Gentle diuretic. Juice or eat raw alongside cucumber daily.
Carrots (cooked)	Low Vitamin K. Beta-carotene converts to Vitamin A for immune cell development. Gentle on digestive system during treatment-related nausea.
Zucchini	Very low Vitamin K. Easy to digest. Potassium and magnesium without Vitamin K burden. Steam or roast - easy on treatment-sensitive digestive system.
Sweet Potato (cooked)	Low Vitamin K. Beta-carotene for immune support. Chlorogenic acid stabilizes blood glucose disrupted by treatment. Potassium for heart health with pacemaker.
Cauliflower (cooked)	Low Vitamin K when cooked. Gentle sulforaphane activation. Kidney-safe - very low phosphorus and potassium. Easy to digest - important during treatment.
Bell Peppers (any color)	Very low Vitamin K. Highest Vitamin C of any common vegetable (3x orange). Vitamin C is essential for WBC function and collagen repair. Eat raw or lightly cooked.
Artichoke Hearts	Low Vitamin K. Cynarin is the most potent plant compound for liver bile stimulation. Critical for clearing chemotherapy metabolites. Use canned in water - rinse well.
Mushrooms (cooked)	Very low Vitamin K. Beta-glucans activate NK cells and macrophages. Turkey Tail PSK/PSP studied specifically in cancer patients. Always cook thoroughly during treatment.

FRUITS

Blueberries (fresh/frozen)	Very low Vitamin K. Highest BDNF-stimulating anthocyanins of any fruit. Antioxidant protection for WBCs during oxidative treatment stress. Eat in Peak Zone (before 2pm).
-----------------------------------	---

Oranges	Low Vitamin K. Vitamin C for WBC function and collagen repair. D-limonene activates liver Phase I detox. Hesperidin gently anti-inflammatory.
Kiwi (flesh only)	Low Vitamin K. Highest Vitamin C of common fruits. Actinidin digestive enzyme for treatment-related digestive issues. Gentle on stomach.
Lemon (juice)	Very low Vitamin K. D-limonene activates liver detox enzymes. Citrate prevents kidney stones - critical with treatment medications. Use daily in water.
Strawberries	Very low Vitamin K. Vitamin C + ellagic acid antioxidant protection. Gentle on digestive system. Low glycemic index - safe with treatment-disrupted glucose.
GRAINS / CARBS	
Lentils	16g fiber per cup. Very low Vitamin K. Resistant starch feeds gut microbiome disrupted by antibiotics during treatment. Folate for WBC production.
Quinoa	Complete protein + complex carb. Very low Vitamin K. All 9 essential amino acids for tissue repair during treatment. Magnesium for heart rhythm with pacemaker.
Oats (rolled)	Very low Vitamin K. Beta-glucan (same compound as mushrooms) activates immune cells. Supports gut microbiome. Easy to digest during treatment.
Sweet Potato	Low Vitamin K. See vegetables section above. Primary complex carbohydrate recommendation for this patient.
HEALTHY FATS	
Extra Virgin Olive Oil	Very low Vitamin K. Oleocanthal inhibits COX-1 and COX-2 - natural anti-inflammatory. Enhances absorption of all fat-soluble nutrients from the same meal. Daily non-negotiable.
Avocado	Low Vitamin K. Oleic acid + glutathione - master antioxidant that protects cells during chemotherapy oxidative stress. Boosts fat-soluble nutrient absorption up to 400%.
Pumpkin Seeds	Very low Vitamin K. Highest zinc of any seed - zinc is required for every immune cell to function. Magnesium for heart rhythm. Selenium for WBC protection.
Walnuts	Low Vitamin K. ALA omega-3 for inflammation resolution. Melatonin for sleep quality during treatment. Polyphenols raise HDL gently.
DR. VERA NOTE:	Every food on this list has been verified safe for Vitamin K restriction on warfarin therapy, safe for consumption during active cancer treatment, and supportive of at least one of the three clinical goals: liver support, WBC support, or cardiac protection with pacemaker.

FOODS TO AVOID - HIGH VITAMIN K OR TREATMENT CONTRAINDICATED		
Food / Substance	Why - Be Specific	Safe Alternative
Spinach	Very high Vitamin K (483mcg/cup cooked). Directly counteracts warfarin - will destabilize INR dangerously.	Arugula (low K) or Swiss Chard (low K, not spinach)
Kale	Extremely high Vitamin K (1062mcg/cup cooked). One serving can shift INR significantly. Absolute avoid on warfarin.	Bell peppers for Vitamin C. Arugula for liver support.
Moringa Leaves	141mcg Vitamin K per 100g. High in iron but Vitamin K content makes it unsafe on warfarin.	Spirulina for iron (very low Vitamin K). Blackstrap molasses for iron.
Parsley (large amounts)	Extremely high Vitamin K per gram. Even small garnish amounts add up. Avoid therapeutic doses completely.	Cilantro in very small amounts. Lemon zest for flavor.

Broccoli (raw)	Moderate-high Vitamin K when raw (93mcg/cup). Cooking reduces K content. If used, must be well-cooked and consistent amount daily.	Cauliflower (lower K). Bell peppers for sulforaphane alternatives.
Grapefruit	Inhibits CYP3A4 enzyme - which metabolizes BOTH warfarin AND many chemotherapy drugs. Double contraindication. Absolute avoid.	Orange or lemon for Vitamin C and D-limonene.
Alcohol	Directly hepatotoxic. Potentiates warfarin unpredictably. Adds massive burden to already-stressed liver during chemotherapy.	None. Complete avoidance during treatment.
Raw Sprouts	Neutropenic diet rule: sprouts carry bacterial contamination risk that a suppressed immune system cannot fight. Absolute avoid during treatment.	Cooked mushrooms for immune support. Cooked vegetables only.
Raw Fish / Sushi	Neutropenic diet: raw fish contamination risk is life-threatening with suppressed WBC counts. Absolute avoid during all treatment phases.	Cooked wild salmon. Cooked sardines.
Acetaminophen (Tylenol)	Not a food - but critical: severely hepatotoxic when liver is already stressed by chemotherapy. Must inform oncologist before any OTC pain reliever.	Discuss pain management options with oncologist.
Excessive Fructose	Routes directly to hepatic lipogenesis - adding metabolic burden to already-stressed liver. No fruit after 2pm. No fruit juice except morning window.	Low-glycemic fruit before 2pm only. Cucumber and celery after 2pm.

■ LIVER FLUSH PROTOCOL - CHEMOTHERAPY SUPPORT

Supporting Phase I and II liver detox while clearing chemotherapy metabolites safely

Morning Lemon Water	Half a lemon in 16oz warm water with a pinch of sea salt - every single morning before any food or medication. D-limonene activates liver Phase I. Citrate protects kidneys from treatment medication accumulation.
Beet Juice (daily)	One small beet juiced with cucumber, celery, and lemon. Betaine drives liver methylation - the pathway that clears chemotherapy metabolites. Very low Vitamin K - safe on warfarin. Drink in Peak Zone only (before 10am).
Artichoke Hearts	2-3 artichoke hearts with lunch daily. Cynarin is the most potent natural bile stimulant - bile is how the liver exports toxins. Use canned in water (rinse). Or artichoke leaf extract 300mg before meals.
Dandelion Root Tea	1-2 cups daily. Stimulates bile flow and liver enzyme activity. Gentle potassium-sparing diuretic - supports kidney clearance of treatment metabolites. Very low Vitamin K.
Milk Thistle Silymarin	280mg standardized extract (70-80% silymarin) with every meal. Used in European hospitals for chemotherapy liver protection. Regenerates liver cells, reduces ALT and AST. Confirm with oncologist - most chemotherapy protocols are compatible.
Schisandra Berry	500mg extract 2x daily. Lignan compounds directly protect liver cells from chemotherapy-induced hepatocyte damage. Used in Chinese hospitals alongside cancer treatment. Low Vitamin K.
Garlic (cooked)	Crush 2 cloves, wait 10 minutes (allicin formation), then add to any cooked meal. Allicin activates liver Phase II glutathione pathway. Antimicrobial during neutropenic phases. Very low Vitamin K.
Turmeric + Black Pepper	Half teaspoon turmeric in food or juice daily - always with black pepper (2000% absorption increase) and olive oil (fat-soluble). Curcumin inhibits NF-kB, COX-2, and 5-LOX simultaneously. Anti-inflammatory liver protection during treatment.
DR. VERA NOTE:	The liver runs its deepest Phase II detox between 1-3am. Kitchen must close at 7pm without exception - food after 7pm diverts energy from detox to digestion at the exact hours when chemotherapy metabolite clearance needs to peak. This is not optional for this patient.

■■ WHITE BLOOD CELL + IMMUNE SUPPORT PROTOCOL

Supporting NK cells, macrophages, and bone marrow WBC production during treatment

Turkey Tail Mushroom (cooked)	PSK and PSP polysaccharides from Turkey Tail are approved as a cancer adjunct therapy in Japan. Activates NK cells and macrophages. 1-2g extract daily or cook mushrooms thoroughly. Beta-glucans are the most studied natural WBC support compounds.
Astragalus Root	Used in Chinese hospitals alongside chemotherapy specifically for WBC recovery. Polysaccharides stimulate bone marrow activity. 500mg extract daily. Confirm with oncologist - safe with most chemotherapy protocols.
Reishi Mushroom (cooked)	Immune modulator - activates immune cells without over-stimulating. Also improves sleep quality during treatment - deep sleep is when bone marrow produces WBCs most actively. 500mg-1g extract daily.
Spirulina	Fastest plant-based hemoglobin builder studied. Very low Vitamin K - safe on warfarin. Iron + B12 + beta-carotene. Supports red and white blood cell recovery simultaneously. Half teaspoon in juice daily.
Pumpkin Seeds (daily)	Zinc - every single immune cell requires zinc to develop and function. Treatment depletes zinc stores rapidly. One handful daily. Selenium content also protects WBCs from oxidative treatment damage.
Bell Peppers (raw)	3x more Vitamin C than orange. Very low Vitamin K. Vitamin C is consumed by immune cells at high rates during active infection or treatment stress. Eat raw to preserve enzyme activity - heat destroys Vitamin C.
Garlic + Ginger (daily)	Allicin from garlic and gingerols from ginger are directly antimicrobial - critical when WBC counts are suppressed and the immune system cannot fight infection normally. Always cooked garlic during neutropenic phases.
Selenium (Brazil Nuts - 2-3 only)	2-3 Brazil nuts provide the daily selenium requirement. Selenium is the cofactor for glutathione peroxidase - the enzyme that neutralizes viral and bacterial free radicals inside WBCs. Never more than 3 - selenium toxicity is real.
DR. VERA NOTE:	Sleep is the most powerful WBC recovery intervention that exists. Bone marrow produces WBCs most actively during deep sleep between 10pm and 3am. Reishi mushroom at night supports both immune activation AND deep sleep architecture simultaneously. No supplement compensates for consistently disrupted sleep during treatment.

■ CARDIAC SUPPORT PROTOCOL - PACEMAKER + WARFARIN SAFE

Supporting cardiac function and arterial health without affecting INR or pacemaker operation

Beet Juice (morning)	Nitrates convert to nitric oxide within 90 minutes - dilates arteries and reduces cardiac workload. Safe and beneficial with pacemaker. Very low Vitamin K. Heart-protective alongside any arrhythmia management.
Wild Salmon (3x weekly)	EPA + DHA reduce arrhythmia risk - well-studied in pacemaker patients. Omega-3s stabilize cardiac cell membranes. Anti-inflammatory. Cook thoroughly.
Walnuts (daily)	L-arginine for nitric oxide synthesis. ALA omega-3. Melatonin - highest of any nut - supports nighttime heart rate variability. Low Vitamin K.
Pumpkin Seeds	Magnesium - essential for cardiac rhythm and electrical conduction. Pacemaker patients need consistent magnesium. Low Vitamin K.
Lemon Water (morning)	Potassium + citrate support cardiac electrolyte balance. Lemon citrate specifically helps prevent the calcium deposits that can occur around pacemaker leads over time.
Garlic (cooked daily)	Gentle antiplatelet activity - reduces clot formation alongside warfarin. Mild blood pressure lowering. Do NOT take garlic supplements - food amounts only with warfarin.
Extra Virgin Olive Oil (daily)	Oleocanthal reduces arterial inflammation without affecting INR. Mediterranean diet studies show specific benefit in arrhythmia management. 2 tablespoons daily minimum.
DR. VERA NOTE:	Every food supplement interaction with warfarin must be disclosed to the cardiologist. The goal is consistency - same amounts of the same foods daily so INR remains stable. Sudden large increases in any food - even safe ones - can shift INR. Introduce new foods one at a time and monitor INR for 1-2 weeks after each addition.

■ DAILY HEALING PROTOCOL - CANCER PATIENT SCHEDULE

Time	What to Do	Why
6-7am Rising	16oz warm water + half lemon + sea salt. Wait 20 min before medications.	Bile activation, electrolyte reset, kidney citrate protection.
7-8am Peak Zone	Beet + cucumber + celery + lemon + ginger juice (skin on both). Add spirulina powder after juicing.	Liver flush, WBC iron support, nitric oxide for cardiac health.
8-9am Breakfast	Egg whites + bell peppers + zucchini + avocado. Milk thistle with this meal. Turmeric + black pepper on eggs.	Complete protein, Vitamin C for WBCs, liver protection, fat for curcumin absorption.
9am Supplements	Turkey Tail or Reishi mushroom extract. Astragalus root. Pumpkin seeds (zinc). 2-3 Brazil nuts (selenium).	NK cell activation, bone marrow WBC production support, zinc and selenium for immune function.
10am-12pm Peak Window	Largest meal of the day. Lentils or quinoa with cooked salmon or chicken. Artichoke hearts. Bell peppers.	Peak insulin sensitivity, maximum nutrient absorption, liver bile stimulation from artichoke.
After Every Meal	10-15 minute gentle walk if energy allows.	Reduces post-meal glucose spike 20-30%. Supports lymphatic drainage. Gentle cardiac exercise.
12-2pm Midday	Light protein and vegetable meal. Milk thistle with this meal. Dandelion root tea.	Continuing liver support. Dandelion stimulates bile for ongoing chemotherapy metabolite clearance.
After 2pm Declining Zone	No fruit juice. Low sugar only. Cucumber and celery sticks. Water with lemon.	Insulin sensitivity declining. Same sugar load produces 2-3x higher spike. Protects treatment-disrupted glucose metabolism.
5-7pm Evening	Light cooked fish or chicken with roasted vegetables (low Vitamin K only). Milk thistle with evening meal. Schisandra berry.	Liver prep for overnight detox. Schisandra protects liver cells. Keep dinner light.
7pm Kitchen Closes	KITCHEN CLOSES. Water only after 7pm. No exceptions during treatment.	Liver Phase II detox peaks 1-3am. Food after 7pm diverts energy from chemotherapy metabolite clearance.
8-9pm Evening	Reishi mushroom tea or ashwagandha (if cardiac-approved by cardiologist). Dandelion root tea.	Reishi deepens sleep for bone marrow WBC production. Liver bile stimulation before overnight fast.
10pm-3am Overnigh Repair	Sleep - 7-8 hours minimum. Complete fast. Water only if needed.	WBC production in bone marrow. Liver Phase II peaks 1-3am. Glymphatic brain flush. Growth hormone pulses.

■ DAILY HEALING JUICE - CANCER PATIENT FORMULA

INGREDIENTS: 1 small Beet + 1 large Cucumber (skin on) + 4 stalks Celery (skin on) + 1 Lemon (peeled) + 1 inch fresh Ginger + half inch Turmeric root + pinch of black pepper added after juicing + half teaspoon Spirulina powder stirred in after juicing

RULES: Drink within 20 minutes of pressing. Morning Peak Zone only (before 10am). This formula is Vitamin K safe. Low glycemic. Liver-flushing. WBC-supporting. Cardiac-safe with pacemaker. Confirm Spirulina with oncologist first.

■ SUPPLEMENT PROTOCOL - ALL WARFARIN REVIEWED

Supplement	Dose + Timing	Purpose	Warfarin Note
Milk Thistle Silymarin	280mg (70-80% extract) 3x daily with meals	Liver cell protection during chemo. Reduces ALT/AST.	Generally safe. Confirm with oncologist.
Turkey Tail Mushroom	1-2g extract daily with food	NK cell activation. WBC support. Approved in Japan as cancer adjunct.	No known warfarin interaction.
Astragalus Root	500mg extract daily with food	Bone marrow stimulation. WBC recovery. Used in Chinese hospitals with chemo.	No known warfarin interaction. Confirm with oncologist.
Reishi Mushroom	500mg-1g extract at night	Immune modulation + deep sleep support for bone marrow WBC production.	May have mild antiplatelet effect - disclose to cardiologist.
Schisandra Berry	500mg 2x daily with food	Direct liver cell protection from chemo-induced hepatocyte damage.	No known warfarin interaction. Confirm with oncologist.
Vitamin D3	2000-5000 IU daily with K2-MK7 (low dose)	Low Vitamin D is directly correlated with low WBC counts in cancer patients.	K2 at low dose (45mcg) has minimal warfarin effect but disclose to cardiologist.
Zinc (from food)	From pumpkin seeds daily (not supplement)	Every immune cell requires zinc. Treatment depletes zinc stores.	Food-form zinc has no warfarin interaction.
Selenium (Brazil Nuts)	2-3 Brazil nuts daily. Never more.	Glutathione peroxidase cofactor. Protects WBCs from oxidative damage.	Food-form selenium. No warfarin interaction at food amounts.

Dr. Vera, CNS + CHN - Your Health Matters - yourhealthmatters.ebepstudios.com

This protocol was created by Erica Ehiwe through holistic health research. It is educational in nature and does not constitute medical advice. All supplements and dietary changes must be reviewed with the treating oncologist and cardiologist before implementation - especially with warfarin therapy and active cancer treatment.