Convergence in rich regime

Contradicts Chizat & Bach (2018)?

25

Theorem (informal) [Chizat & Bach '18]

For smooth activations and the initialisation $\begin{cases} w_i \sim \frac{\lambda}{\sqrt{m}} \mathcal{U}(\mathbb{S}_{d-1}) \\ a_i \sim \frac{\lambda}{\sqrt{m}} \mathcal{U}(\{-1,1\}) \end{cases}$

$$\lim_{m \to \infty} \lim_{t \to \infty} L(\theta_m^t) = 0$$

→ **smoothness allows to preserve omnidirectionality of the neurons**

→ **omnidirectionality is essential to global convergence**

Convergence in rich regime

Contradicts Chizat & Bach (2018)?

Theorem (informal) [Chizat & Bach '18]

For smooth activations and the initialisation $\begin{cases} w_i \sim \frac{\lambda}{\sqrt{m}} \mathcal{U}(\mathbb{S}_{d-1}) \\ a_i \sim \frac{\lambda}{\sqrt{m}} \mathcal{U}(\{-1,1\}) \end{cases}$

$$\lim_{m \to \infty} \lim_{t \to \infty} L(\theta_m^t) = 0$$

→ **smoothness allows to preserve omnidirectionality of the neurons**

→ **omnidirectionality is essential to global convergence**

Training dynamics