

# Homework 2

Erik Brakke

September 16, 2015

**Collaborators:** .

## Answer 1

(a) Thm: If  $x \equiv y \pmod{p-1}$  then for any  $a$ ,  $a^x \equiv a^y \pmod{p}$

Proof:  $x = (p-1)k_x + r$  and  $y = (p-1)k_y + r$  (by the unique fact about division)  
 Therefore,  $a^{(p-1)k_x+r} \equiv a^{(p-1)k_y+r} \pmod{p}$  (using substitution)  
 $a^{(p-1)k_x+r} \pmod{p} = a^{(p-1)k_y+r} \pmod{p}$  (fact about congruency)  
 $(a^{(p-1)})^{k_x} \pmod{p} * a^r \pmod{p} = (a^{(p-1)})^{k_y} \pmod{p} * a^r \pmod{p}$  (property of exponents and proof from HW1 that the order of 'mod' does not matter)  
 $1 * a^r \pmod{p} = 1 * a^r \pmod{p}$  (by Fermat's little theorem)  
 $a^r \equiv a^r$   
 (fact about congruency)  
 $r = x \pmod{p-1} = y \pmod{p-1}$  (by definition of 'mod' and our premise)  
 Therefore, if  $x \equiv y \pmod{p-1}$  then for any  $a$ ,  $a^x \equiv a^y \pmod{p}$

□

(b) Thm: if  $g$  is a generator, then  $g^x \equiv 1$  if and only if  $(p-1) \mid x$

Proof: Assume  $g^x \equiv 1$   
 $g^{p-1} \equiv 1$  ( $g \in \mathbb{Z}_p^*$  by def'n of generator, Fermat's Little Theorem)  
 $x = p-1$  (substitution)  
 Therefore,  $(p-1) \mid x$  (definition of divides)

Now, assume  $(p-1) \mid x$   
 Let  $r = x \pmod{p-1}$  (Def'n of 'mod')  
 $r = 0$  (def'n of divides)  
 Consider  $g^r \pmod{p}$   
 $g^{x \pmod{p-1}}$  (substitution)  
 $g^0 = 1$  (because  $(p-1) \mid x$ )  
 Therefore  $g^x \equiv 1$

□

(c) Thm: if  $g$  is a generator, and  $g^x \equiv g^y$  then  $x \equiv y \pmod{p-1}$

Proof: Let's assume  $g^x \equiv g^y$  and  $x \not\equiv y \pmod{p-1}$

$x \bmod p-1 \neq y \bmod p-1$  (Fact of congruency)

$r_x \neq r_y$  (definition of mod)

This means that  $\exists_{r_x, r_y} r_x r_y \in (1, \dots, p-1), r_x \neq r_y$  and  $g^{r_x} \equiv g^{r_y}$

However,  $g$  is a generator, which means that each element in  $(1, \dots, p-1)$  maps to a distinct element in  $(1, \dots, p-1)$  (def'n of generator)

Therefore,  $g^{r_x} \not\equiv g^{r_y}$  which means  $g^x \not\equiv g^y$

This is a contradiction, therefore the statement must be true  $\square$

(d) Thm: If  $g$  is a generator, and  $a = g^x \pmod{p}$ , and  $x$  is even, then  $a$  has a square root modulo  $p$

Proof: Because  $x$  is even, we can rewrite it as  $2y$  where  $y$  is also a number in  $\{1, \dots, p-1\}$

$a = g^{2y} \pmod{p}$

$a = (g^y \pmod{p}) * (g^y \pmod{p})$  (Splitting exponents with like bases)

Because  $g$  is a generator, we know that  $g^y \in \{1, \dots, p-1\}$  and  $g^y \not\equiv g^x$  (def'n of generator)

Therefore  $g^y$  is the square root of  $a$  (Knowledge of square roots)  $\square$

Thm: if  $a$  has a square root modulo then  $x$  is even

Proof: Let's represent  $a$  as a generator  $g$  raised to some  $x \bmod p$ .  $a = g^x \pmod{p}$

Let's also assume that  $x$  is odd

$g^x \equiv g^y * g^y$  (because we assume that  $a$  has a square root)

$g^x \equiv g^{2y}$

This means that  $x = 2y$

This is a contradiction, because we assumed  $x$  was even

Therefore, if  $a$  has a square root, then  $x$  must be even.  $\square$

(e) Thm: If  $a$  is a square, then  $a^{\frac{p-1}{2}} \equiv 1$

Proof: Let's assume there is a generator  $g$  such that  $g^x \equiv a$

We know that  $x$  must be even (by the previous part)

$x = 2y$  for some  $y \in \{1, \dots, p-1\}$

$a \equiv g^{2y}$

Now consider  $(g^{2y})^{\frac{p-1}{2}}$

$g^{y(p-1)}$  (2's cancel)

Because  $(p-1) \mid y(p-1)$  we know that  $g^{y(p-1)} \equiv 1$  (proof from (b))

Therefore,  $a^{\frac{p-1}{2}} \equiv 1$   $\square$

Thm: If  $a$  is non-square, then  $a^{\frac{p-1}{2}} \not\equiv 1$

Proof: Let's assume there is a generator  $g$  such that  $g^x \equiv a$

We know that  $x$  must be odd (from proof (d))

Now consider  $(g^x)^{\frac{p-1}{2}}$

$g^{\frac{x}{2}(p-1)}$  (using rules of exponents)

$(p-1) \nmid \frac{x}{2}(p-1)$  therefore,  $a^{\frac{p-1}{2}} \not\equiv 1$   $\square$

## References

None