

# 1 Introduction

- "Because two-locus and other types of model cannot describe inheritance and selection for continuously varying traits (such as ornaments and preferences), these models are fundamentally disconnected from the empirical study of sexual selection," [7]
- "The male traits most likely to become exaggerated by such mechanisms are those under weak natural selection and subject to relatively large variance in female sexual preferences, such as some behavioral and morphological elements of courtship and mating," [6]
- quantitative genetics model / like mine: "Variation among females in their preferences selects for increased variance in the male trait; we assume that this effect is weak by requiring that  $\sigma_z^2 \sigma_y^2 \ll 1$ ," [4] WHY?!?!]
- two-allele model: "Neither copying model supports variation at the male trait locus, and copying makes it more difficult for a novel male trait phenotype to spread," [3]. "Since Lill's study, anecdotal evidence for copying has also been found in other species of birds, mammals, and fishes (reviewed by Gibson and Hoglund 1992; Pruett-Jones 1992)," [3]. "Bradbury and Gibson (1983) and Bradbury et al. (1985) found that the large variance of male mating success observed in sage grouse could not be plausibly attributed to independent female choice, and suggested that copying might be responsible," [3]. "Y-linkage is of interest because many of the loci that affect sexually-selected male coloration in guppies are Y-linked (Winge 1927; Haskins et al. 1961)," [3]. "Copying itself does not support polymorphism in the male trait, at least in haploid and Y-linked systems," [3]. "Guppies are a perplexing example in this context. The species is famous for highly polymorphic male coloration (Endler 1983) which is known to be under sexual selection (Endler 1980, 1983; Breden and Stoner 1987; Houde 1987, 1988; Stoner and Breden 1988; Houde and Endler 1990) and is at least in part caused by Y-linked loci (Winge 1927; Haskins et al 1961)," [3].
- "Mate preferences can be genetically determined (Saether et al. 2007), but they can also be earned (Virzijden et al. 2012). For example, in sticklebacks females learn to prefer mates with phenotypes similar to their fathers (Kozak et al. 2011) and in some cichlids females learn to prefer mates with phenotypes similar to their mothers (Verzijden et al. 2008). Recent work suggests that learned preferences for parental phenotypes can promote speciation (Verzijden et al. 2005; Servedio and Dukas 2013)," [1].

# 2 Question

1. H1 Song learning can decelerate speciation by allowing for genetically diverse birds to mate with each other, maintaining gene flow between subpopulations that might either begin to diverge
2. H2 Song learning can accelerate speciation by increasing standing genetic variation, which would allow for quicker divergence once new selection pressures arise [5]
3. H3 Song learning can accelerate speciation because culturally inherited traits can evolve more quickly than genetically inherited ones [2]

Table 1: Summary of choices made in previous models

	Lachlan and Servedio [5]
$\sigma$ trait(s)	allele A / a: song predisposition song
How $\sigma$ trait is learned	
Inherited	
$\varphi$ trait(s)	
How $\varphi$ trait is learned	
Inherited	
Errors	
Population structure	
Mating structure	
$\varphi$ preference	
Selection	

### 3 Goals

### 4 Model

**4.1 Continuous traits** Each male has a song and each female has preference for a particular song. She will mate with males with songs other than her preferred song, but the probability of her doing so decreases as the potential mate’s song gets less similar to her preferred song. Each female mates once and chooses a male according to her preferences and the distribution of songs present in the population. Each male, therefore, may breed multiple times or not at all. We assume that generations are non-overlapping, so once the adults breed they die and we can shift our focus to the new generation. To begin with, we assume that males acquire their songs from their fathers at birth and females acquire their preferences from their mothers at birth. We further assume an unbiased sex ratio. Before the new generation mates, each male has a small probability of “mutating” its song. This can be interpreted as a learning error or as innovation. After mutation, the new males and females mate.

Mathematically, each male has a song  $x \in \mathbb{R}$  and each female has a preference  $y \in \mathbb{R}$ . The probability density of the male songs will be written  $P_m(x)$  and the probability density of female preferences will be written  $P_f(y)$ . The preference of a female with preference  $y$  mates for a male with song  $x$  is

$$f_y(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-y)^2}{2\sigma^2}\right),$$

which is maximal when  $x = y$  and decreases as  $|x - y|$  increases. The probability that a female with preference  $y$  choose a male with song  $x$  is

$$\frac{P_m(x)f_y(x)}{\int_{\mathbb{R}} P_m(x')f_y(x')dx'},$$

Table 2: Parameters

$\sigma^2$	width of female preference function
$\sigma_f^2$	variance of female preferences within each population
$\sigma_m^2$	variance of male traits within each population

so the probability of a  $(x, y)$  pair mating is then

$$P_{\text{mate}}(x, y) = \frac{P_f(y)P_m(x)f_y(x)}{\int_{\mathbb{R}} P_m(x')f_y(x')dx'} = \frac{P_f(y)P_m(x)f_y(x)}{Z_y},$$

where  $Z_y$  is the normalizing factor  $\int_{\mathbb{R}} P_m(x')f_y(x')dx'$ . Each such pair will produce a male with song  $x$  and a female with preference  $y$ . Before mating, the male's song changes to  $x - \delta_{\text{mut}}$  with probability  $p_{\text{mut}}/2$  and to  $x + \delta_{\text{mut}}$  with probability  $p_{\text{mut}}/2$ . Under these assumptions, the probability density of female preferences does not change over time:

$$P_f(y, t + 1) = P_f(y, t).$$

The probability density of male songs in the next generation, before mutating and after mutating, follows

$$P_m(x, t + 1/2) = \int_{\mathbb{R}} P_{\text{mate}}(x, y, t)dy$$

$$P_m(x, t + 1) = (1 - p_{\text{mut}})P_m(x, t + 1/2) + p_{\text{mut}}/2P_m(x - \delta_{\text{mut}}, t + 1/2) + p_{\text{mut}}/2P_m(x + \delta_{\text{mut}}, t + 1/2)$$

(Population dynamics are modeled in Code ???. Global and / or needed parameters are given in Code 2.)

We start with mixtures of normal distributions of both male and female:

$$P_f(y, 0) = p_f \frac{1}{\sqrt{2\pi\sigma_f^2}} \exp\left(-\frac{(y+1)^2}{2\sigma_f^2}\right) + (1 - p_f) \frac{1}{\sqrt{2\pi\sigma_f^2}} \exp\left(-\frac{(y-1)^2}{2\sigma_f^2}\right)$$

$$P_m(x, 0) = p_m \frac{1}{\sqrt{2\pi\sigma_m^2}} \exp\left(-\frac{(x+1)^2}{2\sigma_m^2}\right) + (1 - p_m) \frac{1}{\sqrt{2\pi\sigma_m^2}} \exp\left(-\frac{(x-1)^2}{2\sigma_m^2}\right)$$

Even with a continuous distribution of female preferences, we find that after several generations, the male song distribution is concentrated in several discrete peaks (Figure ??). (Parameters and initial conditions given in Code ??.)

There are three critical parameters:

Mutations give rise to a small number of males with song traits that are slightly higher (lower) than the rest of the population. The “edge” females that most prefer these extreme songs have limited options because they do not like the songs most males are singing. They therefore mate with the new “edge” males with very high probability, which leads to a very fast increase of these songs. As mutations continue to generate new songs, extreme songs will be selected for at the expense of similar songs closer to the middle of the song range. Ultimately, this results in a number of discrete songs at high density.

Ignoring mutations for the moment, the rate of change of the probability density of a particular song is given by

$$\frac{P_m(x, t + 1)}{P_m(x, t)} = \int_{\mathbb{R}} \frac{P_f(y)f_y(x)}{Z_y} dy \quad (1)$$

This shows that the normalizing factor  $Z_y$  affects how the density of male songs changes from one generation to the next: if a male is recognized by females with very few options (small  $Z_y$ ), then the density of that song will increase more than the song a male recognized by females with many options (large  $Z_y$ ) (Figure ??).

## 4.2 Analysis

**Claim 4.1**  $\exp\left(-\frac{(x-\mu_a)^2}{2\sigma_a^2}\right) \times \exp\left(-\frac{(x-\mu_b)^2}{2\sigma_b^2}\right) = \exp\left(-\frac{(x-\mu_{ab})^2}{2\sigma_{ab}^2}\right) \times \exp\left(-\frac{(\mu_a-\mu_b)^2}{2(\sigma_a^2+\sigma_b^2)}\right)$  where  $\mu_{ab} = \frac{\mu_a\sigma_b^2+\mu_b\sigma_a^2}{\sigma_a^2+\sigma_b^2}$  and  $\sigma_{ab}^2 = \frac{\sigma_a^2\sigma_b^2}{\sigma_a^2+\sigma_b^2}$

$$\begin{aligned}
P_m(x)f_y(x) &= \frac{1}{\sqrt{2\pi\sigma_m^2}} \exp\left(-\frac{(x-\mu_m)^2}{2\sigma_m^2}\right) \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-y)^2}{2\sigma^2}\right) \\
&= \frac{1}{\sqrt{2\pi\sigma_m^2\sigma^2}} \exp\left(-\frac{(x-\frac{\mu_m\sigma^2+y\sigma_m^2}{\sigma_m^2+\sigma^2})^2}{2\frac{\sigma_m^2\sigma^2}{\sigma_m^2+\sigma^2}}\right) \exp\left(-\frac{(y-\mu_m)^2}{2(\sigma_m^2+\sigma^2)}\right) \\
&\quad \text{where } \mu_{ab} = \frac{\mu_m\sigma^2+y\sigma_m^2}{\sigma_m^2+\sigma^2} \text{ and } \sigma_{ab}^2 = \frac{\sigma_m^2\sigma^2}{\sigma_m^2+\sigma^2} \\
&= \frac{1}{\sqrt{2\pi(\sigma_m^2+\sigma^2)}} \exp\left(-\frac{(y-\mu_m)^2}{2(\sigma_m^2+\sigma^2)}\right) \frac{1}{\sqrt{2\pi\frac{\sigma_m^2\sigma^2}{\sigma_m^2+\sigma^2}}} \exp\left(-\frac{(x-\frac{\mu_m\sigma^2+y\sigma_m^2}{\sigma_m^2+\sigma^2})^2}{2\frac{\sigma_m^2\sigma^2}{\sigma_m^2+\sigma^2}}\right) \\
\Rightarrow Z_y &= \int_{\mathbb{R}} P_m(x)f_y(x)dx = \frac{1}{\sqrt{2\pi(\sigma_m^2+\sigma^2)}} \exp\left(-\frac{(y-\mu_m)^2}{2(\sigma_m^2+\sigma^2)}\right) \\
&\Rightarrow \frac{P_f(y)P_m(x)f_y(x)}{Z_y} = \frac{1}{\sqrt{2\pi\sigma_f^2}} \exp\left(-\frac{(y-\mu_f)^2}{2\sigma_f^2}\right) \frac{1}{\sqrt{2\pi\frac{\sigma_m^2\sigma^2}{\sigma_m^2+\sigma^2}}} \exp\left(-\frac{(x-\frac{\mu_m\sigma^2+y\sigma_m^2}{\sigma_m^2+\sigma^2})^2}{2\frac{\sigma_m^2\sigma^2}{\sigma_m^2+\sigma^2}}\right) \\
&\quad x - \frac{\mu_m\sigma^2+y\sigma_m^2}{\sigma_m^2+\sigma^2} = \frac{(\sigma_m^2+\sigma^2)x - \mu_m\sigma^2 - y\sigma_m^2}{\sigma_m^2+\sigma^2} = \frac{\frac{\sigma_m^2+\sigma^2}{\sigma_m^2}x - \frac{\sigma_m^2}{\sigma_m^2}\mu_m - y}{\frac{\sigma_m^2+\sigma^2}{\sigma_m^2}} \\
&\Rightarrow \frac{(x - \frac{\mu_m\sigma^2+y\sigma_m^2}{\sigma_m^2+\sigma^2})^2}{\frac{\sigma_m^2\sigma^2}{\sigma_m^2+\sigma^2}} = \frac{\left(y - \left(\frac{\sigma_m^2+\sigma^2}{\sigma_m^2}x - \frac{\sigma_m^2}{\sigma_m^2}\mu_m\right)\right)^2}{(\sigma_m^2+\sigma^2)^2} \times \frac{\sigma_m^4}{\sigma_m^2\sigma^2} = \frac{\left(y - \left(\frac{\sigma_m^2+\sigma^2}{\sigma_m^2}x - \frac{\sigma_m^2}{\sigma_m^2}\mu_m\right)\right)^2}{(\sigma_m^2+\sigma^2)\frac{\sigma_m^2}{\sigma_m^2}}
\end{aligned}$$

$$\begin{aligned}
&\Rightarrow \frac{P_f(y)P_m(x)f_y(x)}{Z_y} = \frac{1}{\sqrt{2\pi\sigma_f^2}} \exp\left(-\frac{(y-\mu_f)^2}{2\sigma_f^2}\right) \frac{1}{\sqrt{2\pi\frac{\sigma_m^2\sigma^2}{\sigma_m^2+\sigma^2}}} \exp\left(-\frac{\left(y-\left(\frac{\sigma_m^2+\sigma^2}{\sigma_m^2}x-\frac{\sigma_m^2}{\sigma_m^2}\mu_m\right)\right)^2}{2\frac{(\sigma_m^2+\sigma^2)\sigma^2}{\sigma_m^2}}\right) \\
&\mu_{ab} = \frac{\mu_a\sigma_b^2 + \mu_b\sigma_a^2}{\sigma_a^2 + \sigma_b^2}, \sigma_{ab}^2 = \frac{\sigma_a^2\sigma_b^2}{\sigma_a^2 + \sigma_b^2} \\
&= \frac{1}{\sqrt{2\pi\sigma_f^2}} \frac{1}{\sqrt{2\pi\frac{\sigma_m^2\sigma^2}{\sigma_m^2+\sigma^2}}} \exp\left(-\frac{(y-\mu_f^*)^2}{2\sigma_f^{*2}}\right) \exp\left(-\frac{\left(\frac{\sigma_m^2+\sigma^2}{\sigma_m^2}x-\frac{\sigma_m^2}{\sigma_m^2}\mu_m-\mu_f\right)^2}{2\left(\sigma_f^2+\frac{(\sigma_m^2+\sigma^2)\sigma^2}{\sigma_m^2}\right)}\right) \\
&\text{where } \mu_f^* = \frac{\mu_f\sigma^2(\sigma_m^2+\sigma^2) + \sigma_f^2(\sigma_m^2+\sigma^2)x - \sigma_f^2\sigma^2\mu_m}{\sigma_f^2\sigma_m^2 + (\sigma_m^2+\sigma^2)\sigma^2} \text{ and } \sigma_f^{*2} = \frac{\sigma_f^2(\sigma_m^2+\sigma^2)\sigma^2}{\sigma_m^2\sigma_f^2 + (\sigma_m^2+\sigma^2)\sigma^2} \\
&= \frac{1}{\sqrt{2\pi\sigma_f^2}} \frac{1}{\sqrt{2\pi\frac{\sigma_m^2\sigma^2}{\sigma_m^2+\sigma^2}}} \exp\left(-\frac{(y-\mu_f^*)^2}{2\sigma_f^{*2}}\right) \exp\left(-\frac{\left(x-\frac{\sigma_m^2}{\sigma_m^2+\sigma^2}\mu_m-\frac{\sigma_m^2}{\sigma_m^2+\sigma^2}\mu_f\right)^2}{2\frac{\sigma_m^2(\sigma_m^2\sigma_f^2+(\sigma_m^2+\sigma^2)\sigma^2)}{(\sigma_m^2+\sigma^2)^2}}\right) \\
&= \frac{1}{\sqrt{2\pi}} \frac{\sqrt{\sigma_m^2+\sigma^2}}{\sqrt{\sigma_f^2\sigma_m^2\sigma^2}} \frac{\sqrt{\sigma_f^2(\sigma_m^2+\sigma^2)\sigma^2}}{\sqrt{\sigma_m^2\sigma_f^2+(\sigma_m^2+\sigma^2)\sigma^2}} \frac{1}{\sqrt{2\pi\sigma_f^{*2}}} \exp\left(-\frac{(y-\mu_f^*)^2}{2\sigma_f^{*2}}\right) \exp\left(-\frac{\left(x-\frac{\sigma_m^2}{\sigma_m^2+\sigma^2}\mu_m-\frac{\sigma_m^2}{\sigma_m^2+\sigma^2}\mu_f\right)^2}{2\frac{\sigma_m^2(\sigma_m^2\sigma_f^2+(\sigma_m^2+\sigma^2)\sigma^2)}{(\sigma_m^2+\sigma^2)^2}}\right) \\
&= \frac{1}{\sqrt{2\pi}} \frac{\sqrt{(\sigma_m^2+\sigma^2)^2}}{\sqrt{\sigma_m^2(\sigma_m^2\sigma_f^2+(\sigma_m^2+\sigma^2)\sigma^2)}} \frac{1}{\sqrt{2\pi\sigma_f^{*2}}} \exp\left(-\frac{(y-\mu_f^*)^2}{2\sigma_f^{*2}}\right) \exp\left(-\frac{\left(x-\frac{\sigma_m^2}{\sigma_m^2+\sigma^2}\mu_m-\frac{\sigma_m^2}{\sigma_m^2+\sigma^2}\mu_f\right)^2}{2\frac{\sigma_m^2(\sigma_m^2\sigma_f^2+(\sigma_m^2+\sigma^2)\sigma^2)}{(\sigma_m^2+\sigma^2)^2}}\right) \\
&\Rightarrow \int_{\mathbb{R}} \frac{P_f(y)P_m(x)f_y(x)}{Z_y} dy = \frac{1}{\sqrt{2\pi}} \frac{\sqrt{(\sigma_m^2+\sigma^2)^2}}{\sqrt{\sigma_m^2(\sigma_m^2\sigma_f^2+(\sigma_m^2+\sigma^2)\sigma^2)}} \exp\left(-\frac{\left(x-\frac{\sigma_m^2\mu_m+\sigma_m^2\mu_f}{\sigma_m^2+\sigma^2}\right)^2}{2\frac{\sigma_m^2(\sigma_m^2\sigma_f^2+(\sigma_m^2+\sigma^2)\sigma^2)}{(\sigma_m^2+\sigma^2)^2}}\right)
\end{aligned}$$

OR =

$$\begin{aligned}
\frac{P_{\mathbf{f}}(y)f_y(x)}{Z_y} &= \frac{\sqrt{\sigma_{\mathbf{m}}^2 + \sigma^2}}{\sqrt{2\pi\sigma_{\mathbf{f}}^2\sigma^2}} \exp\left(-\frac{(y-\mu_{\mathbf{f}})^2}{2\sigma_{\mathbf{f}}^2}\right) \exp\left(-\frac{(y-x)^2}{2\sigma^2}\right) \exp\left(-\frac{(y-\mu_{\mathbf{m}})^2}{2(-1)(\sigma_{\mathbf{m}}^2 + \sigma^2)}\right) \\
&= \frac{\sqrt{\sigma_{\mathbf{m}}^2 + \sigma^2}}{\sqrt{2\pi\sigma_{\mathbf{f}}^2\sigma^2}} \exp\left(-\frac{\left(y - \frac{-\mu_{\mathbf{f}}(\sigma_{\mathbf{m}}^2 + \sigma^2) + \mu_{\mathbf{m}}\sigma_{\mathbf{f}}^2}{\sigma_{\mathbf{f}}^2 - (\sigma_{\mathbf{m}}^2 + \sigma^2)}\right)^2}{2\frac{-\sigma_{\mathbf{f}}^2(\sigma_{\mathbf{m}}^2 + \sigma^2)}{\sigma_{\mathbf{f}}^2 - (\sigma_{\mathbf{m}}^2 + \sigma^2)}}\right) \exp\left(-\frac{(\mu_{\mathbf{m}} - \mu_{\mathbf{f}})^2}{2(\sigma_{\mathbf{f}}^2 - (\sigma_{\mathbf{m}}^2 + \sigma^2))}\right) \\
&= \frac{\sqrt{\sigma_{\mathbf{m}}^2 + \sigma^2}}{\sqrt{2\pi\sigma_{\mathbf{f}}^2\sigma^2}} \exp\left(-\frac{\left(y - \frac{-\mu_{\mathbf{f}}(\sigma_{\mathbf{m}}^2 + \sigma^2) + \mu_{\mathbf{m}}\sigma_{\mathbf{f}}^2}{\sigma_{\mathbf{f}}^2 - (\sigma_{\mathbf{m}}^2 + \sigma^2)}\right)^2}{2\frac{\sigma_{\mathbf{f}}^2(\sigma_{\mathbf{m}}^2 + \sigma^2)}{\sigma_{\mathbf{m}}^2 + \sigma^2 - \sigma_{\mathbf{f}}^2}}\right) \exp\left(-\frac{(\mu_{\mathbf{m}} - \mu_{\mathbf{f}})^2}{2(\sigma_{\mathbf{f}}^2 - (\sigma_{\mathbf{m}}^2 + \sigma^2))}\right) \\
&= \frac{\sqrt{\sigma_{\mathbf{m}}^2 + \sigma^2}}{\sqrt{2\pi\sigma_{\mathbf{f}}^2\sigma^2}} \exp\left(-\frac{(y - \bar{\mu}_{\mathbf{f}})^2}{2\frac{\sigma_{\mathbf{f}}^2\sigma_{\mathbf{m}}^2(\sigma_{\mathbf{m}}^2 + \sigma^2)}{\sigma_{\mathbf{f}}^2(\sigma_{\mathbf{m}}^2 + \sigma^2) + \sigma^2(\sigma_{\mathbf{m}}^2 + \sigma^2 - \sigma_{\mathbf{f}}^2)}}\right) \exp\left(-\frac{(\mu_{\mathbf{m}} - \mu_{\mathbf{f}})^2}{2(\sigma_{\mathbf{f}}^2 - (\sigma_{\mathbf{m}}^2 + \sigma^2))}\right) \\
&= \frac{\sqrt{\sigma_{\mathbf{m}}^2 + \sigma^2}}{\sqrt{2\pi\sigma_{\mathbf{f}}^2\sigma^2}} \exp\left(-\frac{(y - \bar{\mu}_{\mathbf{f}})^2}{2\left(\frac{\sigma_{\mathbf{f}}^2\sigma_{\mathbf{m}}^2(\sigma_{\mathbf{m}}^2 + \sigma^2)}{\sigma_{\mathbf{f}}^2(\sigma_{\mathbf{m}}^2 + \sigma^2) + \sigma^2(\sigma_{\mathbf{m}}^2 + \sigma^2 - \sigma_{\mathbf{f}}^2)}\right)}\right) \exp\left(-\frac{(\mu_{\mathbf{m}} - \mu_{\mathbf{f}})^2}{2(\sigma_{\mathbf{f}}^2 - (\sigma_{\mathbf{m}}^2 + \sigma^2))}\right) \\
&= \frac{\sqrt{\sigma_{\mathbf{m}}^2 + \sigma^2}}{\sqrt{2\pi\sigma_{\mathbf{f}}^2\sigma^2}} \exp\left(-\frac{(y - \bar{\mu}_{\mathbf{f}})^2}{2\frac{\sigma_{\mathbf{f}}^2\sigma_{\mathbf{m}}^2(\sigma_{\mathbf{m}}^2 + \sigma^2)}{\sigma_{\mathbf{f}}^2\sigma_{\mathbf{m}}^2 + \sigma^2(\sigma_{\mathbf{m}}^2 + \sigma^2 - \sigma_{\mathbf{f}}^2)}}\right) \exp\left(-\frac{(\mu_{\mathbf{m}} - \mu_{\mathbf{f}})^2}{2(\sigma_{\mathbf{f}}^2 - (\sigma_{\mathbf{m}}^2 + \sigma^2))}\right) \\
&\Rightarrow \int_{\mathbb{R}} \frac{P_{\mathbf{f}}(y)f_y(x)}{Z_y} dy = \frac{\sqrt{(\sigma_{\mathbf{m}}^2 + \sigma^2)^2}}{\sqrt{\sigma_{\mathbf{f}}^2\sigma_{\mathbf{m}}^2 + \sigma^2(\sigma_{\mathbf{m}}^2 + \sigma^2)}} \exp\left(-\frac{\left(x - \frac{\mu_{\mathbf{f}}(\sigma_{\mathbf{m}}^2 + \sigma^2) - \mu_{\mathbf{m}}\sigma_{\mathbf{f}}^2}{\sigma_{\mathbf{m}}^2 + \sigma^2 - \sigma_{\mathbf{f}}^2}\right)^2}{2\left(\frac{\sigma^2(\sigma_{\mathbf{m}}^2 + \sigma^2) + \sigma_{\mathbf{f}}^2\sigma_{\mathbf{m}}^2}{\sigma_{\mathbf{m}}^2 + \sigma^2 - \sigma_{\mathbf{f}}^2}\right)}\right) \exp\left(-\frac{(\mu_{\mathbf{m}} - \mu_{\mathbf{f}})^2}{2(\sigma_{\mathbf{f}}^2 - (\sigma_{\mathbf{m}}^2 + \sigma^2))}\right)
\end{aligned}$$



$$\begin{aligned}
P_m(x) \int_{\mathbb{R}} \frac{P_f(y)f_y(x)}{Z_y} dy &= \frac{\sqrt{(\sigma_m^2 + \sigma^2)^2}}{\sqrt{2\pi\sigma_m^2(\sigma_f^2\sigma_m^2 + \sigma^2(\sigma_m^2 + \sigma^2))}} \exp\left(-\frac{(x - \mu_m)^2}{2\sigma_m^2}\right) \exp\left(-\frac{\left(x - \frac{\mu_f(\sigma_m^2 + \sigma^2) - \mu_m\sigma_f^2}{\sigma_m^2 + \sigma^2 - \sigma_f^2}\right)^2}{2\left(\frac{\sigma_m^2(\sigma_m^2 + \sigma^2) + \sigma_f^2\sigma_m^2}{\sigma_m^2 + \sigma^2 - \sigma_f^2}\right)}\right) \\
&= \frac{\sqrt{(\sigma_m^2 + \sigma^2)^2}}{\sqrt{2\pi\sigma_m^2(\sigma_f^2\sigma_m^2 + \sigma^2(\sigma_m^2 + \sigma^2))}} \exp\left(-\frac{(x - \mu'_m)^2}{x - \sigma_m'^2}\right) \exp\left(-\frac{\left(\mu_m - \frac{\mu_f(\sigma_m^2 + \sigma^2) - \mu_m\sigma_f^2}{\sigma_m^2 + \sigma^2 - \sigma_f^2}\right)^2}{2\left(\sigma_m^2 + \frac{\sigma_m^2(\sigma_m^2 + \sigma^2) - \sigma_f^2\sigma_m^2}{\sigma_m^2 + \sigma^2 - \sigma_f^2}\right)}\right) \\
&= \frac{\sqrt{(\sigma_m^2 + \sigma^2)^2}}{\sqrt{2\pi\sigma_m^2(\sigma_f^2\sigma_m^2 + \sigma^2(\sigma_m^2 + \sigma^2))}} \exp\left(-\frac{(x - \mu'_m)^2}{x - \sigma_m'^2}\right) \exp\left(-\frac{\left(\frac{\mu_m(\sigma_m^2 + \sigma^2) - \mu_f(\sigma_m^2 + \sigma^2)}{\sigma_m^2 + \sigma^2 - \sigma_f^2}\right)^2}{2\left(\frac{(\sigma_m^2 + \sigma^2)^2}{\sigma_m^2 + \sigma^2 - \sigma_f^2}\right)}\right) \\
&= \frac{\sqrt{(\sigma_m^2 + \sigma^2)^2}}{\sqrt{2\pi\sigma_m^2(\sigma_f^2\sigma_m^2 + \sigma^2(\sigma_m^2 + \sigma^2))}} \exp\left(-\frac{(x - \mu'_m)^2}{x - \sigma_m'^2}\right) \exp\left(-\frac{(\mu_m - \mu_f)^2}{2(\sigma_f^2 - (\sigma_m^2 + \sigma^2))}\right) \\
&= \frac{\sqrt{(\sigma_m^2 + \sigma^2)^2}}{\sqrt{2\pi\sigma_m^2(\sigma_f^2\sigma_m^2 + \sigma^2(\sigma_m^2 + \sigma^2))}} \exp\left(-\frac{(x - \mu'_m)^2}{x - \sigma_m'^2}\right) \exp\left(-\frac{(\mu_m - \mu_f)^2}{2(\sigma_m^2 + \sigma^2 - \sigma_f^2)}\right) \\
&= \frac{\sqrt{(\sigma_m^2 + \sigma^2)^2}}{\sqrt{2\pi\sigma_m^2(\sigma_f^2\sigma_m^2 + \sigma^2(\sigma_m^2 + \sigma^2))}} \exp\left(-\frac{(x - \mu'_m)^2}{x - \sigma_m'^2}\right) \exp\left(-\frac{(\mu_m - \mu_f)^2}{2(\sigma_m^2 + \sigma^2 - \sigma_f^2)}\right) \\
&= \frac{\sqrt{(\sigma_m^2 + \sigma^2)^2}}{\sqrt{2\pi\sigma_m^2(\sigma_f^2\sigma_m^2 + \sigma^2(\sigma_m^2 + \sigma^2))}} \exp\left(-\frac{(x - \mu'_m)^2}{x - \sigma_m'^2}\right) \exp\left(-\frac{(\mu_m - \mu_f)^2}{2(\sigma_m^2 + \sigma^2 - \sigma_f^2)}\right) \\
&= \frac{\mu_m}{\mu_m} \frac{\left(\frac{\sigma^2(\sigma_m^2 + \sigma^2) + \sigma_f^2\sigma_m^2}{\sigma_m^2 + \sigma^2 - \sigma_f^2}\right) + \left(\frac{\mu_f(\sigma_m^2 + \sigma^2) - \mu_m\sigma_f^2}{\sigma_m^2 + \sigma^2 - \sigma_f^2}\right) \sigma_m^2}{\sigma_m^2 + \frac{\sigma^2(\sigma_m^2 + \sigma^2) + \sigma_f^2\sigma_m^2}{\sigma_m^2 + \sigma^2 - \sigma_f^2}} \\
&= \frac{\mu_m\sigma^2(\sigma_m^2 + \sigma^2) + \mu_f\sigma_m^2(\sigma_m^2 + \sigma^2)}{(\sigma_m^2 + \sigma^2)^2} = \frac{\mu_m\sigma^2 + \mu_f\sigma_m^2}{\sigma_m^2 + \sigma^2} \\
&= \frac{\mu_m}{\mu_m} \frac{\frac{\sigma^2(\sigma_m^2 + \sigma^2) + \sigma_f^2\sigma_m^2}{\sigma_m^2 + \sigma^2 - \sigma_f^2}}{\mu_m + \frac{\sigma^2(\sigma_m^2 + \sigma^2) + \sigma_f^2\sigma_m^2}{\sigma_m^2 + \sigma^2 - \sigma_f^2}} \\
&= \frac{\mu_m(\sigma_f^2\sigma_m^2 + \sigma^2(\sigma_m^2 + \sigma^2))}{(\sigma_m^2 + \sigma^2)^2} \\
\Rightarrow P_m(x) \int_{\mathbb{R}} \frac{P_f(y)f_y(x)}{Z_y} dy &= \frac{1}{\sqrt{2\pi\sigma_m'^2}} \exp\left(-\frac{(x - \mu'_m)^2}{2\sigma_m'^2}\right) \text{ and } \int_{\mathbb{R}} \frac{P_f(y)P_m(x)f_y(x)}{Z_y} dx = P_f(y)
\end{aligned}$$

### 4.3 Multivariate model



$$P_m(x, y) \sim N \left( \begin{pmatrix} \mu_{mx} \\ \mu_{my} \end{pmatrix}, \begin{pmatrix} \sigma_{mx}^2 & \rho_m \sigma_{mx} \sigma_{my} \\ \rho_m \sigma_{mx} \sigma_{my} & \sigma_{my}^2 \end{pmatrix} \right)$$

$$P_f(x, y) \sim N \left( \begin{pmatrix} \mu_{fx} \\ \mu_{fy} \end{pmatrix}, \begin{pmatrix} \sigma_{fx}^2 & \rho_f \sigma_{fx} \sigma_{fy} \\ \rho_f \sigma_{fx} \sigma_{fy} & \sigma_{fy}^2 \end{pmatrix} \right)$$

$$), (x_2, y_2)) = \frac{P_f(x_2, y_2) P_m(x_1, y_1) f_{y_2}(x_1)}{\int_{\mathbb{R}} \int_{\mathbb{R}} P_m(x, y) f_{y_2}(x) dx dy}$$

$$\begin{aligned} P_m(x_1, y_1) &= \frac{1}{2\pi \sqrt{(1 - \rho_m)^2 \sigma_{mx}^2 \sigma_{my}^2}} \exp \left( -\frac{(x_1 - (\mu_{mx} + \frac{\rho_m \sigma_{mx}}{\sigma_{my}}(y_1 - \mu_{my})))^2}{2(1 - \rho_m^2) \sigma_{mx}^2} \right) \exp \left( -\frac{(y_1 - \mu_{my})^2}{2\sigma_{my}^2} \right) \\ &= \frac{1}{\sqrt{2\pi(1 - \rho_m)^2 \sigma_{mx}^2}} \exp \left( -\frac{(x_1 - (\mu_{mx} + \frac{\rho_m \sigma_{mx}}{\sigma_{my}}(y_1 - \mu_{my})))^2}{2(1 - \rho_m^2) \sigma_{mx}^2} \right) \frac{1}{\sqrt{2\pi \sigma_{my}^2}} \exp \left( -\frac{(y_1 - \mu_{my})^2}{2\sigma_{my}^2} \right) \end{aligned}$$

$$\mu'_{mx} = \mu_{mx} + \frac{\rho_m \sigma_{mx}}{\sigma_{my}}(y_1 - \mu_{my}) \text{ and } \sigma'^2_{mx} = (1 - \rho_m^2) \sigma_{mx}^2$$

$$y_1) f_{y_2}(x_1) = \frac{1}{\sqrt{2\pi(\sigma^2 + \sigma_{mx}^2)}} \exp \left( -\frac{(y_2 - \mu'_{mx})^2}{2(\sigma^2 + \sigma_{mx}^2)} \right) \frac{1}{\sqrt{2\pi \frac{\sigma_{mx}^2 \sigma^2}{\sigma_{mx}^2 + \sigma^2}}} \exp \left( -\frac{(x_1 - \frac{\mu'_{mx} \sigma^2 + y_2 \sigma_{mx}^2}{\sigma_{mx}^2 + \sigma^2})^2}{2\pi \sigma_{my}^2} \right) \frac{1}{\sqrt{2\pi \sigma_{my}^2}} \exp \left( -\frac{(y_1 - \mu_{my})^2}{2\sigma_{my}^2} \right)$$

$$y_2) dx dy = \frac{1}{\sqrt{2\pi(\sigma^2 + \sigma_{mx}^2)}} \exp \left( -\frac{(y_2 - \mu'_{mx})^2}{2(\sigma^2 + \sigma_{mx}^2)} \right)$$

$$P_f(x_2, y_2) = \frac{1}{\sqrt{2\pi(1 - \rho_f)^2 \sigma_{fy}^2}} \exp \left( -\frac{(y_2 - (\mu_{fy} + \frac{\rho_f \sigma_{fy}}{\sigma_{fx}}(x_2 - \mu_{fx})))^2}{2(1 - \rho_f^2) \sigma_{fy}^2} \right) \frac{1}{\sqrt{2\pi \sigma_{fx}^2}} \exp \left( -\frac{(x_2 - \mu_{fx})^2}{2\sigma_{fx}^2} \right)$$

$$\mu'_{fy} = \mu_{fy} + \frac{\rho_f \sigma_{fy}}{\sigma_{fx}}(x_2 - \mu_{fx}) \text{ and } \sigma'^2_{fy} = (1 - \rho_f^2) \sigma_{fy}^2$$

$$\begin{aligned} ), (x_2, y_2)) &= \frac{1}{\sqrt{2\pi \sigma_{fy}^2}} \exp \left( -\frac{(y_2 - \mu'_{fy})^2}{2\sigma_{fy}^2} \right) \frac{1}{\sqrt{2\pi \sigma_{fx}^2}} \exp \left( -\frac{(x_2 - \mu_{fx})^2}{2\sigma_{fx}^2} \right) \frac{1}{\sqrt{2\pi \frac{\sigma_{fx}^2 \sigma^2}{\sigma_{fx}^2 + \sigma^2}}} \exp \left( -\frac{(x_1 - \frac{\mu'_{fx} \sigma^2 + y_2 \sigma_{fx}^2}{\sigma_{fx}^2 + \sigma^2})^2}{2\pi \sigma_{my}^2} \right) \frac{1}{\sqrt{2\pi \sigma_{my}^2}} \exp \left( -\frac{(y_1 - \mu_{my})^2}{2\sigma_{my}^2} \right) \\ &= \frac{1}{\sqrt{2\pi \sigma_{fy}^2}} \exp \left( -\frac{(y_2 - \mu'_{fy})^2}{2\sigma_{fy}^2} \right) \frac{1}{\sqrt{2\pi \sigma_{fx}^2}} \exp \left( -\frac{(x_2 - \mu_{fx})^2}{2\sigma_{fx}^2} \right) \frac{1}{\sqrt{2\pi \frac{\sigma_{fx}^2 \sigma^2}{\sigma_{fx}^2 + \sigma^2}}} \exp \left( -\frac{(y_2 - (\frac{\sigma'^2_{mx} + \sigma^2}{\sigma_{mx}^2} x_1 - \frac{\sigma^2}{\sigma_{mx}^2} \mu'_{mx}))^2}{2 \frac{(\sigma'^2_{mx} + \sigma^2) \sigma^2}{\sigma_{mx}^2}} \right) \frac{1}{\sqrt{2\pi \sigma_{my}^2}} \exp \left( -\frac{(y_1 - \mu_{my})^2}{2\sigma_{my}^2} \right) \end{aligned}$$

$$\mu''_{fy} = \frac{\mu'_{fy} \left( \frac{(\sigma'^2_{mx} + \sigma^2) \sigma^2}{\sigma_{mx}^2} \right)}{\left( \frac{(\sigma'^2_{mx} + \sigma^2) \sigma^2}{\sigma_{mx}^2} + \sigma_{fx}^2 \right) + \left( \frac{\sigma'^2_{mx} + \sigma^2}{\sigma_{mx}^2} x_1 - \frac{\sigma^2}{\sigma_{mx}^2} \mu'_{mx} \right) \sigma_{fy}^2} \text{ and } \sigma'^2_{fy} = \frac{\sigma'^2_{fy} \sigma^2 (\sigma_{mx}^2 + \sigma^2)}{\sigma_{fy}^2 \sigma_{mx}^2 + (\sigma'^2_{mx} + \sigma^2) \sigma^2}$$

$$\begin{aligned}
& \frac{(y_2 - \mu''_{ty})^2}{2\sigma_{ty}''^2} \exp\left(-\frac{(x_2 - \mu_{tx})^2}{2\sigma_{tx}^2}\right) \frac{1}{\sqrt{2\pi\sigma_{tx}^2}} \exp\left(-\frac{(x_2 - \mu_{tx})^2}{2\sigma_{tx}^2}\right) \frac{1}{\sqrt{2\pi\frac{\sigma_{tx}'^2\sigma^2}{\sigma_{tx}^2} + \sigma^2}} \exp\left(-\frac{\left(-\frac{\frac{\sigma_{tx}'^2 + \sigma^2}{\sigma_{tx}^2}x_1 - \frac{\sigma_{tx}'^2}{\sigma_{tx}^2}\mu'_{mx} - \mu'_{ty}\right)^2}{2\left(\sigma_{ty}^{\prime 2} + \frac{(\sigma_{tx}'^2 + \sigma^2)\sigma^2}{\sigma_{tx}^2}\right)}\right) \frac{1}{\sqrt{2\pi\sigma_{my}^2}} \exp\left(-\frac{(y_1 - \mu_{my})^2}{2\sigma_{my}^2}\right) \\
& \frac{(y_2 - \mu''_{ty})^2}{2\sigma_{ty}''^2} \exp\left(-\frac{(x_2 - \mu_{tx})^2}{2\sigma_{tx}^2}\right) \frac{1}{\sqrt{2\pi\frac{\sigma_{tx}'^2\sigma^2}{\sigma_{tx}^2} + \sigma^2}} \exp\left(-\frac{\left(x_1 - \frac{\sigma^2}{\sigma_{tx}'^2 + \sigma^2}\mu'_{mx} - \frac{\sigma_{tx}'^2}{\sigma_{tx}'^2 + \sigma^2}\mu'_{ty}\right)^2}{2\frac{\sigma_{tx}'^2(\sigma_{ty}^{\prime 2}\sigma_{tx}^{\prime 2} + (\sigma_{tx}'^2 + \sigma^2)\sigma^2)}{(\sigma_{tx}^{\prime 2} + \sigma^2)^2}}\right) \frac{1}{\sqrt{2\pi\sigma_{my}^2}} \exp\left(-\frac{(y_1 - \mu_{my})^2}{2\sigma_{my}^2}\right) \\
& \frac{(\sigma^2)}{\sigma^2} \exp\left(-\frac{(y_2 - \mu''_{ty})^2}{2\sigma_{ty}^{\prime 2}}\right) \frac{1}{\sqrt{2\pi\sigma_{tx}^2}} \exp\left(-\frac{(x_2 - \mu_{tx})^2}{2\sigma_{tx}^2}\right) \frac{1}{\sqrt{2\pi\frac{\sigma_{tx}'^2(\sigma_{ty}^{\prime 2}\sigma_{tx}^{\prime 2} + (\sigma_{tx}'^2 + \sigma^2)\sigma^2)}{(\sigma_{tx}^{\prime 2} + \sigma^2)^2}}} \exp\left(-\frac{\left(x_1 - \frac{\sigma^2}{\sigma_{tx}'^2 + \sigma^2}\mu'_{mx} - \frac{\sigma_{tx}'^2}{\sigma_{tx}'^2 + \sigma^2}\mu'_{ty}\right)^2}{2\frac{\sigma_{tx}'^2(\sigma_{ty}^{\prime 2}\sigma_{tx}^{\prime 2} + (\sigma_{tx}'^2 + \sigma^2)\sigma^2)}{(\sigma_{tx}^{\prime 2} + \sigma^2)^2}}\right) \frac{1}{\sqrt{2\pi\sigma_{my}^2}} \exp\left(-\frac{(y_1 - \mu_{my})^2}{2\sigma_{my}^2}\right)
\end{aligned}$$

$$\begin{aligned}
P_{\text{mate}}((x_1, y_1), (x_2, y_2)) &= \frac{1}{\sqrt{2\pi\sigma_{\text{mx}}'^2}} \exp\left(-\frac{(x_1 - \mu_{\text{mx}}'')^2}{2\sigma_{\text{mx}}'^2}\right) \frac{1}{\sqrt{2\pi\sigma_{\text{my}}^2}} \exp\left(-\frac{(y_1 - \mu_{\text{my}})^2}{2\sigma_{\text{my}}^2}\right) \frac{1}{\sqrt{2\pi\sigma_{\text{tx}}'^2}} \exp\left(-\frac{(x_2 - \mu_{\text{tx}})^2}{2\sigma_{\text{tx}}'^2}\right) \frac{1}{\sqrt{2\pi\sigma_{\text{ty}}'^2}} \exp\left(-\frac{(y_2 - \mu_{\text{ty}}'')^2}{2\sigma_{\text{ty}}'^2}\right) \\
\mu_{\text{mx}}' &= \mu_{\text{mx}} + \frac{\rho_{\text{m}}\sigma_{\text{mx}}}{\sigma_{\text{my}}} (y_1 - \mu_{\text{my}}) \\
\sigma_{\text{mx}}'^2 &= (1 - \rho_{\text{m}}^2)\sigma_{\text{mx}}^2 \\
\mu_{\text{ty}}' &= \mu_{\text{ty}} + \frac{\rho_{\text{f}}\sigma_{\text{ty}}}{\sigma_{\text{tx}}} (x_2 - \mu_{\text{tx}}) \\
\sigma_{\text{ty}}'^2 &= (1 - \rho_{\text{f}}^2)\sigma_{\text{ty}}^2 \\
\mu_{\text{mx}}'' &= \frac{\sigma^2}{\sigma_{\text{mx}}'^2 + \sigma^2} \mu_{\text{mx}}' + \frac{\sigma_{\text{mx}}'^2}{\sigma_{\text{mx}}'^2 + \sigma^2} \mu_{\text{ty}}' = \frac{\sigma^2 \mu_{\text{mx}}' + \sigma_{\text{mx}}'^2 \mu_{\text{ty}}'}{\sigma_{\text{mx}}'^2 + \sigma^2} \\
&= \frac{\sigma^2}{\sigma_{\text{mx}}'^2 + \sigma^2} \mu_{\text{mx}} + \frac{\rho_{\text{m}}\sigma_{\text{mx}}\sigma^2}{\sigma_{\text{my}}(\sigma_{\text{mx}}'^2 + \sigma^2)} y_1 - \frac{\rho_{\text{m}}\sigma_{\text{mx}}\sigma^2}{\sigma_{\text{my}}(\sigma_{\text{mx}}'^2 + \sigma^2)} \mu_{\text{my}} + \frac{\sigma_{\text{mx}}'^2}{\sigma_{\text{mx}}'^2 + \sigma^2} \mu_{\text{ty}} + \frac{\rho_{\text{f}}\sigma_{\text{ty}}\sigma_{\text{mx}}'^2}{\sigma_{\text{tx}}(\sigma_{\text{mx}}'^2 + \sigma^2)} x_2 - \frac{\rho_{\text{f}}\sigma_{\text{ty}}\sigma_{\text{mx}}'^2}{\sigma_{\text{tx}}(\sigma_{\text{mx}}'^2 + \sigma^2)} \mu_{\text{tx}} \\
\sigma_{\text{mx}}''^2 &= \frac{\sigma_{\text{mx}}'^2 (\sigma_{\text{ty}}'^2 \sigma_{\text{mx}}'^2 + (\sigma_{\text{mx}}'^2 + \sigma^2) \sigma^2)}{(\sigma_{\text{mx}}'^2 + \sigma^2)^2} \\
\mu_{\text{ty}}'' &= \frac{\mu_{\text{ty}}' \left( \frac{(\sigma_{\text{mx}}'^2 + \sigma^2) \sigma^2}{\sigma_{\text{mx}}'^2} \right) + \left( \frac{\sigma_{\text{mx}}'^2 + \sigma^2}{\sigma_{\text{mx}}'^2} x_1 - \frac{\sigma^2}{\sigma_{\text{mx}}'^2} \mu_{\text{mx}}' \right) \sigma_{\text{ty}}'^2}{\frac{(\sigma_{\text{mx}}'^2 + \sigma^2) \sigma^2}{\sigma_{\text{mx}}'^2} + \sigma_{\text{ty}}'^2} = \frac{\mu_{\text{ty}}' \sigma^2 (\sigma_{\text{mx}}'^2 + \sigma^2) + ((\sigma_{\text{mx}}'^2 + \sigma^2) x_1 - \sigma^2 \mu_{\text{mx}}') \sigma_{\text{ty}}'^2}{(\sigma_{\text{mx}}'^2 + \sigma^2) \sigma^2 + \sigma_{\text{mx}}'^2 \sigma_{\text{ty}}'^2} \\
&= \frac{\sigma^2 (\sigma_{\text{mx}}'^2 + \sigma^2)}{(\sigma_{\text{mx}}'^2 + \sigma^2) \sigma^2 + \sigma_{\text{mx}}'^2 \sigma_{\text{ty}}'^2} \mu_{\text{ty}} + \frac{\rho_{\text{f}}\sigma_{\text{ty}}\sigma^2 (\sigma_{\text{mx}}'^2 + \sigma^2)}{\sigma_{\text{tx}}((\sigma_{\text{mx}}'^2 + \sigma^2) \sigma^2 + \sigma_{\text{mx}}'^2 \sigma_{\text{ty}}'^2)} x_2 - \frac{\rho_{\text{f}}\sigma_{\text{ty}}\sigma^2 (\sigma_{\text{mx}}'^2 + \sigma^2)}{\sigma_{\text{tx}}((\sigma_{\text{mx}}'^2 + \sigma^2) \sigma^2 + \sigma_{\text{mx}}'^2 \sigma_{\text{ty}}'^2)} \mu_{\text{tx}} + \\
&\quad \frac{\sigma_{\text{ty}}'^2 (\sigma_{\text{mx}}'^2 + \sigma^2)}{(\sigma_{\text{mx}}'^2 + \sigma^2) \sigma^2 + \sigma_{\text{mx}}'^2 \sigma_{\text{ty}}'^2} x_1 - \frac{\sigma^2 \sigma_{\text{ty}}'^2}{(\sigma_{\text{mx}}'^2 + \sigma^2) \sigma^2 + \sigma_{\text{mx}}'^2 \sigma_{\text{ty}}'^2} \mu_{\text{mx}} - \frac{\rho_{\text{m}}\sigma_{\text{mx}}\sigma^2 \sigma_{\text{ty}}'^2}{\sigma_{\text{my}}((\sigma_{\text{mx}}'^2 + \sigma^2) \sigma^2 + \sigma_{\text{mx}}'^2 \sigma_{\text{ty}}'^2)} y_1 + \frac{\rho_{\text{m}}\sigma_{\text{mx}}\sigma^2 \sigma_{\text{ty}}'^2}{\sigma_{\text{my}}((\sigma_{\text{mx}}'^2 + \sigma^2) \sigma^2 + \sigma_{\text{mx}}'^2 \sigma_{\text{ty}}'^2)} \mu_{\text{my}} \\
\sigma_{\text{ty}}''^2 &= \frac{\sigma_{\text{ty}}'^2 \sigma^2 (\sigma_{\text{mx}}'^2 + \sigma^2)}{\sigma_{\text{ty}}'^2 \sigma_{\text{mx}}'^2 + (\sigma_{\text{mx}}'^2 + \sigma^2) \sigma^2}
\end{aligned}$$

$$\begin{aligned} \text{es with } x_2 \text{ female)} &= \int \int P_{\text{mate}}((x_1, y_1), (x_2, y_2)) dy_1 dy_2 \\ &= \int \frac{1}{\sqrt{2\pi\sigma''^2_{mx}}} \exp\left(-\frac{(x_1 - \mu'_{mx})^2}{2\sigma'^2_{mx}}\right) \frac{1}{\sqrt{2\pi\sigma^2_{my}}} \exp\left(-\frac{(y_1 - \mu_{my})^2}{2\sigma^2_{my}}\right) \frac{1}{\sqrt{2\pi\sigma'^2_{fx}}} \exp\left(-\frac{(x_2 - \mu_{fx})^2}{2\sigma'^2_{fx}}\right) \\ \mu''_{mx} &= \frac{\sigma^2}{\sigma'^2_{mx} + \sigma^2} \mu_{mx} + \frac{\rho_m \sigma_{mx} \sigma^2}{\sigma_{my} (\sigma'^2_{mx} + \sigma^2)} y_1 - \frac{\rho_m \sigma_{mx} \sigma^2}{\sigma_{my} (\sigma'^2_{mx} + \sigma^2)} \mu_{my} + \frac{\sigma'^2_{mx}}{\sigma'^2_{mx} + \sigma^2} \mu_{fy} + \frac{\rho_t \sigma_{fy} \sigma'^2_{mx}}{\sigma_{tx} (\sigma'^2_{mx} + \sigma^2)} x_2 - \frac{\rho_t \sigma_{fy} \sigma'^2_{mx}}{\sigma_{tx} (\sigma'^2_{mx} + \sigma^2)} \mu_{fx} \\ &= \frac{\sigma^2}{\sigma'^2_{mx} + \sigma^2} \left( \mu_{mx} + \frac{\rho_m \sigma_{mx}}{\sigma_{my}} y_1 - \frac{\rho_m \sigma_{mx}}{\sigma_{my}} \mu_{my} + \frac{\sigma'^2_{mx}}{\sigma'^2_{mx} + \sigma^2} \mu_{fy} + \frac{\rho_t \sigma_{fy} \sigma'^2_{mx}}{\sigma_{tx} \sigma^2} x_2 - \frac{\rho_t \sigma_{fy} \sigma'^2_{mx}}{\sigma_{tx} \sigma^2} \mu_{fx} \right) \\ &= \frac{\sigma^2}{\sigma'^2_{mx} + \sigma^2} \frac{\rho_m \sigma_{mx}}{\sigma_{my}} \left( y_1 + \frac{\rho_t \sigma_{fy} \sigma_{my} \sigma'^2_{mx}}{\rho_m \sigma_{mx} \sigma_{fx} \sigma^2} x_2 + \frac{\sigma_{my}}{\rho_m \sigma_{mx}} \mu_{mx} - \mu_{my} - \frac{\rho_t \sigma_{fy} \sigma_{my} \sigma'^2_{mx}}{\rho_m \sigma_{mx} \sigma_{fy} \sigma^2} \mu_{fx} + \frac{\sigma_{my} \sigma'^2_{mx}}{\rho_m \sigma_{mx} \sigma^2} \mu_{fy} \right) \\ &\Rightarrow \frac{(x_1 - \mu''_{mx})^2}{\sigma''^2_{mx}} = \frac{\left( y_1 + \frac{\rho_t \sigma_{fy} \sigma_{my} \sigma'^2_{mx}}{\rho_m \sigma_{mx} \sigma_{fx} \sigma^2} x_2 + \frac{\sigma_{my}}{\rho_m \sigma_{mx}} \mu_{mx} - \mu_{my} - \frac{\rho_t \sigma_{fy} \sigma_{my} \sigma'^2_{mx}}{\rho_m \sigma_{mx} \sigma_{fy} \sigma^2} \mu_{fx} + \frac{\sigma_{my} \sigma'^2_{mx}}{\rho_m \sigma_{mx} \sigma^2} \mu_{fy} \right)^2}{\sigma''^2_{mx} \frac{(\sigma'^2_{mx} + \sigma^2)^2}{\sigma^4} \frac{\sigma^2_{my}}{\rho^2 \sigma^2_{mx}}} \\ &= \exp\left(-\frac{(y_1 - \mu_{my})^2}{2\sigma^2_{my}}\right) \exp\left(-\frac{(y_1 - *)^2}{2 \frac{\sigma^2_{my} \sigma'^2_{mx} (\sigma'^2_{mx} + \sigma^2)^2}{\sigma^4 \rho^2 \sigma^2_{mx} + \sigma'^2_{mx} (\sigma'^2_{mx} + \sigma^2)^2}}\right) \exp\left(-\frac{\left( \mu_{my} + \frac{\rho_t \sigma_{fy} \sigma_{my} \sigma'^2_{mx}}{\rho_m \sigma_{mx} \sigma_{fx} \sigma^2} x_2 + \frac{\sigma_{my}}{\rho_m \sigma_{mx}} \mu_{mx} - \mu_{my} - \frac{\rho_t \sigma_{fy} \sigma_{my} \sigma'^2_{mx}}{\rho_m \sigma_{mx} \sigma_{fy} \sigma^2} \mu_{fx} + \frac{\sigma_{my} \sigma'^2_{mx}}{\rho_m \sigma_{mx} \sigma^2} \mu_{fy} - \frac{\sigma'^2_{mx} + \sigma^2}{\sigma^2} \frac{\sigma_{my}}{\rho_m \sigma_{mx}} x_1 \right)^2}{2(\sigma^2_{my} + \sigma'^2_{mx} \frac{(\sigma'^2_{mx} + \sigma^2)^2}{\sigma^4} \frac{\sigma^2_{my}}{\rho^2 \sigma^2_{mx}})}\right) \\ &= \exp\left(-\frac{(y_1 - *)^2}{2 \frac{\sigma^2_{my} \sigma'^2_{mx} (\sigma'^2_{mx} + \sigma^2)^2}{\sigma^4 \rho^2 \sigma^2_{mx} + \sigma'^2_{mx} (\sigma'^2_{mx} + \sigma^2)^2}}\right) \exp\left(-\frac{\left( \frac{\rho_t \sigma_{fy} \sigma_{my} \sigma'^2_{mx}}{\rho_m \sigma_{mx} \sigma_{fx} \sigma^2} x_2 + \frac{\sigma_{my}}{\rho_m \sigma_{mx}} \mu_{mx} - \frac{\rho_t \sigma_{fy} \sigma_{my} \sigma'^2_{mx}}{\rho_m \sigma_{mx} \sigma_{fy} \sigma^2} \mu_{fx} + \frac{\sigma_{my} \sigma'^2_{mx}}{\rho_m \sigma_{mx} \sigma^2} \mu_{fy} - \frac{\sigma'^2_{mx} + \sigma^2}{\sigma^2} \frac{\sigma_{my}}{\rho_m \sigma_{mx}} x_1 \right)^2}{2 \frac{\rho^2 \sigma^2_{mx} (\sigma'^2_{mx} + \sigma^2)^2}{\sigma^4} \frac{\sigma^2_{my}}{\rho^2 \sigma^2_{mx}}}\right) \\ &= \exp\left(-\frac{(y_1 - *)^2}{2 \frac{\sigma^2_{my} \sigma'^2_{mx} (\sigma'^2_{mx} + \sigma^2)^2}{\sigma^4 \rho^2 \sigma^2_{mx} + \sigma'^2_{mx} (\sigma'^2_{mx} + \sigma^2)^2}}\right) \exp\left(-\frac{\left( \frac{\rho_t \sigma_{fy} \sigma'^2_{mx}}{\sigma_{tx} (\sigma'^2_{mx} + \sigma^2)} x_2 + \frac{\sigma^2}{\sigma'^2_{mx} + \sigma^2} \mu_{mx} - \frac{\rho_t \sigma_{fy} \sigma'^2_{mx}}{\sigma_{ty} (\sigma'^2_{mx} + \sigma^2)} \mu_{fx} + \frac{\sigma'^2_{mx}}{\sigma'^2_{mx} + \sigma^2} \mu_{fy} - x_1 \right)^2}{2 \frac{\sigma^4 \rho^2 \sigma^2_{mx} + \sigma'^2_{mx} (\sigma'^2_{mx} + \sigma^2)^2}{(\sigma'^2_{mx} + \sigma^2)^2}}\right) \end{aligned}$$

father, female gets preference from father.

This model was built to study whether song diversity can be maintained when two populations come back together. However, even within one population there are interesting dynamics.

1. There can be many discrete “popular” songs when  $\sigma_m^2$  is small,  $\sigma_f^2$  is high, and  $\sigma^2$  is intermediate.
2. When the equilibrium population is unimodal, there is an interesting interaction between  $\sigma^2$  and  $\sigma_f^2$ : At low  $\sigma^2$  and low  $\sigma_f^2$ , the males are even more narrowly distributed than the females because the male trait preferred by the most females does way better than anyone else. At low  $\sigma^2$  and high  $\sigma_f^2$ , the male distribution matches the female distribution. At high  $\sigma^2$ ,  $\sigma_f^2$  does not matter and the male distribution is moderately narrow because average males can mate with most females.

## References

- [1] R. Tucker Gilman and Genevieve M. Kozak. Learning to speciate: The biased learning of mate preferences promotes adaptive radiation. *Evolution*, 69(11):3004–3012, 2015.
- [2] D.E. Irwin. Culture in songbirds and its contribution toward the evolution of new species. In E. Slingerland and M. Collard, editors, *Creating Consilience: Integrating the Sciences and the Humanities*, pages 163–178. Oxford University Press, 2012.
- [3] Mark Kirkpatrick and Lee Alan Dugatkin. Sexual selection and the evolutionary effects of copying mate choice. *Behavioral Ecology and Sociobiology*, 34:443–449, 1994.
- [4] Mark Kirkpatrick, Trevor Price, and Stevan J. Arnold. The darwin-fisher theory of sexual selection in monogamous birds. *Evolution*, 44(1):180–193, 1990.
- [5] R. Lachlan and M. Servedio. Song learning accelerates allopatric speciation. *Evolution*, 58(9):2049–2063, 2004.
- [6] Russell Lande. Models of speciation by sexual selection on polygenic traits. *Proceedings of the National Academy of Sciences*, 78(6):3721–3725, 1981.
- [7] Louise S. Mead and Stevan J. Arnold. Quantitative genetic models of sexual selection. *TRENDS in Ecology and Evolution*, 19(5):264–271, 2004.