

Student Name: \_\_\_\_\_

Math 131

Homework 8

Due Date: November 9, 2018

## Homework 8

---

**Problem 1.**

---

**Solution.**

The terms *limit* and *limit point* mean different things, but the notions are easy to confuse. For example, the constant sequence  $1, 1, \dots, 1, \dots$  is convergent with *limit* 1. As a subset of the real line, however, its values are just equal to the set  $\{1\}$ , which cannot have a limit point.

1. Explain why the set  $\{1\}$  in  $\mathbb{R}$  has no limit points.
2. To clarify the notions of limit and limit point, prove the following statement: If a convergent sequence in a metric space has infinitely many distinct points, then its limit is a limit point of the set of points of the sequence.

---

**Problem 2.**

Give an example of a sequence  $\{x_n\}$  with values in  $[0, 1]$  that has the following property: For every  $x \in [0, 1]$ , we can find a subsequence  $\{x_{n_k}\}$  such that  $x_{n_k} \rightarrow x$  as  $k \rightarrow \infty$ .

---

**Solution.**

---

**Problem 3.**

This exercise reveals additional interesting qualities of the Cantor set.

Let  $C$  be the Cantor set as defined in Rudin 2.44. Further, note that the sum  $C + C = \{x + y : x, y \in C\}$ . The goal of this exercise is to prove that  $C + C = [0, 2]$ . Since  $C \subset [0, 1]$ , it follows immediately that  $C + C \subset [0, 2]$ . Thus, it remains only to show the inclusion in the other direction, namely that  $[0, 2] \subset \{x + y : x, y \in C\}$ . That is, given any  $s \in [0, 2]$ , we must show there exist numbers  $x, y \in C$  that satisfy  $x + y = s$ . Do so through the following steps:

1. Recall the notation from class in the construction of the Cantor set:  $C = \bigcap_{n=0}^{\infty} K_n$  where  $K_n$  is the union of  $2^n$  closed subintervals of  $[0, 1]$  in which successive open middle third subintervals have been removed. Namely,  $K_0 = [0, 1]$ ,  $K_1 = [0, 1/3] \cup [2/3, 1]$ ,  $K_2 = [0, 1/9] \cup [2/9, 3/9] \cup [6/9, 7/9] \cup [8/9, 1]$ , and so forth. Show that there exist  $x_1, y_1 \in K_1$  for which  $x_1 + y_1 = s$ .
2. Use induction to show in general that for an arbitrary  $n \in \mathbb{N}$ , we can always find  $x_n, y_n \in K_n$  for which  $x_n + y_n = s$ .
3. Show that we can find  $x$  and  $y$  in  $C$  satisfying  $x + y = s$ . (Note: The sequences  $\{x_n\}$  and  $\{y_n\}$  do not necessarily converge. Nonetheless, these can be used to produce the desired result.)

---

**Solution.**