

***In vitro* evaluation of the genotoxicity of CeO₂ nanoparticles in human peripheral blood lymphocytes using cytokinesis-block micronucleus test, comet assay, and gamma H2AX**

Toxicology and Industrial Health
2018, Vol. 34(5) 293–300
© The Author(s) 2018
Reprints and permissions:
sagepub.co.uk/journalsPermissions.nav
DOI: 10.1177/0748233717753780
journals.sagepub.com/home/tih


Serpil Könen-Adıgüzel and Serap Ergene

Abstract

Engineered nanoparticles (ENPs) are used in a wide range of applications because of their unique properties. Cerium dioxide nanoparticles (CeO₂ NPs) are one of the important ENPs, and they can cause negative health effects, such as genotoxicity, in humans and other living organisms. The aim of this work was to analyze the genotoxic effects of short-term (3–24 h) CeO₂ NPs exposure to cultured human blood lymphocytes. Three genotoxicity systems “cytokinesis-block micronucleus test, comet assay, and gamma H2AX test” were used to show the genotoxic potential of CeO₂ NPs (particle size <25 nm, concentrations: 6, 12, and 18 µg/mL). Hydrogen peroxide was selected as the positive-control genotoxic agent. Our results indicate that CeO₂ NPs have genotoxic potential on human peripheral blood lymphocytes cells even at 3–24 h exposure under *in vitro* conditions.

Keywords

CeO₂ NPs, comet assay, micronucleus test, gamma H2AX, blood lymphocytes, *in vitro*

Received 31 July 2017; Revised 20 November 2017; Accepted 16 December 2017

Introduction

The use of nanomaterials (size between 1 nm and 100 nm) in the world has become an important component of consumer products in recent years. This fact has gradually led to an increase in concerns and debates about the impact of nanomaterials on different biological systems (Courbiere et al., 2013; Dhawan and Sharma, 2010). Nanoparticles (NPs) can be classified into two groups such as natural (soil colloids, fullerenes, carbon nanotubes, biogenic magnetite, and airborne nanocrystals of sea salts) and anthropogenic NPs (carbon black, carbon nanotubes, platinum, and engineered/manufactured NPs) (Dhawan and Sharma, 2010). According to assessments in recent years, engineered NPs, which have small size, large specific surface area and surface reactivity, are extensively used in many industries such as electronic, textile, cosmetic, pharmacology, and medicine (Courbiere et al., 2013). However, their effects on health and potential biological impacts need to be determined for risk assessment.

Nanomaterials have frequently been used in nanotechnology products. Lanthanide cerium dioxide nanoparticles (CeO₂ NPs) have a wide application in a variety of fields, such as catalysts, solid fuel cells, ultraviolet absorbents, gas sensors, glass and ceramic applications (Srinivas et al., 2011). Therefore, the risk of both occupational and environmental exposure to CeO₂ NPs has risen for humans. According to an OECD report, CeO₂ NPs are one of the most important NPs for which human and environmental impacts need to be assessed (Srinivas et al., 2011). Although some researchers reported that CeO₂ NPs exhibit antioxidant properties (Colon et al., 2010; Rubio et al.,

Department of Biology, Faculty of Science and Letters, Mersin University, Mersin, Turkey

Corresponding author:

Serpil Könen-Adıgüzel, Department of Biology, Faculty of Science and Letters, Mersin University, Mersin 33343, Turkey.
Email: serpillkonen@gmail.com

2016; Schubert et al., 2006), some recent reports have shown that CeO₂ NPs' exposure can give rise to negative effects on human health such as genotoxicity, toxicity, and apoptosis (Aalapati et al., 2014; Mittal and Pandey, 2014; Marzi et al., 2013).

Some tests, including the micronucleus test, comet assay, and gamma H2AX, are commonly used to assess the genotoxicity of many different chemicals (drug, pesticide, NPs, and phthalates) *in vivo* as well as *in vitro* studies (Balasubramanyam et al., 2009; Jugan et al., 2012; Tsaousi et al., 2010).

The *in vitro* micronucleus test is a standard cytogenetic test for genotoxicity. It is simple, useful, and applicable in different cell types (Kirklanda et al., 2011; Lindberga et al., 2009; Meschini et al., 2015). The type performed using cytochalasin B is the most useful micronucleus assay which produces binucleated cells (BNCs) (Battal et al., 2015). The comet assay, also called single-cell gel electrophoresis (SCGE) assay, is extensively used as a marker for the assessing of DNA break in single cell (Tice et al., 2000). This technique can even be applied to a limited number of cells. The alkaline type of SCGE has also been used to evaluate single-stranded DNA breaks and DNA repair of individual cells (Collins, 2004). Additionally, phosphorylated histone H2AX, also known as γH2AX, is susceptible and fast in revealing DNA double-strand breaks in the cells. This test evaluates genotoxic damage, which is DNA damage caused by various agents in interphase cells (Cavaş et al., 2012).

There is limited knowledge related to genotoxicity of cerium oxide NPs (Li et al., 2016). In this study, we aimed to evaluate *in vitro* genotoxic potential of CeO₂ NPs in human blood lymphocytes exposed to different concentrations (concentrations: 6, 12, and 18 µg/mL; particle size <25 nm). During our studies, the micronucleus test, the SCGE assay, and gamma H2AX foci formation assays were used as genotoxicity endpoints.

Materials and methods

Chemicals

Pure cerium dioxide NPs, phosphate buffered saline (PBS), Ethylenediaminetetraacetic acid (EDTA), Triton X-100, normal melting point agarose, and low melting point agarose were purchased from Sigma-Aldrich (St. Louis, USA). Anti-gamma H2AX (phospho S139) antibody (9f3) primary and Alexa Fluor[®] 488 Goat anti-Mouse IgG secondary antibodies were supplied by Abcam (Cambridge, MA, USA) and




Figure 1. SEM images of CeO₂ NPs.

BioLegend (San Diego, CA, USA). All other chemicals used in the experiments were also purchased from Bio Ind (Cromwell, CT, USA). Human blood lymphocytes were treated with 100 µM hydrogen peroxide (H₂O₂) as positive control. CeO₂ NPs dissolved in sterile distilled water and were used at concentration of 0.5% (v/v).

Characterization of NPs

CeO₂ NPs in stock solution were dispersed by ultrasonication for 20 min, before each treatment dose was prepared. Once the particle stock solution was prepared in the cell culture medium, the addition of NPs to the cell culture medium caused discoloration at high concentration. Therefore, distilled water was used as solvent of NPs. Moreover, others researchers used distilled water for NP stock solution (Estevez et al., 2011; Molina et al., 2014; Selvaraj et al., 2015). The morphology and the size of powdered NP were determined using scanning electron microscopy (SEM) (Figure 1) and X-ray diffraction analysis. The average particle size was confirmed by BET (≤25 nm).

Blood sampling

Human blood lymphocytes were attained from healthy nonsmoking donors (four men and four women, mean age 24 years). The donors were interviewed about the possible influence of this study. Each person read the questionnaire and signed it. The study protocol was designed according to ethical standards and was approved by the Ethical Committee of the Medical Faculty at Mersin University (document

number: 2012/81). The genotoxicity tests were evaluated using three culture tubes for each individual. Each test was applied three times in order to avoid inaccuracy (three repeated, total individual: 8).

Cell culture

Blood sample (3 mL) and Histopaque-1077 (3 mL) was centrifuged at 2000 rpm for 30 min. The visible opaque layer was used as lymphocyte suspension after centrifugation. Subsequently, the lymphocyte suspension (0.5 mL) was added to RPMI1640 cell culture medium (4.5 mL) supplemented with 20% fetal calf serum, 2 mM L-glutamine, 10 mg/mL phytohemagglutinin, 100 U/mL penicillin, and 100 mg/mL streptomycin. The cell cultures were incubated at 37°C with 5% CO₂ for 24 h.

Using doses

CeO₂ NPs doses were determined and selected due to the available data of NPs dose ranges humans are exposed to both occupationally and environmentally (Gehlhaus et al., 2009; Hussain et al., 2012). Three doses were used, a low dose (6 µg/mL), a middle dose (12 µg/mL), and a high dose (18 µg/mL).

Cytokinesis-block micronucleus test

The cytokinesis-block micronucleus test was utilized to evaluate chromosomal damages. Lymphocytes, which were transferred in sterile cell culture dishes (15 mL), were exposed to different concentrations of CeO₂ NPs and H₂O₂ (100 µM) for 24 h. Following this exposure, cells were resuspended in 0.075 M KCl, and incubated for 3 min. The cells were fixed three times in Carnoy fixative (methanol: glacial acetic acid, 3:1) and then dropped onto clean slides. Finally, the number of micronuclei in BNCs was evaluated under light microscope in 2000 exposed to cells after the nucleus was stained by 5 % giemsa for 10 min. (Figure 2).

Alkaline comet assay

The comet assay was performed to detect the DNA strand breaks induced by CeO₂ NPs. After exposure to different concentrations of CeO₂ NPs for 3 h, 100 µL of lymphocyte, the suspension was mixed with 2% low melting temperature agarose and embedded on slides precoated with 5% normal melting point agarose. Subsequently, the slides were placed in a bottle containing the lysis solution (pH 10), including 2.5 M NaCl, 0.1 M EDTA, 10 mM Tris Base, and 1% Triton




Figure 2. Binucleated cell including micronucleus in lymphocyte cultures exposed to CeO₂ NPs.




Figure 3. Comet assay views in lymphocyte cultures exposed to CeO₂ NPs.

X for 1 h at 4°C. The slides were denatured in alkaline buffer (0.3 M NaCl, 1 mM EDTA) for 30 min and performed at 25 V and 300 mA for 20 min, and then washed with neutralization buffer (0.5 M Tris-HCl, pH 7.5) for 10 min. After neutralization, the slides were dehydrated by 70% ethanol, air dried, stained with ethidium bromide, and analyzed with a fluorescence microscope (Figure 3). All processes were performed in dark to prevent DNA damage.

Immunofluorescence for γ-H2AX foci formation

γH2AX foci were used to show that CeO₂ NPs lead to double-strand DNA breaks. All experiments were designed according to Scarpato et al. (2013). Briefly, cells that were exposed to CeO₂ NPs were centrifuged and resuspended in 0.075 M KCl, and incubated for 3 min. Then the cells were centrifuged again and




Figure 4. XRD patterns of CeO₂ NPs.

gradually fixed in methanol: glacial acetic acid (5:1), methanol and methanol: glacial acetic acid (3:1). Following this fixation, the cell solution was transplanted onto pre-cleaned slides. After the slides were washed twice in 1X PBS, the cells were blocked in 1X PBS/0.3% Triton X-100 (PBS/Triton; Sigma-Aldrich) and washed again three times in 1X PBS. The slides were steeped overnight at 4°C with primary antibody diluted 1:100 PBS/Triton. The following day, the cells were washed three times in 1X PBS and then incubated at room temperature for 2 h with secondary antibody diluted 1:100 in PBS/Triton. After three washes in 1X PBS, the slides were counterstained with 3–4 drops of 4',6-Diamidino-2-phenylindole. Then the slides were investigated using a fluorescence microscope.

Statistical analysis

The effect of the NP treatment on genetic material (nuclei) was analyzed using the SPSS 11.5 program.

Results

Characterization of NPs

We evaluated the size of individual NPs by SEM, which was approximately 20.7 ± 2.7 nm. X-ray diffraction analysis confirmed the crystalline size of CeO₂ NPs from diffraction peak (Figure 4). According to XRD analysis, the crystalline phase was determined as cubic. The BET result of CeO₂ NPs was determined by Sigma-Aldrich and it was indicated as ≤ 25 nm at the product information sheet. So the BET of NPs was not evaluated.

Table 1. Effects of CeO₂ NPs on the frequencies of MNBN% and CBPI values in human lymphocyte cells.

Dose	Micronucleus (%), mean \pm standard deviation	CBPI (%), mean \pm standard deviation
Negative control	1.00 \pm 0.76	2.08 \pm 0.05
Positive control	11.25 \pm 1.03 ^a	0.89 \pm 0.01 ^a
6 μ g/mL CeO ₂ NPs	10.00 \pm 1.31 ^a	1.19 \pm 0.01 ^a
12 μ g/mL CeO ₂ NPs	15.00 \pm 1.30 ^a	1.16 \pm 0.01 ^a
18 μ g/mL CeO ₂ NPs	18.87 \pm 1.12 ^a	1.13 \pm 0.02 ^a

MNBN: micronucleated binucleated cells; CBPI: cytokinesis-block proliferation index; CeO₂ NPs: cerium dioxide nanoparticles.

^aStatistical difference from negative control $p < 0.0001$.

Cytokines-blocked micronucleus test

The frequency of micronuclei was determined in human lymphocytes treated with various doses of CeO₂ NPs. The frequency of micronuclei was 1% in the negative control group. The micronucleated cell frequency enhanced to 11.25% in the H₂O₂ (100 μ M) positive control group. The lowest concentration of CeO₂ NPs (6 μ g/mL) caused micronucleus frequency (10%) as well as positive control (11.25%). The frequency of micronuclei increased to 15% and 18.87% when CeO₂ NP doses increased to 12 and 18 μ g/mL, respectively (Table 1).

Comet assay

Table 2 shows the genetic damage index (GDI) values in human peripheral lymphocytes treated with CeO₂ NPs. Two doses used of CeO₂ NPs (12 and 18 μ g/mL) significantly increased the control GDI value of 0.5 to 1.01 and 1.19, respectively ($p < 0.05$). The lowest CeO₂ NP dose did not significantly increase the GDI value in human peripheral lymphocytes. However, treatment with positive control (H₂O₂) significantly increased the GDI value in human lymphocytes ($p < 0.001$), while the solvent control group did not significantly increase the GDI value. Our results showed that the GDI value induced by positive control was similar to that of the highest concentration of CeO₂ NPs (Table 2).

Gamma H2AX foci formation

Treatment with CeO₂ NPs was evaluated with gamma H2AX foci formation that shows DNA double-strand breaks in human lymphocytes. Any significant

Table 2. Comet assay results in human lymphocyte cells exposed to CeO₂ NPs.

Exposed dose	Exposed duration	Proportion of damaged nuclei (%)						Percentage of damaged cells	GDI
		T0	T1	T2	T3	T4	T5		
NC	3 h	71.62 ± 2.28	15.25 ± 1.27	3.00 ± 0.53	2.12 ± 0.54	8.00 ± 1.52	13.12 ± 1.93	0.59 ± 0.07	
PC	3 h	25.87 ± 2.15	20.62 ± 2.07	10.62 ± 1.19	12.25 ± 1.23	30.37 ± 0.80	53.25 ± 1.76 ^b	2.00 ± 0.04 ^b	
6 µg/mL CeO ₂ NPs	3 h	60.25 ± 3.28	20.00 ± 2.25	6.50 ± 1.25	2.62 ± 0.41	10.62 ± 1.79	19.75 ± 2.78	0.83 ± 0.08	
12 µg/mL CeO ₂ NPs	3 h	51.12 ± 3.66	23.00 ± 2.85	10.75 ± 1.87	3.37 ± 0.62	11.75 ± 1.90	25.87 ± 2.00	1.01 ± 0.07 ^a	
18 µg/mL CeO ₂ NPs	3 h	45.87 ± 2.80	23.50 ± 2.55	10.62 ± 1.53	4.87 ± 0.98	15.12 ± 2.86	30.62 ± 3.40 ^a	1.19 ± 0.11 ^a	

NC: negative control; PC: positive control (hydrogen peroxide); CeO₂ NPs: cerium dioxide nanoparticles; GDI: genetic damage index.

^aStatistical difference from negative control $p < 0.05$.

^bStatistical difference from negative control $p < 0.001$.

Table 3. Gamma H2AX results in human lymphocyte cells exposed to CeO₂ NPs.

Group	Concentration	Gamma H2AX ± SE
Negative control	—	2.12 ± 0.35
Positive control	100 µM	20.37 ± 0.80 ^a
Cerium dioxide	6 µg/mL	7.75 ± 0.36 ^a
Cerium dioxide	12 µg/mL	10.25 ± 0.52 ^a
Cerium dioxide	18 µg/mL	33.50 ± 0.82 ^a

CeO₂ NPs: cerium dioxide nanoparticles; SE: Standard Error.

^aStatistical difference from negative control $p < 0.0001$.

increase in gamma H2AX foci formation was not detected for the solvent control group, while the positive control H₂O₂ also significantly induced double-strand DNA breaks ($p < 0.0001$; Table 3). Furthermore, the negative control group did not induce the gamma H2AX formation. Treatment with three doses of CeO₂ NPs significantly increased DNA double-strand breaks in lymphocytes. Moreover, the highest dose of CeO₂ NPs induced a higher DNA damage than in positive control.

Discussion

Because of their unique properties, NPs may be useful in a diverse range of applications, such as the biomedical field, the use of nano vaccines and nano drugs. (Saptarshi et al., 2013). Hence, NPs' increased use in nanotechnology enhances the risk of its exposure to humans and other living organisms. These risks may cause cytotoxic, genotoxic, and biochemical adverse effects on the biological systems. Therefore, there is an urgent need to demonstrate these adverse effects of NPs on biological systems (Kumari et al., 2014a).

Because of the increasing evidence of NPs adverse effect on biological systems, scientists became concerned. As a consequence, studies on the possible adverse effects of NPs, especially genotoxicity, have increased (Gonzalez et al., 2008). Genotoxic effects of different types of NPs have been evaluated by several researchers (Asha Rani et al., 2009; Ghosh et al., 2010; Kumari et al., 2009; Trouiller et al., 2009; Wise et al., 2010). Shukla et al. (2011) investigated titanium dioxide NPs (TiO₂ NPs) in human epidermal cells which cause reactive oxygen species (ROS)-mediated genotoxicity. Their data demonstrated that TiO₂ NPs have a mild cytotoxic potential. Moreover, both *in vitro* and *in vivo* genotoxicity research of oxide NPs has increased recently. Singh et al. (2016) made a

research on genotoxic effects of chromium oxide NPs and microparticles in *Wistar* rats after 28 days of repeated oral exposure. Their finding provided initial data on the probable genotoxicity.

The genotoxic effects of CeO₂ NPs on different organism have been reported by some researchers (Ali et al., 2015; Gaiser et al., 2012; Kawai et al., 2015). However, these reports are conflicting. For example, several researchers showed (Aalapathi et al., 2014; Marzi et al., 2013; Mittal and Pandey, 2014) that CeO₂ has genotoxic effects, whereas some others have claimed (Colon et al., 2010; Rubio et al., 2016; Schubert et al., 2006) that CeO₂ has antigenotoxic effects. Auffan et al. (2009) measured the potential *in vitro* cyto- and genotoxicity of nano-sized CeO₂ (7 nm) on human dermal fibroblasts. Nano-sized CeO₂ at these dosages caused strong DNA lesions and these genotoxic effects occurred at very low doses. Kumari et al. (2014b) evaluated genotoxicity of cerium oxide NPs and cerium oxide micro in *Wistar* rats after 28 days of repeated oral administration. They used concentrations of 30, 300, and 600 mg/kg body weight (bw)/day of cerium oxide NPs. Their results showed that DNA damage has significantly increased in peripheral blood leukocytes and liver. Also, MN and CA have increased in peripheral blood and bone marrow. Benameur et al. (2015) evaluated the genotoxic effects of cerium oxide NPs' environmentally relevant doses on primary human dermal fibroblasts. They used a cytokinesis-blocked micronucleus assay for the genotoxicity study. The results showed that nano-CeO₂ increased the frequency of micronuclei in dermal fibroblasts. Moreover, the frequency of centromere negative micronuclei increased in cells, which were exposed to nano-CeO₂ for 48 h. In addition, Kaki et al. (2015) studied the effects of cerium NPs (CeNP) and CeO₂ on oxidative toxic stress in human lymphocytes. Although cerium NPs have an interesting potential in drug delivery, gene therapy, and molecular medicine, they found that CeO₂ is able to induce oxidative toxicity in human lymphocytes (24, 48, and 72 h exposure; 15, 30, 60, 120 µmol/mL; 30 nm). Similar results were also observed in different studies (Aalapathi et al., 2014; Mittal and Pandey, 2014; Marzi et al., 2013). Different from these studies, Pierscioneck et al. (2009) claimed that nanoceria did not cause any DNA damage and significant increases in the number of sister chromatid exchanges. Additionally, Pedram et al. (2015) used different concentrations of CeO₂ (0, 0.5, 1, and 2 ng/mL) in

human lymphocytes. Their results showed that CeO₂NPs have protective effects in very low concentrations against CP.

The difference of this study is that it used different doses and genotoxicity tests in human lymphocytes. The earlier studies showed that CeO₂ NPs have a cytotoxic effect, oxidative damage, and decrease of cell viability in human blood lymphocytes. In our study, a gamma H2AX foci formation test has been used for the first time to investigate the genotoxic effects of CeO₂ NPs in human lymphocytes *in vitro*. We used three doses of CeO₂ NPs (6, 12, and 18 µg/mL) in our study. Being exposed to NPs for 72 h significantly decreased the viability of human lymphocytes. Therefore, the lymphocytes for comet and gamma H2AX foci formation tests have been exposed to NPs for 3 h. However, the lymphocytes for micronucleus test have been exposed to NPs for 24 h. Exposure to three doses of NPs significantly increased the potential genotoxic effect. All the test systems showed that CeO₂ NPs have a potential genotoxic effect.

Although several researchers have demonstrated the protective and antioxidant properties of CeO₂ NPs in their work, we reported a potential genotoxic effect of CeO₂ NPs. Prior to our study, Auffan et al. (2009) investigated the genotoxic effects of these NPs on human dermal fibroblasts *in vitro*. Moreover, Srinivas et al. (2011) evaluated a 24-h, 48-h, and 14-day postexposure period on cytotoxicity, oxidative stress, and inflammation in rats with CeO₂ NPs. They found a significant reduction in cell viability and their results suggested that acute exposure to CeO₂ NPs through an inhalation route may cause cytotoxicity via oxidative stress.

Consequently, our results showed that the genotoxic effect of CeO₂ NPs is strictly dose dependent, and human lymphocytes are the sensitive cells. Additionally, the cells are useful tools to show DNA damage produced by nanomaterial samples. The genotoxic effect of CeO₂ NPs is thought to be due to the presence of ROS and oxidative DNA damage (Benameur et al., 2015; Kaki et al., 2015; Kawai et al., 2015; Mittal and Pandey, 2014). However, the genotoxic effect of CeO₂ NPs depends on cell types and from time to time these NPs have antigenotoxic effects when another genotoxic compound is also present (Marzi et al., 2013). Therefore, more studies are necessary to evaluate the genotoxic effect of CeO₂ NPs in specific cells.

Acknowledgment

The authors want to thank the Mersin University Scientific Research Projects Coordination Unit for providing devices used in this study (project no: BAP-FBE B (SKA) 2011-7 DR).

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The authors are indebted to Mersin University for its financial support of the study. This academic work was linguistically supported by the Mersin Technology Transfer Office Academic Writing Center of Mersin University.

References

- Aalapathi S, Ganapathy S, Manapuram S, et al. (2014) Toxicity and bio-accumulation of inhaled cerium oxide nanoparticles in CD1 mice. *Nanotoxicology* 8(7): 786–798.
- Ali D, Alarifi S, Alkahtani S, et al. (2015) Cerium oxide nanoparticles induce oxidative stress and genotoxicity in human skin melanoma cells. *Cell Biochemistry and Biophysics* 71: 1643–1651.
- Asha Rani PV, Kah Mun GL, Hande MP, et al. (2009) Cytotoxicity and genotoxicity of silver nanoparticles in human cells. *American Chemical Society NANO* 3(2): 279–290.
- Auffan M, Rose J, Orsiere T, et al. (2009) CeO₂ nanoparticles induce DNA damage towards human dermal fibroblasts in vitro. *Nanotoxicology* 3(2): 161–171.
- Balasubramanyam A, Sailaja N, Mahboob M, et al. (2009) In vivo genotoxicity assessment of aluminium oxide nanomaterials in rat peripheral blood cells using the comet assay and micronucleus test. *Mutagenesis* 24(3): 245–251.
- Battal D, Çelik A, Güler G, et al. (2015) SiO₂ nanoparticle-induced size-dependent genotoxicity—an in vitro study using sister chromatid exchange, micronucleus and comet assay. *Drug and Chemical Toxicology* 38(2): 196–204.
- Benameur L, Auffan M, Cassien M, et al. (2015) DNA damage and oxidative stress induced by CeO₂ nanoparticles in human dermal fibroblasts: evidence of a clastogenic effect as a mechanism of genotoxicity. *Nanotoxicology* 9(6): 696–705.
- Cavaş T, Cinkılıç N, Vatan Ö, et al. (2012) In vitro genotoxicity evaluation of acetamiprid in CaCo-2 cells using the micronucleus, comet and cH2AX foci assays. *Pesticide Biochemistry and Physiology* 104: 212–217.
- Collins AR (2004) The comet assay for DNA damage and repair: principles, applications, and limitations. *Molecular Biotechnology* 26: 249–261.
- Colon J, Hsieh N, Ferguson A, et al. (2010) Cerium oxide nanoparticles protect gastrointestinal epithelium from radiation-induced damage by reduction of reactive oxygen species and upregulation of superoxide dismutase 2. *Nanomedicine: Nanotechnology, Biology, and Medicine* 6: 698–705.
- Courbiere B, Auffan M, Rollais R, et al. (2013) Ultrastructural interactions and genotoxicity assay of cerium dioxide nanoparticles on mouse oocytes. *International Journal of Molecular Sciences* 14: 21613–21628.
- Dhawan A and Sharma V (2010) Toxicity assessment of nanomaterials: methods and challenges. *Analytical and Bioanalytical Chemistry* 398: 589–605.
- Estevez AY, Pritchard S, Harper K, et al. (2011) Neuroprotective mechanisms of cerium oxide nanoparticles in a mouse hippocampal brain slice model of ischemia. *Free Radical Biology & Medicine* 51: 1155–1163.
- Gaiser BK, Fernandes TF, Jepson MA, et al. (2012) Interspecies comparisons on the uptake and toxicity of silver and cerium dioxide nanoparticles. *Environmental Toxicology and Chemistry* 31: 144–154.
- Gehlhaus M, Osier M, Lladós F, et al. (2009) Toxicological Review of cerium oxide and cerium compounds (CAS No. 1306-38-3) In support of summary information on the integrated risk information system (IRIS). Washington, DC: US Environmental Protection Agency. EPA/635/R-08/002F.
- Ghosh M, Bandyopadhyay M, and Mukherjee A (2010) Genotoxicity of titanium dioxide (TiO₂) nanoparticles at two trophic levels: plant and human lymphocytes. *Chemosphere* 81: 1253–1262.
- Gonzalez L, Lison D, and Kirsch-Volders M (2008) Genotoxicity of engineered nanomaterials: a critical review. *Nanotoxicology* 2(4): 252–273.
- Hussain S, Al-Nsour F, Rice AB, et al. (2012) Cerium dioxide nanoparticles induce apoptosis and autophagy in human peripheral blood monocytes. *ACS NANO* 6(7): 5820–5829.
- Jugan ML, Barillet S, Simon-Deckers A, et al. (2012) Titanium dioxide nanoparticles exhibit genotoxicity and impair DNA repair activity in A549 cells. *Nanotoxicology* 6(5): 501–513.
- Kaki A, Younesi H, Hosseini SA, et al. (2015) Comparison the effects of cerium nanoparticles (CeNP) and cerium oxide (CeO₂) on oxidative toxic stress in human lymphocytes. *Research in Molecular Medicine* 3(3): 28–32.

- Kawai K, Li YS, Kawasaki Y, et al. (2015) Oxidative DNA damage in rat lung after exposure to cerium oxide nanoparticles. *European Respiratory Journal* 46: 59.
- Kirklanda D, Reeve L, Gatehousec D, et al. (2011) A core in vitro genotoxicity battery comprising the Ames test plus the in vitro micronucleus test is sufficient to detect rodent carcinogens and in vivo genotoxins. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 721: 27–73.
- Kumari M, Mukherjee A, and Chandrasekaran N (2009) Genotoxicity of silver nanoparticles in *Allium cepa*. *Science of the Total Environment* 407: 5243–5246.
- Kumari M, Singh SP, Chinde S, et al. (2014a) Toxicity study of cerium oxide nanoparticles in human neuroblastoma cells. *International Journal of Toxicology* 33: 1–12.
- Kumari M, Kumari SI, and Grover P (2014b) Genotoxicity analysis of cerium oxide micro and nanoparticles in Wistar rats after 28 days of repeated oral administration. *Mutagenesis* 2(6): 467–479.
- Li Y, Li P, Yu H, et al. (2016) Recent advances (2010–2015) in studies of cerium oxide nanoparticles' health effects. *Environmental Toxicology and Pharmacology* 44: 25–29.
- Lindberga HK, Falcka GCM, Suhonena S, et al. (2009) Genotoxicity of nanomaterials: DNA damage and micronuclei induced by carbon nanotubes and graphite nanofibres in human bronchial epithelial cells in vitro. *Toxicology Letters* 186: 166–173.
- Marzi LD, Monaco A, Lapuente JD, et al. (2013) Cytotoxicity and genotoxicity of ceria nanoparticles on different cell lines in vitro. *International Journal of Molecular Sciences* 14(2): 3065–3077.
- Meschini R, Berni A, Filippi S, et al. (2015) The micronucleus assay in mammalian cells in vitro to assess health benefits of various phytochemicals. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 793: 79–85.
- Mittal S and Pandey AK (2014) Cerium oxide nanoparticles induced toxicity in human lung cells: role of ROS mediated DNA damage and apoptosis. *Hindawi Publishing Corporation BioMed Research International* 2014: 14.
- Molina RM, Konduru NV, Jimenez RJ, et al. (2014) Bioavailability, distribution and clearance of tracheally instilled, gavaged or injected cerium dioxide nanoparticles and ionic cerium. *Environmental Science Nano* 1: 561–573.
- Pedram S, Mohammadirad A, Rezvanfar MA, et al. (2015) On the protection by the combination of CeO₂ nanoparticles and sodium selenite on human lymphocytes against chlorpyrifos-induced apoptosis *in vitro*. *Cell Journal* 17(2): 361–371.
- Pierscionek BK, Li Y, Yasseen AA, et al. (2009) Nanoceria have no genotoxic effect on human lens epithelial cells. *Nanotechnology* 21: 3.
- Rubio L, Annangi B, Vila L, et al. (2016) Antioxidant and anti-genotoxic properties of cerium oxide nanoparticles in a pulmonary like cell system. *Archives Toxicology* 90: 269–278.
- Saptarshi SR, Duschl A, and Lopata AL (2013) Interaction of nanoparticles with proteins: relation to bio-reactivity of the nanoparticle. *Journal of Nanobiotechnology* 11: 26.
- Scarpato R, Castagna S, Aliotta R, et al. (2013) Kinetics of nuclear phosphorylation (γ-H2AX) in human lymphocytes treated in vitro with UVB, bleomycin and mitomycin C. *Mutagenesis* 28(4): 465–473.
- Schubert D, Dargusch R, Raitano J, et al. (2006) Cerium and yttrium oxide nanoparticles are neuroprotective. *Biochemical and Biophysical Research Communications* 342: 86–91.
- Selvaraj V, Nepal N, Rogers S, et al. (2015) Inhibition of MAP kinase/NF-κB mediated signaling and attenuation of lipopolysaccharide induced severe sepsis by cerium oxide nanoparticles. *Biomaterials* 59: 160–171.
- Shukla RK, Sharma V, Pandey AK, et al. (2011) ROS-mediated genotoxicity induced by titanium dioxide nanoparticles in human epidermal cells. *Toxicology In Vitro* 25: 231–241.
- Singh SP, Chinde S, Kamal SSK, et al. (2016) Genotoxic effects of chromium oxide nanoparticles and microparticles in Wistar rats after 28 days of repeated oral exposure. *Environmental Science and Pollution Research* 23: 3914–3924.
- Srinivas A, Jaganmohan Rao P, Selvam G, et al. (2011) Acute inhalation toxicity of cerium oxide nanoparticles in rats. *Toxicology Letters* 205: 105–115.
- Tice RR, Agurell E, Anderson D, et al. (2000) Single cell gel/comet assay: guidelines for in vitro and in vivo genetic toxicology testing. *Environmental and Molecular Mutagenesis* 35: 206–221.
- Trouiller B, Reliene R, Westbrook A, et al. (2009) Titanium dioxide nanoparticles induce DNA damage and genetic instability in vivo in mice. *Cancer Research* 69(22): 8784–8789.
- Tsaousi A, Jones E and Case CP (2010) The in vitro genotoxicity of orthopaedic ceramic (Al₂O₃) and metal (CoCr alloy) particles. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 697: 1–9.
- Wise JPSr, Goodale BC, Wise SS, et al. (2010) Silver nanospheres are cytotoxic and genotoxic to fish cells. *Aquatic Toxicology* 97: 34–41.