

Particulate matter in the environment: pulmonary and cardiovascular effects

Ernesto Alfaro-Moreno^{a,b}, Tim S. Nawrot^a, Abderrahim Nemmar^{a,c} and Benoit Nemery^a

Purpose of review

The mechanisms related to adverse respiratory and cardiovascular effects in populations exposed to particulate matter are under debate and different models have been used to further our understanding of the various aspects of those effects. In this review we present some studies that may give new insights into the cellular and systemic mechanisms related to particulate matter toxicity.

Recent findings

Strong epidemiological evidence is now available regarding exposure markers and health effects. This is evident in the correlation between carbon content in macrophages and decrease in lung function, an increase in the risk of chronic obstructive pulmonary disease, lung cancer and postnatal mortality. The role of outdoor temperature and a missing allele for *GSTM1* and the impact of these factors on cardiovascular effects are also reported. At the experimental level, the effects of particulate matter and the interactions between different cell types, the role of toll-like receptor-2 and 4, the translocation of particles through cell monolayers and the activation of endothelial cells by particulate matter are also discussed. The role of composition is under intense debate, and different statistical analyses have been proposed.

Summary

Experimental studies on the effects of particulate matter are giving plausibility to the epidemiological findings, but the possible mechanisms of action are also becoming a hot topic.

Keywords

cardiovascular effects, PM₁₀, PM_{2.5}, pulmonary effects, ultrafine particles

Abbreviations

COPD chronic obstructive pulmonary disease
DEP diesel exhaust particle
TLR toll-like receptor
UFP ultrafine particle

© 2007 Lippincott Williams & Wilkins
1070-5287

Introduction

Airborne particulate matter has been associated with increases in morbidity and mortality. The historical events of the Meuse valley fog of 1930 [1] and the London smog of 1952 [2] triggered successful efforts and legislations to reduce air pollution. Nevertheless, a relation between urban air pollution and morbidity and mortality persists even at present levels of urban pollution [3]. Understanding the mechanisms of the adverse health effects caused by particulate matter represents an important scientific challenge. One of these challenges is to explain which properties of particulate matter are critical in causing injury. Particles capable of penetrating into the lower respiratory tract, and even into the alveolar region, are those with a mean aerodynamic size of up to 10 µm (PM₁₀), up to 2.5 µm (PM_{2.5}) and up to 0.1 µm (ultrafine particles, UFPs) [4]. Particulate matter can trigger multiple cellular responses in the lung such as cytotoxicity, inflammation, and mutagenesis. Moreover, the adverse effects of particulate matter are not confined to the lung, but there are also extrapulmonary manifestations of exposure to particulate matter. These systemic effects may be due to a 'spill-over effect' of pulmonary inflammation or to translocation of a fraction of particles into the circulation.

We review the recent advances in the epidemiologic and mechanistic research on particulate matter pollution, with some bias towards our own contributions to the field. An excellent and extensive general review on the topic of air pollution has been recently published by Pope and Dockery [5**].

Respiratory effects of particulate matter


Exposure to particulate matter may have different outcomes, including lung function deterioration, chronic obstructive lung disease, allergic illness and lung cancer (Fig. 1).

Curr Opin Pulm Med 13:98–106. © 2007 Lippincott Williams & Wilkins.

^aLung Toxicology Unit, Laboratory of Pneumology, KU Leuven, Belgium, ^bBasic Research, Nacional Cancer Institute, Mexico and ^cDepartment of Physiology, College of Medicine & Health Sciences, Sultan Qaboos University, Sultanate of Oman

Correspondence to Benoit Nemery, Research Unit of Lung Toxicology, Laboratory of Pneumology, KU Leuven, Herestraat 49 (O&N 706), B-3000 Leuven, Belgium
Tel: +32 16 347121; fax: +32 16 347124; e-mail: ben.nemery@med.kuleuven.be

Current Opinion in Pulmonary Medicine 2007, 13:98–106

Figure 1 Diagram of the possible targets and effects of particulate matter (PM) on the airways

Cell death (necrosis, apoptosis), proinflammatory factors secretion, DNA damage, tissue damage and remodelling. COPD, chronic obstructive pulmonary disease; IL, interleukin; TLR, toll-like receptor; TNF, tumor necrosis factor; UFP, ultrafine particle.

Pre and postnatal effects of air pollution

Several studies have shown adverse effects of particulate matter on pregnancy outcomes, including preterm delivery, lower birth weight and an increase in postneonatal infant mortality. A PM₁₀ increase of 10 $\mu\text{g}/\text{m}^3$ is associated with a 5% increase in postneonatal mortality for all causes and around 22% for postneonatal mortality from respiratory diseases [6 \bullet]. The critical periods and the mechanisms remain unclear.

Lung function

Exposure to air pollution during childhood may impair the development of the lung and thus lead to a lower level of peak pulmonary function in early adulthood. Kulkarni *et al.* [7 $\bullet\bullet$] were able to show, in children, that the load of carbon particles found in airway macrophages, recovered from induced sputum, correlated with their

exposure to particulate matter and that this biologic index of exposure was inversely correlated with lung function. Thus, each 1.0 $\mu\text{g}/\text{m}^3$ increase in primary PM₁₀ was associated with an increase of 0.10 μm^2 in the carbon content of airway macrophages, and each increase of 1.0 μm^2 in carbon content was associated with a reduction of 17% in forced expiratory volume in 1 s, of 12.9% in forced vital capacity, and of 34.7% in the forced expiratory flow between 25 and 75% of the forced vital capacity. These results strengthen the observations from Gauderman *et al.* [8] who reported that increased exposure to PM₁₀ was associated with impaired growth of lung function.

Allergic illness

Animal studies, as well as controlled human exposure studies, have shown that diesel exhaust particles (DEPs)

can increase the risk of allergy development and exacerbate allergic reactions [9,10]. The evidence from population studies to support this observation is still conflicting, possibly as a result of differences in the mix of transport-related air pollutants [11].

Chronic obstructive lung disease

Acute exacerbations of chronic obstructive pulmonary disease (COPD) have been associated with short-term variations in air pollution [12]. Schikowski *et al.* [13^{••}] investigated the influence of long-term exposure to air pollution on respiratory symptoms and pulmonary function in middle-aged women. A $7 \mu\text{g}/\text{m}^3$ increase in 5-year means of PM_{10} (interquartile range) was associated with a 33% increase in the risk of COPD. These findings were confirmed when replacing the estimates of PM_{10} by distance of home to a major road.

Lung cancer

Until now, six prospective studies have indicated that long-term exposure to urban pollution is likely to increase the risk of lung cancer [14,15[•],16–19]. Compared with the risk associated with active smoking, the risk associated with air pollution is low (+10–20% for each $10 \mu\text{g}/\text{m}^3$ increase in PM_{10} or $\text{PM}_{2.5}$). A recent European cohort study (European Prospective Investigation into Cancer and Nutrition) yielded a positive but nonsignificant association between lung cancer incidence and residence near heavy traffic roads (1.46, CI 0.89–2.40) [14]. In the same cohort, however, a significant interaction was found between the exposure variable ‘distance from heavy traffic road’ and polymorphisms in genes coding for base excision repair (*XRCC1-Arg399Gln*) and double-strand break repair (*BRCA2-Asn372His*) [20^{••}]. An ecological study suggests that some of the differences in lung cancer mortality within Europe could be explained by exposure to particulate matter pollution [21].

Experimental studies

Various outcomes such as cell necrosis and apoptosis [22–26], secretion of cytokines (tumor necrosis factor α (TNF α), interleukin (IL)-1 β , IL-6, IL-8) [22,25–27] and DNA damage (mutation assays, single and double-strand breaks, DNA degradation, and comet assay) [25,28,29] have been described upon incubation of various types of cells with particulate matter. Two main questions remain: what are the mechanisms involved and which components of the particulate matter are responsible for the effects?

Rats instilled with PM_{10} present a mild oedematous response 3 days after exposure [30[•]]. This response was linked to an increase in the expression of genes for the ion channel-acetylcholine receptor, IL-2 and serine/threonine kinase. The lack of other markers was explained by an earlier response that may not last long.

In-vitro evaluation of particulate matter on epithelial cells, macrophages and lymphocytes provided new evidence in this direction: proinflammatory mediators such as TNF α , IL-1 β , IL-8 are secreted in parallel to anti-inflammatory mediators such as IL-6 and transforming growth factor β (TGF β) [31[•]], and this response is constant up to 72 h of exposure. In another study, UFPs showed cytoskeletal dysfunction in macrophages [32[•]]. The balance between proinflammatory/antiinflammatory effects and an inefficient clearance of particles (due to dysfunction of macrophages) could lead to chronic oedema.

The upregulation of IL-4 and interferon γ (IFN γ), and the expression of CD69 on T cells exposed to particulate matter could explain exacerbations of asthma [33[•]]. In a recent study, epithelial cells exposed to particulate matter in the presence of a supernatant of previously exposed alveolar macrophages to the same particles led to a stronger upregulation of several proinflammatory factors [RANTES, TNF α , intercellular adhesion molecule-1 (ICAM-1), IL-1 β , membrane cofactor protein, leukemia inhibitory factor and vascular endothelial growth factor] and an enhanced activity of the nuclear factor κB when compared with the response to particulate matter alone [34] correlating with TNF α and IL-1 β . Later on, the same group [35^{••}] reported that cocultures of alveolar macrophages with epithelial cells have a wide response to particulate matter presenting an upregulation in the expression of macrophage inflammatory protein 1 β (MIP-1 β), granulocyte-macrophage colony stimulating factor (GM-CSF) and IL-6 after 2 h and in the secretion of MIP-1 β , GM-CSF, M-CSF and IL-6 after 24 h of exposure. Interestingly, the interaction between alveolar macrophages and epithelial cells seems to be independent of cross-linking between these two cells.

Recently, toll-like receptor-2 (TLR-2) and TLR-4 have been identified as candidates of how cells interact with particulate matter [36^{••}]. According to this report, the secretion of IL-8 by airway epithelial cells can be inhibited when an antibody against TLR-2 is added, but no effect is observed on alveolar macrophages using the same antibody. In contrast, the response of macrophages to particulate matter seems to be related to TLR-4 but is not affected by the inhibition of TLR-2. These findings are important considering that biological components of the particles have been previously related to their effects [27,37].

Composition of particulate matter and its relation to biologic effects

During the 1990s the toxicity of particulate matter was mostly related to size, and variations in composition were considered less important. Nevertheless, many publications have indicated that composition plays a

very important role on particle toxicity [25–27], and the observed differences were empirically related to endotoxin, metals and organic compounds. Differences in composition related to particulate size, city and season have recently been reported [38[•]]. The variations in composition have more impact on the effects (TNF α and IL-6 release) induced by the coarse fraction than the differences in surface area. Independently, a study of different soil-derived particulate matter also presented a high correlation between differences in composition (organic carbon and elemental carbon content) and the secretion of IL-6 and IL-8 in BEAS-2B cells [39^{••}].

Different statistical approaches may be used to evaluate the importance of independent components of particulate matter. The use of multiple regression analysis followed by a stepwise analysis showed that PM₁₀ from different regions of Mexico City are mainly related to nitrogen with regard to cytotoxicity, and to lead with regard to proinflammatory effects (E. Alfaro-Moreno, S. Ponce-de-León-Rosales, A.R. Osornio-Vargas, *et al.*, in preparation). In another study using principal component analysis and factorial analysis, independent combinations of components were found to correlate with viability (S/K/Ca/Ti/Mn/Fe/Zn/Pb), TNF α and IL-6 secretion (S/K/Ca/Ti/Mn/Fe/Zn/Pb and endotoxins/organic carbon/black carbon), and p53 expression (high levels of endotoxins/organic carbon/black carbon and low levels of S/K/Ca/Ti/Mn/Fe/Zn/Pb) [40]. In a study evaluating the secretion of cytokines and reactive oxygen species (ROS) induced by particulate matter collected in different months [41^{••}], correlations of IL-6 with Fe and Si were observed, while IL-8 correlated with Cr. In a comprehensive evaluation of the effects of engine emissions (134 components) on 15 different biological outcomes, it was found that re-grouping components by chemical class or subclass allowed for the correlation of lung toxicity with organic carbon, hopanes and steranes; mutagenicity correlated with particle-bound nitrated polycyclic aromatic hydrocarbons [42].

In relation to the genotoxic effects of particulate matter as a possible factor for developing lung cancer (see above), ROS play an important role in DNA damage [43[•]]. Oxidative DNA damage evaluated on circulating mononuclear cells of healthy nonsmokers was related to outdoor–indoor exposure to UFPs, but no correlation was found with other gaseous pollutants, even considering that correlations among them exist [44^{••}]. An in-vitro study shows that the organic fraction and the washed fraction are less potent inducers of DNA damage than the total particles [45^{••}]. Mutagenic compounds, mainly associated with particulate matter from urban or rural regions (related to a chemical plant) were demonstrated using the Ames test [46[•]].

Cardiovascular effects of particulate matter

Epidemiologists have linked air pollution with acute and chronic cardiovascular endpoints [47[•]]. The plausibility of these observations, however, was questionable because little or no experimental studies provided evidence for effects of particulates outside the lungs (Fig. 2).


Acute cardiovascular effects

Dozens of studies have investigated the acute effects of peaks of air pollution [3]. Initially these studies used time-series analyses relating day-to-day variations in atmospheric pollutants with daily variations in deaths or disease. Weather-related differences over the seasons, however, may modify or even confound the association between air pollution and mortality, because both high and low temperatures increase mortality, and temperature is closely associated with pollution. Due to these complex relations, simple statistical adjustments may be inadequate and some effects of pollutants may be associated primarily with outdoor temperature. A recent analysis [48] of Belgian mortality data, stratified across seasons and across quintiles of outdoor temperature, revealed that the short-term effect of PM₁₀ on mortality strongly depends on season or outdoor temperature, even in a temperate climate. Thus, the percentage increases in total mortality on days in the highest season-specific PM₁₀ quartile versus the lowest season-specific PM₁₀ quartile were 7.8% in summer, 6.3% in spring, 2.2% in autumn and 1.4% in winter. This pattern was also observed for cardiovascular mortality. The component-specific toxicity of PM₁₀ may differ across the temperature range as suggested in an in-vitro study [38[•]] in which PM₁₀ samples collected in the summer were more potent to induce inflammatory cytokines (IL-6 and TNF α) than samples collected in winter.

At the level of the population, acute effects of air pollution on cardiovascular endpoints have been noted to be significant from middle age on. The highest risk appears to be in people with preexisting cardiac disease [49]. Drugs that modify oxidant defences may influence the susceptibility to particle-induced inflammatory or pro-oxidative responses. In addition to their cholesterol-lowering effect, statins also have potent antiinflammatory properties. Schwartz and colleagues [50^{••}] found that the effect of PM₁₀ on heart rate variability was confined to persons missing the allele for *GSTM1* (lower oxidative stress defence), but the association was only apparent in those who were not under statin treatment.

Chronic cardiovascular effects

In the Harvard Six Cities study [14] the adjusted overall mortality rate for the most polluted city versus the least polluted city was 1.26. Cardiovascular deaths accounted for the largest single category of the increased mortality. Among air pollutants, elevation of PM_{2.5} was most

Figure 2 Diagram of the possible mechanisms related to systemic effects induced by ultrafine particles (UFPs)

Proinflammatory factors secretion, particle translocation, activation of endothelial cells and prothrombotic effects. ICAM, intercellular adhesion molecule; IL, interleukin; PECAM, platelet–endothelial adhesion molecule; TNF, tumor necrosis factor.

strongly associated with mortality. Pope *et al.* [51] linked mortality statistics over a 16-year period to chronic exposure to multiple air pollutants in approximately 500 000 adults who resided in all 50 states of the USA. Mean annual PM_{2.5} concentration linearly enhanced cardiovascular mortality, with a significant increase in the risk of death from ischaemic heart disease of 1.18 for an increase in PM_{2.5} of 10 µg/m³. The risks for arrhythmia, heart failure, or cardiac arrest mortality were also increased [relative risk (RR) 1.13]. Intriguingly (and hardly discussed in that article), air pollution was associated with a significantly lower risk of dying from COPD (RR 0.84), suggesting either a wrong coding of the cause of death in COPD patients or that patients with COPD are more susceptible to die from cardiovascular complications than from COPD when they have been exposed to high pollution.

Extrapulmonary translocation of ultrafine particles

The impact of inhaled particles on the autonomic nervous system leading to changes in the pattern of

breathing, heart rate and heart rate variability is under investigation. Elder *et al.* [52^{••}] demonstrated that the olfactory neuronal pathway represents a significant exposure route of central nervous system (CNS) tissue to inhaled UFPs. These authors showed that, in rats, which are obligatory nose breathers, translocation of inhaled nanosized particles along neurons is a more efficient pathway to the CNS than via the blood circulation across the blood–brain barrier. Given that this neuronal translocation pathway was also demonstrated in nonhuman primates, it is likely to be operative in humans as well [52^{••},53].

In relation to the possibility that UFPs translocate from the lungs into the blood circulation, various studies have been conducted in different animal models. The amount of UFPs that translocated into blood and extrapulmonary organs differed amongst these studies [54–57]. Recent studies [58^{••},59,60] have provided morphological data illustrating that inhaled particles are transported into the pulmonary capillary space,

presumably by transcytosis. It has also been shown that, following intranasal delivery, polystyrene microparticles (1.1 μm) can translocate to tissues in the systemic compartment [61].

In an ex-vivo model of isolated perfused rat lungs, lacking both lymph flow and recruited inflammatory cells [62[•]], it was confirmed that UFPs can translocate from the lung into the circulation, when pulmonary microvascular permeability was increased by pharmacological mediation (H_2O_2 , vascular histamine administration). Using a similar model in rabbit lungs, we found a passage of UFPs from the blood vessels to the alveolar spaces, but only after histamine administration [63[•]].

The issue of particle translocation in humans is still controversial. In a recent study using an aerosol of technetium-99m-labelled carbon particles (10 nm in diameter, but these rapidly formed aggregates of 100 nm in diameter in the inhaled aerosol), no translocation into the systemic circulation could be demonstrated, and no significant radioactivity was found over the liver. The nature of the radioactivity found in blood (4.4%) consisted mainly of pertechnetate, as analysed by thin layer chromatography [64[•]]. These results contrast with previous findings also based on inhaling an aerosol of technetium-99m-labelled carbon particles [65], in which particle-bound radioactivity (also assessed by thin-layer chromatography) was detected in blood after 1 min, reaching a maximum between 10 and 20 min, and remaining at this level up to 60 min. Gamma camera images showed substantial radioactivity over the liver and other areas of the body. The discrepancies between these two studies may be related to the chromatography technique or the size of the inhaled particles. Further studies with other types of radioactive labelling should clarify the issue.

Effect of particles on thrombogenesis

The effects of particles on thrombosis constitute a relevant cardiovascular endpoint. Using a hamster model of vascular thrombosis in the femoral vein, we found a marked increase in prothrombotic tendency after intravenous injection of positively charged (amine-modified) polystyrene particles of 60 nm diameter, which resulted, at least in part, from platelet activation [66]. Similar effects were obtained after the intratracheal administration of these positively charged polystyrene particles [67]. In complete agreement with our findings, Silva *et al.* [68[•]], who used a rat model of ear vein thrombosis, also observed a significant dose-dependent enhancement of thrombus formation after intravenous and intratracheal administration of positively charged polystyrene particles but not with negatively charged polystyrene particles of the same size.

Using real pollutant particles, that is DEPs, we showed a marked pulmonary inflammation, enhanced venous and arterial thrombosis and platelet activation within 1 h of their deposition in the lungs [69]. The prothrombotic tendency, activation of circulating blood platelets, as well as lung inflammation persisted up to 24 h after instillation [70]. Pretreatment with an H1-receptor antagonist prevented the effects at 6 and 24 h, but the prothrombotic tendency observed 1 h after DEP exposure did not appear to correlate with pulmonary inflammation. The latter is compatible with direct platelet activation by DEPs (or their constituents), having penetrated into the circulation [70]. In subsequent experiments, we found that 24 h after DEP exposure, pretreatment with dexamethasone or with cromoglycate blocked the DEP-induced pulmonary inflammation, prothrombotic events and histamine release in bronchoalveolar lavage fluid and plasma. We concluded that the systemic inflammatory and prothrombotic effects found 24 h after DEP administration are secondary to lung inflammation, and that they can be prevented by mast cell stabilization [71]. More recently, using silica particles, we provided novel evidence for a critical role of macrophage–neutrophil cross-talk during lung inflammation induced by particles, leading to the release of neutrophil elastase into the systemic circulation. We found that neutrophil enzymes may be responsible for the priming of platelet activation and contribute to the development of a thrombotic tendency, when such primed platelets encounter a (mildly) injured vessel wall [72^{••}]. Although the mechanism involved in the effect of silica particles is different from that triggered by DEPs, this study provided further evidence that inflammation is an integral component of thrombosis.

In support of this concept, it has been shown, using repeated exposures to particulate air pollution, that PM_{10} increased the volume and advanced the phenotype of the coronary atherosclerotic lesions in Watanabe heritable hyperlipidaemic rabbits [73]. Moreover, Sun *et al.* [74^{••}] showed, in an apoE^{−/−} mouse model, that long-term exposure to low concentrations of $\text{PM}_{2.5}$ altered vasomotor tone, induced vascular inflammation, and potentiated atherosclerosis.

Some in-vitro studies have evaluated the possible mechanisms of translocation and the vascular effects of particles. Geys *et al.* [75[•]] showed that nanoparticles are capable of translocating from the apical to the basolateral side of a confluent culture of Calu-3 cells. Studies with human vein endothelial cells have shown that urban particles are able to induce the expression of adhesion molecules such as E-selectin, ICAM-1, vascular cell adhesion molecule 1 and platelet–endothelial adhesion molecule 1 [76,77]. These effects appear to be partially related to the presence of endotoxin in the insoluble

fraction [76], while the PM₁₀ particles seem to induce a stronger effect than the PM_{2.5} fraction [77]. Urban particles were also shown to induce oxidative stress (H₂O₂ production, activation of extracellular signal regulated kinase 1/2 and p38 mitogen-activated protein kinase) in human pulmonary endothelial cells [78**].

Conclusion

The adverse effects of exposure to particulate matter are manifested from early life onwards and at different levels. So far, the effects on lung function, COPD, allergic illness and lung cancer have been reasonably reproduced at an experimental level with regard to systemic, cellular and molecular effects. The importance of particulate matter translocation, both *in vivo* and *in vitro* has become a hot topic. The interactions between different cells have shown that the previous approach studying single cell lines may underestimate the effect of particulate matter. Regardless of the great effort to explain the impact of particulate matter on public health and at the experimental level, we are facing a great challenge to control the emission and composition of particulate matter. This is of special interest in mega cities, like Mexico City, presenting an average level of PM₁₀ of 50 µg/m³ influenced by industry, traffic and natural sources.

Acknowledgements

EAM is supported by European Respiratory Society, Fellowship number 80. TSN is a fellow of the Flemish Scientific Fund (FWO).

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

Additional references related to this topic can also be found in the Current World Literature section in this issue (p. 143).

- 1 Nemery B, Hoet PHM, Nemmar A. The Meuse Valley fog of 1930: an air pollution disaster. *Lancet* 2001; 357:704–708.
- 2 Logan WPD. Mortality in the London fog incident. *Lancet* 1953; 261:336–338.
- 3 Brunekreef B, Holgate ST. Air pollution and health. *Lancet* 2002; 360:1233–1242.
- 4 Aust AE, Ball JC, Hu AA, *et al.* Particle characteristics responsible for effects on human lung epithelial cells. *HEI Research Report* 110; 2002.
- 5 Pope AC III, Dockery DW. Health effects of fine particulate air pollution: lines that connect. *J Air Waste Manage Assoc* 2006; 56:709–742.
- A very comprehensive review of more than 500 articles.
- 6 Lacasana M, Esplugues A, Ballester F. Exposure to ambient air pollution and prenatal and early childhood health effects. *Eur J Epidemiol* 2005; 20:183–199.
- This article reviews publications from 1999 to 2003, which have evaluated the possible impact of air pollution on the fetus and the first year of life.
- 7 Kulkarni N, Pierse N, Rushton L, Grigg J. Carbon in airway macrophages and lung function in children. *N Engl J Med* 2006; 355:21–30.
- Cross-sectional study using carbon in airway macrophages as exposure variable. This study is very original by using an internal marker of external exposure at an individual level.
- 8 Gauderman WJ, Avol E, Gilliland F, *et al.* The effect of air pollution on lung development from 10 to 18 years of age. *N Engl J Med* 2004; 351:1057–1067.
- 9 Riedl M, Diaz-Sanchez D. Biology of diesel exhaust effects on respiratory function. *J Allergy Clin Immunol* 2005; 115:221–228.

- 10 Gilliland FD, Li YF, Saxon A, Diaz-Sanchez D. Effect of glutathione-S-transferase M1 and P1 genotypes on xenobiotic enhancement of allergic responses: randomised, placebo-controlled crossover study. *Lancet* 2004; 363:119–125.
- 11 Polosa R, Salvi S, Di Maria GU. Allergic susceptibility associated with diesel exhaust particle exposure: clear as mud. *Arch Environ Health* 2002; 57:188–193.
- 12 Atkinson RW, Anderson HR, Sunyer J, *et al.* Acute effects of particulate air pollution on respiratory admissions: results from Aphea 2 project: Air Pollution and Health: A European Approach. *Am J Respir and Crit Care Med* 2001; 164:1860–1866.
- 13 Schikowski T, Sugiri D, Ranft U, *et al.* Long-term air pollution exposure and living close to busy roads are associated with COPD in women. *Respir Res* 2005; 6:152.
- An original prospective investigation on the development of COPD in association with chronic air pollution.
- 14 Dockery DW, Pope CA III, Xu X, *et al.* An association between air pollution and mortality in six US cities. *N Engl J Med* 1993; 329:1753–1759.
- 15 Vineis P, Hoek G, Krzyzanowski M, *et al.* Air pollution and risk of lung cancer in a prospective study in Europe. *Int J Cancer* 2006; 119:169–174.
- A European multicentre nested-case control study on the association between chronic traffic-related exposure and lung cancer risk.
- 16 Beeson WL, Abbey DA, Knutsen SF. Long-term concentrations of ambient air pollutants and incident lung cancer in California adults: results from the AHSOG study. *Environ Health Perspect* 1998; 106:813–823.
- 17 Pope CA, Burnett RT, Thun MJ, *et al.* Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *JAMA* 2002; 287:1132–1141.
- 18 Hoek G, Brunekreef B, Goldbohm S, *et al.* Association between mortality and indicators of traffic-related air pollution in the Netherlands: a cohort study. *Lancet* 2002; 360:1203–1209.
- 19 Nafstad P, Haheim LL, Oftedal B, *et al.* Lung cancer and air pollution: a 27 year follow up of 16209 Norwegian men. *Thorax* 2003; 58:1071–1076.
- 20 Manuguerra M, Matullo G, Veglia F, *et al.* Multifactor dimensionality reduction applied to a large prospective investigation on gene–gene and gene–environment interactions. *Carcinogenesis* 2006; 6 September [Epub ahead of print].
- Shows a gene–environment interaction between polymorphisms of DNA repair and the distance of living from a major road.
- 21 Nawrot TS, Nackaerts K, Hoet PH, Nemery B. Lung cancer mortality and fine particulate air pollution in Europe. *Int J Cancer* (in press).
- 22 Moon C, Becker S. Cytotoxicity and induction of proinflammatory cytokines from human monocytes exposed to fine (PM_{2.5}) and coarse (PM₁₀₋₂₅) in outdoor and indoor air. *Toxicol Appl Pharmacol* 1999; 155:245–252.
- 23 Holian A, Hamilton RF, Morandi MT, *et al.* Urban particle induced apoptosis and phenotype shifts in human alveolar macrophages. *Environ Health Perspect* 1998; 106:127–132.
- 24 Obot CJ, Morandi MT, Beebe TP, *et al.* Surface components of airborne particulate matter induce macrophage apoptosis through scavenger receptors. *Toxicol Appl Pharmacol* 2002; 184:98–106.
- 25 Alfaro-Moreno E, Martínez L, García-Cuellar C, *et al.* Biologic effects induced in vitro by PM₁₀ from three different zones of Mexico City. *Environ Health Perspect* 2002; 110:715–720.
- 26 Osornio-Vargas AR, Bonner JC, Alfaro-Moreno E, *et al.* Proinflammatory and cytotoxic effects of Mexico City air pollution particulate matter in vitro are dependent on particle size and composition. *Environ Health Perspect* 2003; 111:1289–1293.
- 27 Bonner JC, Rice AB, Lindroos PM, *et al.* Induction of the lung myofibroblast PDGF receptor system by urban ambient particles from Mexico City. *Am J Respir Cell Mol Biol* 1998; 19:672–680.
- 28 Villalobos R, Blanco S, Gomez S. Mutagenicity assessment of airborne particles in Mexico City. *Atmospheric Environ* 1995; 29:517–524.
- 29 García-Cuellar C, Alfaro-Moreno E, Martínez-Romero F, *et al.* DNA damage induced by PM₁₀ from different zones of Mexico City. *Ann Occup Hygiene* 2002; 46 (Suppl 1):425–428.
- 30 Wise H, Balharry D, Reynolds LJ, *et al.* Conventional and toxicogenomic assessment of the acute pulmonary damage induced by the instillation of Cardiff PM₁₀ into the rat lung. *Sci Total Environ* 2006; 360:60–67.
- This paper shows that 3 days after instillation, rats present a mild oedematous response.

- 31 Dagher Z, Garçon G, Gosset P, *et al.* Pro-inflammatory effects of Dunkerque City air pollution particulate matter 2.5 in human epithelial lung cells (L132) in culture. *J Appl Toxicol* 2005; 25:166–175.
In this paper, the authors present an evaluation of urban particle effects in a human lung epithelial cell line under proliferative conditions, showing that the proinflammatory response lasts up to 3 days after exposure.
- 32 Möller W, Brown DM, Kreyling WG, Stone V. Ultrafine particles cause cytoskeletal dysfunctions in macrophages: role of intracellular calcium. *Particle Fibre Toxicol* 2005; 2:7.
This study gives evidence of the mechanism related to cytoskeletal dysfunction in macrophages exposed to particulate matter.
- 33 Mamessier E, Nieves A, Vervloet D, Magnan A. Diesel exhaust particles enhance T-cell activation in severe asthmatics. *Allergy* 2006; 61:581–588.
This paper provides evidence of the possible mechanism involved in exacerbation of asthma related to particulate matter exposure.
- 34 Ishii H, Fuji T, Hogg JC, *et al.* Contribution of $\text{IL-1}\beta$ and $\text{TNF}\alpha$ to the initiation of the peripheral lung response to atmospheric particulates (PM10). *Am J Physiol Lung Cell Mol Physiol* 2004; 287:L176–L183.
- 35 Ishii H, Hayashi S, Hogg JC, *et al.* Alveolar macrophage–epithelial cell interaction following exposure to atmospheric particles induces the release of mediators involved in monocyte mobilization and recruitment. *Respir Res* 2005; 6:87.
This is a study presenting the interactions between macrophages and epithelial cells in relation to the cytokine secretion profile after exposure to particulate matter.
- 36 Becker S, Dailey L, Soukup JM, *et al.* TLR-2 is involved in airway epithelial cell response to air pollution particles. *Toxicol Appl Pharmacol* 2005; 203:45–52.
This is an excellent study providing strong evidence of the role of TLR-2 in the cellular response to particulate matter.
- 37 Becker S, Fenton MJ, Soukup JM. Involvement of microbial components and toll-like receptors 2 and 4 in cytokine responses to air pollution particles. *Am J Respir Cell Mol Biol* 2002; 27:611–618.
- 38 Hetland RB, Cassee FR, Låg M, *et al.* Cytokine release from alveolar macrophages exposed to ambient particulate matter: heterogeneity in relation to size, city and season. *Particle Fibre Toxicol* 2005; 2:4.
Describes the differences of in-vitro effects of particles related to the city in which the particles were collected.
- 39 Veranth JM, Moss TA, Chow JC, *et al.* Correlation of in vitro cytokine responses with the chemical composition of soil-derived particulate matter. *Environ Health Perspect* 2006; 114:341–349.
An excellent article presenting the role of composition on the biologic effect of soil-derived particulate matter.
- 40 Rosas-Pérez I, Serrano J, Alfaro-Moreno E, *et al.* Relations between PM10 composition and cell toxicity: a multivariate and graphical approach. *Chemosphere* (in press).
- 41 Becker S, Dailey L, Soukup JM, *et al.* Seasonal variations in air pollution particle-induced inflammatory mediator release and oxidative stress. *Environ Health Perspectives* 2005; 113:1032–1038.
An excellent article that deals with differences in composition related to seasonal variations and different biologic effects.
- 42 McDonald JD, Eide I, Seagrave J, *et al.* Relationship between composition and toxicity of motor vehicle emission samples. *Environ Health Perspect* 2004; 112:1527–1538.
- 43 Risom L, Møller P, Loft S. Oxidative stress-induced DNA damage by particulate air pollution. *Mutation Res* 2005; 592:119–137.
A comprehensive review of the evidence that correlates particulate matter with DNA damage.
- 44 Vinzents PS, Møller P, Sørensen M, *et al.* Personal exposure to ultrafine particles and oxidative DNA damage. *Environ Health Perspect* 2005; 113:1485–1490.
A very good article presenting evidence that particulate matter correlates with oxidative stress and DNA damage, but other pollutants do not.
- 45 Healey K, Lingard JJN, Tomlin S, *et al.* Genotoxicity of size-fractionated samples of urban particulate matter. *Environ Mol Mutagen* 2005; 45:380–387.
The data presented in this article strongly indicate that the genotoxic effects of particulate matter may be related to interactions among particulate matter components. This is concluded considering that fractions isolated from the total particles had a less strong genotoxic effect.
- 46 Du Four VA, Janssen CR, Brits E, Larebeke NV. Genotoxic and mutagenic activity of environmental air samples from different rural, urban and industrial sites in Flanders. *Belgium Mutation Res* 2005; 588:106–117.
This study presents evidence of the mutagenic effects of air samples using the Ames test.
- 47 Nawrot TS, Nemmar A, Nemery B. Air pollution: to the heart of the matter. *Eur Heart J* 2006; 27:2269–2271.
Brief overview of the air pollution epidemiology.
- 48 Nawrot TS, Torfs R, Fierens F, *et al.* Stronger associations between daily mortality and fine particulate air pollution in summer than in winter: evidence from a heavily polluted region in western Europe. *J Epidemiol Comm Health* (in press).
- 49 Goldberg MS, Burnett RT, Bailar JC, *et al.* Identification of persons with cardiorespiratory conditions who are at risk of dying from the acute effects of ambient air particles. *Environ Health Perspect* 2001; 109 (Suppl 4):487–494.
- 50 Schwartz J, Park SK, O'Neill MS, *et al.* Glutathione-S-transferase M1, obesity, statins, and autonomic effects of particles: gene-by-drug-by-environment interaction. *Am J Respir Crit Care Med* 2005; 172:1529–1533.
Original concept that shows in an epidemiological context the role of the susceptibility to inflammation as an effect modifier in the association between heart rate variability and air pollution.
- 51 Pope CA III, Burnett RT, Thurston GD, *et al.* Cardiovascular mortality and long-term exposure to particulate air pollution: epidemiological evidence of general pathophysiological pathways of disease. *Circulation* 2004; 109: 71–77.
- 52 Elder A, Gelein R, Silva V, *et al.* Translocation of inhaled ultrafine manganese oxide particles to the central nervous system. *Environ Health Perspect* 2006; 114:1172–1178.
This is a very original study demonstrating that the olfactory neuronal pathway is efficient for translocating inhaled UFPs to the CNS and that this can result in inflammatory changes.
- 53 Oberdorster G, Sharp Z, Atudorei V, *et al.* Translocation of inhaled ultrafine particles to the brain. *Inhal Toxicol* 2004; 16:437–445.
- 54 Kreyling W, Semmler M, Erbe F, *et al.* Translocation of ultrafine insoluble iridium particles from lung epithelium to extrapulmonary organs is size dependent but very low. *J Toxicol Environ Health A* 2002; 65:1513–1530.
- 55 Nemmar A, Vanbilloen H, Hoylaerts MF, *et al.* Passage of intratracheally instilled ultrafine particles from the lung into the systemic circulation in hamster. *Am J Respir Crit Care Med* 2001; 164:1665–1668.
- 56 Oberdorster G, Sharp Z, Atudorei V, *et al.* Extrapulmonary translocation of ultrafine carbon particles following whole-body inhalation exposure of rats. *J Toxicol Environ Health A* 2002; 65:1531–1543.
- 57 Takenaka S, Karg E, Roth C, *et al.* Pulmonary and systemic distribution of inhaled ultrafine silver particles in rats. *Environ Health Perspect* 2001; 109 (Suppl 4):547–551.
- 58 Geiser M, Rothen-Rutishauser B, Kapp N, *et al.* Ultrafine particles cross cellular membranes by nonphagocytic mechanisms in lungs and in cultured cells. *Environ Health Perspect* 2005; 113:1555–1560.
This study reports that inhaled ultrafine titanium dioxide particles can be found on the luminal side of airways and alveoli, in all major lung tissue compartments and cells, and within capillaries, and the particle uptake *in vitro* into cells does not occur by endocytosis, but by diffusion or adhesive interactions.
- 59 Kapp N, Kreyling W, Schulz H, *et al.* Electron energy loss spectroscopy for analysis of inhaled ultrafine particles in rat lungs. *Microsc Res Tech* 2004; 63:298–305.
- 60 Kato T, Yashiro T, Murata Y, *et al.* Evidence that exogenous substances can be phagocytized by alveolar epithelial cells and transported into blood capillaries. *Cell Tissue Res* 2003; 311:47–51.
- 61 Eyles JE, Bramwell VW, Williamsson ED, Alpar HO. Microsphere translocation and immunopotential in systemic tissues following intranasal administration. *Vaccine* 2001; 19:4732–4742.
- 62 Meiring J, Borm PJA, Bagate K, *et al.* The influence of hydrogen peroxide and histamine on lung permeability and translocation of iridium nanoparticles in the isolated perfused rat lung. *Part Fibre Toxicol* 2005; 2.
This study reports that permeability of the lung barrier to UFPs is controlled both at the epithelial and endothelial level, and is influenced by the inflammation.
- 63 Nemmar A, Hamoir J, Nemery B, Gustin P. Evaluation of particle translocation across the alveolo-capillary barrier in isolated perfused rabbit lung model. *Toxicology* 2005; 208:105–113.
This study demonstrates that UFPs diffuse from the vascular compartment into the lungs in an isolated perfused rabbit lung model, when the microvascular permeability is increased by histamine.
- 64 Mills NL, Amin N, Robinson SD, *et al.* Do inhaled carbon nanoparticles translocate directly into the circulation in man? *Am J Respir Crit Care Med* 2006; 173:426–431.
This article did not confirm the earlier findings by Nemmar *et al.* [66].
- 65 Nemmar A, Hoet PH, Vanquickenborne B, *et al.* Passage of inhaled particles into the blood circulation in humans. *Circulation* 2002; 105: 411–414.
- 66 Nemmar A, Hoylaerts M, Hoet PHM, *et al.* Ultrafine particles affect experimental thrombosis in an *in vivo* hamster model. *Am J Respir Crit Care Med* 2002; 166:998–1004.

- 67** Nemmar A, Hoylaerts MF, Hoet PHM, *et al.* Size effect of intratracheally instilled particles on pulmonary inflammation and vascular thrombosis. *Toxicol Appl Pharmacol* 2003; 186:38–45.
- 68** Silva VM, Corson N, Elder A, Oberdorster G. The rat ear vein model for investigating in vivo thrombogenicity of ultrafine particles (Ufp). *Toxicol Sci* 2005; 85:983–989.
- This is an excellent study that validated the use of a noninvasive ear vein model in rats to evaluate the thrombogenic effects of UFP deposition in the respiratory tract.
- 69** Nemmar A, Hoet PH, Dinsdale D, *et al.* Diesel exhaust particles in lung acutely enhance experimental peripheral thrombosis. *Circulation* 2003; 107:1202–1208.
- 70** Nemmar A, Nemery B, Hoet PHM, *et al.* Pulmonary inflammation and thrombogenicity caused by diesel particles in hamsters – role of histamine. *Am J Respir Crit Care Med* 2003; 168:1366–1372.
- 71** Nemmar A, Hoet PHM, Vermeylen J, *et al.* Pharmacological stabilization of mast cells abrogates late thrombotic events induced by diesel exhaust particles in hamsters. *Circulation* 2004; 110:1670–1677.
- 72** Nemmar A, Nemery B, Hoet PH, *et al.* Silica particles enhance peripheral thrombosis: key role of lung macrophage-neutrophil cross-talk. *Am J Respir Crit Care Med* 2005; 171:872–879.
- This is a mechanistic study showing the importance of pulmonary macrophage–neutrophil cross-talk, in releasing neutrophil elastase into the blood, and causing platelet activation and thrombotic events on mildly damaged vasculature.
- 73** Suwa T, Hogg JC, Quinlan KB, *et al.* Particulate air pollution induces progression of atherosclerosis. *J Am Coll Cardiol* 2002; 39:935–942.
- 74** Sun Q, Wang A, Jin X, *et al.* Long-term air pollution exposure and acceleration of atherosclerosis and vascular inflammation in an animal model. *JAMA* 2005; 294:3003–3010.
- An outstanding article showing, in an apoE^{−/−} mouse model, that long-term exposure to low concentration of PM_{2.5} altered vasomotor tone, induced vascular inflammation, and potentiated atherosclerosis.
- 75** Geys J, Coenegrachts L, Vercammen V, *et al.* In vitro study of the pulmonary translocation of nanoparticles: a preliminary study. *Toxicol Lett* 2006; 160:218–226.
- An interesting preliminary study on in-vitro translocation of UFPs, reporting that the Calu-3 cell line can be used in an in-vitro translocation study.
- 76** Alfaro-Moreno E, López-Marure R, Montiel-Dávalos A, *et al.* E-selectin expression in human endothelial cells exposed to PM₁₀: the role of endotoxin and insoluble fraction. *Environ Res* (in press).
- 77** Montiel-Dávalos A, Alfaro-Moreno E, López-Marure R. PM_{2.5} and PM₁₀ induce the expression of adhesion molecules and the adhesion of monocytic cells to human umbilical vein endothelial cells. *Inhalation Toxicol* (in press).
- 78** Li Z, Hyseni X, Carter JD, *et al.* Pollutant particles enhanced H₂O₂ production from NAD(P)H oxidase and mitochondria in human pulmonary artery endothelial cells. *Am J Physiol Cell Physiol* 2006; 291:C357–C365.
- An excellent mechanistic study of the effects of urban particles on pulmonary artery endothelial cells.