

PAPER

Chloroquine increases low-density lipoprotein removal from plasma in systemic lupus patients

JC Sachet, EF Borba*, E Bonfá, CGC Vinagre, VM Silva and RC Maranhão

Rheumatology Division and Lipid Metabolism Laboratory of the Heart Institute (InCor) of the Medical School Hospital and Faculty of Pharmaceutical Sciences, University of São Paulo, São Paulo, Brazil

Low-density lipoprotein (LDL) pathway in systemic lupus erythematosus (SLE) patients taking chloroquine diphosphate (CDP) was evaluated through the kinetic behavior of a radioactive cholesterol-rich nanoemulsion (LDE) that resembles the LDL lipidic structure. LDE was labeled with ^{14}C -cholesteryl ester (^{14}C -CE), then IV injected in inactive female SLE patients: 10 taking CDP (CDP), 10 without therapy (NO THERAPY); and 10 normal subjects (CONTROL). Groups were age-matched and followed rigorous selection criteria of conditions that interfere in the lipid profile. Blood samples were collected in pre-established intervals after infusion for radioactivity measurement. Fasting lipoproteins were determined in the beginning of kinetic studies. Fractional clearance rate (FCR) of ^{14}C -CE was significantly different in the three groups ($P = 0.03$). In fact, a greater FCR of ^{14}C -CE was observed in CDP compared to NO THERAPY (0.076 ± 0.037 versus $0.046 \pm 0.021 \text{ h}^{-1}$; $P < 0.05$) and to CONTROL ($0.0516 \pm 0.0125 \text{ h}^{-1}$; $P < 0.05$). Accordingly, a significant lower total and LDL cholesterol were observed in CDP (156 ± 16 and $88 \pm 16 \text{ mg/dl}$) compared to NO THERAPY (174 ± 15 and $108 \pm 17 \text{ mg/dl}$; $P < 0.05$) and to CONTROL (200 ± 24 and $118 \pm 23 \text{ mg/dl}$; $P < 0.05$). In contrast, no difference in (FCR) of ^{14}C -CE of NO THERAPY and CONTROL groups was observed. This is the first *in vivo* demonstration that LDE removal by LDL receptor from plasma is increased in SLE patients taking CDP with a consequent beneficial decrease in LDL-c levels. *Lupus* (2007) 16, 273–278.

Key words: atherosclerosis; chloroquine; dyslipoproteinemia; emulsions; lipoproteins; low-Density lipoprotein; systemic lupus erythematosus

Introduction

Antimalarial drugs containing the four amino-quinilone radicals have a modulating effect on the inflammatory process in rheumatic diseases.^{1–3} In fact, hydroxychloroquine (HCQ) and chloroquine diphosphate (CDP) are widely used in systemic lupus erythematosus (SLE).^{4–8}

Furthermore, antimalarials have a plasma lipid lowering effect in SLE^{9–11} that is therapeutically relevant due to the increased risk of premature atherosclerosis in this disease.¹² In fact, dyslipidemias are very frequent in SLE^{12–14} and certainly play a pivotal role in the 50 times greater risk of developing coronary artery

disease (CAD),¹⁵ an important cause of mortality in SLE.^{12,16,17}

SLE dyslipoproteinemias are known to be aggravated by disease activity^{18–21} and therapy.^{22–24} The latter is characterized by an increased total and LDL cholesterol, a dyslipidemia pattern observed in more than half of lupus patients.^{12–14} Importantly, LDL is recognized as one of the main risk factor for atherosclerosis²⁵ and therefore the reported beneficial effect of antimalarials in reducing this lipoprotein^{9,10,26–31} is of great interest.

Since LDL removal by its specific receptor is highly effective in decreasing this lipoprotein fraction,³² we evaluated the LDL pathway in lupus patients with or without CDP and in controls through the determination of plasma kinetics of a radioactively labeled artificial cholesterol-rich nanoemulsion (LDE) that resembles LDL lipidic structure.^{33–35} LDE has been used as a surrogate for native LDL to investigate in clinical studies the process of LDL removal from plasma.^{36–38}

Correspondence: Dr EF Borba, Faculdade de Medicina USP, Reumatologia, Av. Dr. Arnaldo, 455, 3º andar, Sala 3133, São Paulo, CEP:01246-903, SP, Brazil. E-mail: reumato@edu.usp.br
Received 25 October 2006; accepted 31 January 2007

Patients and methods

Study population

Twenty consecutive female SLE patients were selected in the Outpatient Lupus Clinics at the Rheumatology Division of the University of São Paulo. All patients were under 50 years old and fulfilled four or more of the revised American College of Rheumatology criteria for the classification of SLE.³⁹ SLE patients were divided in two age-matched groups according to current therapy at the time of evaluation: CDP – exclusive use of 250 mg of CDP daily for at least one year; and NO THERAPY – inactive disease without any CDP drug therapy or have stopped this drug for at least one year. SLE patients were not under corticosteroids nor immunosuppressive drugs in the last year, and none of them were in use of any lipid-raising drugs, such as anti-convulsants, anti-hypertensives (beta blockers and diuretics), estrogen and/or progesterone containing agents, or lipid-lowering agents. Exclusion criteria were presence of clinical and/or biochemical evidence of Diabetes Mellitus, CAD, liver and thyroid disease, and history of alcohol use. No patient was pregnant or menopausal. All patients had serum creatinine levels below 1.5 mg/dl and proteinuria less than 0.3 g/L/day. Clinical activity was measured according to the Systemic Lupus Erythematosus Disease Activity Index (SLEDAI)⁴⁰ and all SLE patients have inactive disease with SLEDAI scores of 0. Ten healthy and age-matched female subjects (CONTROL) were selected for comparison and were studied in parallel with lupus patients. None of the SLE patients and controls had a body mass index (BMI) greater than 30 kg/m².

LDE preparation

LDE was prepared from a lipid mixture composed of 40 mg cholesteryl oleate, 20 mg egg phosphatidylcholine, 1 mg triolein and 0.5 mg cholesterol purchased from Nu-Check Prep (Elysian). [¹⁴C] cholesteryl ester purchased from Amersham International (Amersham, UK) was added to the mixture. The emulsification of lipids by prolonged ultrasonic irradiation in aqueous media and the two-step ultracentrifugation procedure of the crude emulsion with density adjustment by addition of KBr to obtain LDE microemulsion was carried out by the method of Ginsburg⁴¹ modified by Maranhão *et al.*³³ LDE was dialysed against saline solution and passed through 0.22 µm filter for injection into the patients.

LDE plasma kinetics

The participants were fasting for 12 h at the beginning of the test at approximately 8:30 a.m., but they were

allowed two standard meals (total of 1800 kcal) during the study at ~12:30 p.m. and 7 p.m. LDE containing 37 kBq [¹⁴C]cholesteryl ester in a total lipid mass of 5–6 mg (in a 200 µl volume) was intravenously injected in a bolus. Plasma samples were collected during 24 h, in intervals of 5 min, 1, 2, 4, 6, 8 and 24 h after the injection. Aliquots (1 ml) of blood plasma were transferred to counting vials containing 7.0 ml scintillation solution (ULTIMA GOLD XR, Packard Bioscience, Meriden, USA). Radioactivity was counted using a Packard 1660 TR (Meriden, USA) Liquid Scintillation Analyser.

Calculation of FCR of the emulsion radioactive lipids

FCR of the LDE [¹⁴C]cholesteryl ester, was calculated according to the method described by Matthews as FCR, where a_1 , a_2 , b_1 and b_2 were estimated from biexponential curves obtained from the remaining radioactivity found in plasma after injection, fitted by least squares procedure:

$$y = (a_1 \cdot e^{-b_1 t}) + (a_2 \cdot e^{-b_2 t}),$$

where y represents the radioactivity plasma decay in function of time (t); a indicates the linear coefficient and b , the angular coefficient, which represents the FCR in h⁻¹. The FCR were estimated from parameters a_1 , b_1 and b_2 by the following equation: $\text{FCR} = (a_1/b_1 + a_2/b_2)^{-1}$. Calculations were performed by compartmental analysis using the ANACOMP software.⁴²

Informed consent and radiological safety

The experimental protocol was approved by the local Ethical Committee and a written informed consent was given by all participants. A pregnancy test was performed one day before the kinetic study and all tests were negative. The safety of the radioactive dose intravenously injected into the patients was assured according to the regulations of the International Commission on Radiological Safety as described in our previous study.⁴³ The equivalent absorbed dose on each experiment was 0.03 mSV which is below the 50 mSV annual limit for intake of radionuclides.

Lipid profile

All SLE patients and controls were continued with their normal diet and were fasting for at least 12-h at the beginning of the study. Fasting lipoproteins were determined on the same day of the kinetic studies. Plasma total cholesterol (TC) and triglycerides (TGs) were measured enzymatically (Boehringer Mannheim,

Argentina and Merck, Germany, respectively) on a RA 1000 analyser (Technicon Instruments Corp).^{44,45} High density lipoprotein cholesterol (HDL-c) was obtained after precipitation of very LDL cholesterol (VLDL-c) and LDL cholesterol (LDL-c) by phosphotungstic acid and magnesium chloride.⁴⁶ VLDL-c and LDL-c were estimated since all subjects had TGs less than 400 mg/dl.⁴⁷ VLDL-c levels were calculated from the division of serum TG by 5 (TG/5).⁴⁷ LDL-c levels were estimated using the equation:⁴⁷

$$TC = HDL-c + TG/5 + LDL-c$$

Statistical analysis

Results are presented as the mean \pm standard deviation (SD). Data were analysed by one-way analysis of variance (ANOVA) and if a significant result was obtained ($P < 0.05$), Tukey's test was performed to evaluate differences between groups. Correlations among plasma lipid fractions and FCRs were calculated using Spearmann's rank correlation test. Statistical significance was set as $P < 0.05$.

Results

The distribution of age, weight, height and BMI of the three groups is shown in Table 1. As expected, age was similar in CDP, NO THERAPY and CONTROL groups ($P = 0.93$). In addition, no significant difference was found between groups concerning weight ($P = 0.39$) and height ($P = 0.96$). Likewise, BMI did not significantly differ between groups ($P = 0.31$). SLE patients of CDP group were in use of this drug for 7.1 ± 2.9 years and had a similar disease duration compared to NO THERAPY (9.5 ± 4.6 and 8.2 ± 6.6 years, respectively, $P = 0.10$). Importantly, SLE patients of both groups had a SLEDAI score of zero.

Table 1 Demographic features, disease duration and SLEDAI in SLE groups (CDP and NO THERAPY) and CONTROL

Group	CDP (n = 10)	NO THERAPY (n = 10)	CONTROL (n = 10)	P
Age (years)	35.4 ± 7.5	36.5 ± 6.9	35.6 ± 8.9	0.93
Weight (kg)	60.2 ± 6.8	63.0 ± 8.8	57.5 ± 10.4	0.39
Height (m)	1.61 ± 0.04	1.60 ± 0.06	1.60 ± 0.10	0.96
BMI (kg/m ²)	23.0 ± 2.5	24.3 ± 2.4	22.2 ± 3.7	0.31
Disease duration (years)	9.5 ± 4.6	8.2 ± 6.6	–	0.10
SLEDAI	0	0	–	–

Values were expressed in mean \pm SD; BMI = body mass index; SLEDAI = systematic lupus erythematosus disease activity index.

The lipoprotein profile of CDP, NO THERAPY and CONTROL are shown in Table 2. CDP had lower LDL-c levels compared to NO THERAPY (88 ± 16 versus 108 ± 17 mg/dl; $P < 0.05$) and CONTROL (88 ± 16 versus 118 ± 23 mg/dl; $P < 0.05$). Similarly, TC levels in CDP (156 ± 16 mg/dl) were lower compared to NO THERAPY (174 ± 15 mg/dl; $P < 0.05$) and healthy controls (200 ± 24 mg/dl; $P < 0.05$). In contrast, higher HDL-c levels were found in CDP compared to NO THERAPY (54 ± 8 versus 46 ± 9 mg/dl; $P < 0.05$) although no difference was observed compared to the controls ($P > 0.05$). In addition, NO THERAPY group also had lower HDL-c compared to healthy controls, as well as a lower TC levels ($P < 0.05$). TG and VLDL-c tended to be lower in CDP compared to the other two groups, although this difference did not reach statistical significance ($P = 0.06$ and $P = 0.06$, respectively).

LDE plasma kinetics

Figure 1 shows the plasma decaying curves of the radioactively labeled LDE cholesteryl ester during the 24 h of the study. It should be noticed that the curve obtained in CDP group was pronouncedly faster than NO THERAPY and CONTROL curves (Figure 1). Indeed, as shown in Table 3, the [¹⁴C]cholesteryl ester FCR was significantly different in the three groups studied ($P = 0.03$). In fact, the [¹⁴C]cholesteryl ester FCR in CDP group was greater compared to NO THERAPY ($P < 0.05$) and to CONTROL ($P < 0.05$). In contrast, the [¹⁴C]cholesteryl ester FCR was similar in the comparison of NO THERAPY and CONTROL ($P > 0.05$).

Discussion

This is the first *in vivo* demonstration that in SLE patients treated with CDP the LDE plasma kinetics is accelerated with consequent decrease in LDL cholesterol.

Table 2 Plasma lipid profile of SLE groups (CDP and NO THERAPY) and CONTROL

Parameter	CDP (n = 10)	NO THERAPY (n = 10)	CONTROL (n = 10)	P
Cholesterol				
Total	$156 \pm 16^{a,b}$	174 ± 15^a	200 ± 24	<0.001
HDL-c	54 ± 8^b	46 ± 9^a	64 ± 7	<0.001
LDL-c	$88 \pm 16^{a,b}$	108 ± 17	118 ± 23	<0.001
VLDL-c	13 ± 4	20 ± 7	17 ± 7	0.06
TGs	68 ± 19	103 ± 38	86 ± 33	0.06

Values expressed in mean \pm SD, in mg/dl.

^a $P < 0.05$ versus CONTROL.

^b $P < 0.05$ versus NO THERAPY.




Figure 1 Disappearance curves of the emulsion [^{14}C]cholesteryl ester of CDP, NO THERAPY and CONTROL groups. LDE clearance in CDP patients (continuous line), NO THERAPY patients (dashed line) and CONTROL subjects (dotted line) during 24 hs after intravenous infusion. Disappearance curves of the emulsion [^{14}C]cholesteryl ester of CDP were faster than those of NO THERAPY and CONTROL ($p = 0.03$), indicating a significant increase of clearance in SLE patients taking CDP.

To achieve these findings, rigorous selection criteria was established to eliminate the main systemic diseases that promote specific alterations on lipid metabolism.^{48–52} Additionally, the progressive increase in total and LDL cholesterol with aging⁵³ and menopause⁵⁴ was avoided by the exclusion criteria. All patients included in the study had an inactive form of the disease since it is known that activity can aggravate lupus dyslipoproteinemia¹⁹ and were not using drugs that could interfere in lipoprotein metabolism.⁵⁵ Also importantly, the CDP administration period was long enough to assure that the changes in lipid metabolism by CDP were sustained, which occurs after one year of treatment.^{31,56}

The great advantage of the LDE system used herein is to specifically evaluate the *in vivo* function of LDL receptors that remove the lipoprotein from plasma into cells. LDE is made without protein but when injected into the circulation and in contact with plasma it acquires several small molecular weight apolipoproteins (apo), including apo E.³³ Apo E endows LDE particles to bind to the LDL receptors, since those receptors recognize not only the apo B present in LDL-c, but also apo E that is not found in the LDL fraction.^{35,57} Apo E binds to the LDL receptor with a greater affinity than apo B100 and is thus removed faster than the native LDL.^{35,57} Moreover, the known differences in LDL sub-classes among subjects^{58–60} which may influence the removal of the lipoprotein from the plasma are excluded by the present approach since LDE is a standard preparation allowing a more accurate definition of receptor function.^{37,38}

Table 3 FCR of [^{14}C]cholesteryl ester in SLE groups (CDP and NO THERAPY) and CONTROL

FCR	CDP	NO THERAPY	CONTROL	P
[^{14}C]cholesteryl ester	$0.076 \pm 0.037^{\text{a,b}}$	0.046 ± 0.021	0.051 ± 0.012	0.03

Values in h^{-1} , expressed in mean \pm SD.

^a $p < 0.05$ versus CONTROL.

^b $p < 0.05$ versus NO THERAPY.

Of interest, accelerated LDE removal has been described for acute myelocytic leukemia^{61,62} and a defective function was detected in familial hypercholesterolemia.³⁷ In contrast, we identified herein for the first time that LDL receptor function is normal in SLE supported by the similar LDE removal in lupus patients without therapy and healthy controls. As expected, LDL levels in these two groups were alike in accordance with previous report in inactive untreated lupus patients.¹⁹

Our *in vivo* study also identified that CDP up-regulation of LDL-receptor is a very efficient mechanism for LDL reduction in SLE. A possible explanation for our findings is the known inhibitory effect of chloroquine in lysosomal hydrolysis of cholesterol ester reducing the release of repressive cholesterol with consequent increase in LDL receptor synthesis, as has been demonstrated by *in vitro* experiments in fibroblast and smooth-muscle cells.^{63,64} Reinforcing this hypothesis, the approximately 50% greater LDE removal from the plasma in our CDP was paralleled by a 20% reduction in the LDL-c level compared to untreated patients. Indeed, a decrease in LDL-c levels was also identified in randomized trials after the use of hydroxychloroquine in diabetic patients.^{65,66}

In SLE, it has previously been demonstrated that hydroxychloroquine use elicits lower LDL cholesterol levels, particularly in patients taking corticosteroids.^{9,10} In this regard, the known increase in lipoprotein hepatic synthesis induced by steroids^{10,13,67–69} may be counterbalanced by the anti-malarial enhanced LDL removal. Remarkably, statins (or HMG-CoA) also increase the removal from the plasma of both native LDL and LDE^{37,38} in addition to inhibiting the rate-limiting step of cholesterol synthesis.⁷⁰

Our data provide consistent evidence that CPD powerfully influence the status of LDL receptor function, a known determinant of LDL levels,³² emphasizing its use in lupus patients with high LDL levels, particularly those taking steroids. The possible synergistic effect between CDP and statins should be explored in future studies.

Acknowledgements

Dr. Bonfá and Dr. Maranhão are recipients of Research Awards from Conselho Nacional de Desenvolvimento Científico e Tecnológico (grants 304756/2003-2 and 300771/2003-7, respectively). This research was supported by Fundação do Amparo à Pesquisa do Estado de São Paulo (FAPESP), grant 04/08048-3.

References

- Avina-Zubieta JA, Galindo-Rodriguez G, Newman S *et al.* Long-term effectiveness of antimalarial drugs in rheumatic diseases. *Ann Rheum Dis* 1998; **57**: 582–587.
- Rynes RI. Antimalarial drugs in the treatment of rheumatological diseases. *Br J Rheumatol* 1997; **36**: 799–805.
- Wallace DJ. Antimalarials – the ‘real’ advance in lupus. *Lupus* 2001; **10**(6): 385–387.
- Rothfield N. Efficacy of antimalarials in systemic lupus erythematosus. *Am J Med* 1988; **85**: 53–56.
- Rudnicki RD, Gresham GE, Rothfield NF. The efficacy of antimalarials in systemic lupus erythematosus. *J Rheumatol* 1975; **2**: 323–330.
- Meinao IM, Sato EI, Andrade LE *et al.* Controlled trial with chloroquine diphosphate in systemic lupus erythematosus. *Lupus* 1996; **5**: 237–241.
- The Canadian Hydroxychloroquine Study Group: A randomized study of the effect of withdrawing hydroxychloroquine sulfate in systemic lupus erythematosus. *N Engl J Med* 1991; **324**: 150–154.
- Tsakonas E, Joseph L, Esdaile JM *et al.* A long-term study of hydroxychloroquine withdrawal on exacerbations in systemic lupus erythematosus. The Canadian Hydroxychloroquine Study Group. *Lupus* 1998; **7**: 80–85.
- Wallace DJ, Metzger AL, Stecher VJ *et al.* Cholesterol-lowering effect of hydroxychloroquine in patients with rheumatic disease: reversal of deleterious effects of steroids on lipids. *Am J Med* 1990; **89**: 322–326.
- Petri M, Lakata C, Magder L *et al.* Effect of prednisone and hydroxychloroquine on coronary artery disease risk factors in systemic lupus erythematosus: a longitudinal data analysis. *Am J Med* 1994; **96**: 254–259.
- Petri M. Hydroxychloroquine use in the Baltimore Lupus Cohort: effects on lipids, glucose and thrombosis. *Lupus* 1996; **5** Suppl 1: S16–S22.
- Bruce IN. ‘Not only...but also’: factors that contribute to accelerated atherosclerosis and premature coronary heart disease in systemic lupus erythematosus. *Rheumatology* 2005; **44**: 1492–1502.
- Petri M, Perez-Gutthann S, Spence D *et al.* Risk factors for coronary artery disease in patients with systemic lupus erythematosus. *Am J Med* 1992; **93**: 513–519.
- Bruce IN, Gladman DD, Urowitz MB. Premature atherosclerosis in systemic lupus erythematosus. *Rheum Dis Clin North Am* 2000; **26**: 257–278.
- Manzi S, Meilahn EN, Rairie JE *et al.* Age-specific incidence rates of myocardial infarction and angina in women with systemic lupus erythematosus. *Am J Epidemiol* 1997; **145**: 408–415.
- Ward MM. Premature morbidity from cardiovascular and cerebrovascular diseases in women with systemic lupus erythematosus. *Arthritis Rheum* 1999; **42**: 338–346.
- Ward MM. Outcomes of hospitalizations for myocardial infarctions and cerebrovascular accidents in patients with systemic lupus erythematosus. *Arthritis Rheum* 2004; **50**: 3170–3176.
- Ilowite NT, Samuel P, Ginzler E *et al.* Dyslipoproteinemia in pediatric systemic lupus erythematosus. *Arthritis Rheum* 1988; **31**: 859–863.
- Borba EF, Bonfá E. Dyslipoproteinemias in systemic lupus erythematosus: influence of disease, activity and anticardiolipin antibodies. *Lupus* 1997; **6**: 533–539.
- Borba EF, Bonfá E, Vinagre CGC *et al.* Chylomicron metabolism is markedly altered in systemic lupus erythematosus. *Arthritis Rheum* 2000; **43**: 1033–1040.
- de Carvalho JF, Borba EF, Viana VS *et al.* Anti-lipoprotein lipase antibodies: a new player in the complex atherosclerotic process in systemic lupus erythematosus? *Arthritis Rheum* 2004; **50**: 3610–3615.
- Ettinger WH, Goldberg AP, Applebaum-Bowden D *et al.* Dyslipoproteinemia in systemic lupus erythematosus: effect of corticosteroids. *Am J Med* 1987; **83**: 503–508.
- Ettinger WH Jr, Hazzard WR. Elevated apolipoprotein-B levels in corticosteroid-treated patients with systemic lupus erythematosus. *J Clin Endocrinol Metab* 1988; **67**: 425–428.
- MacGregor AJ, Dhillon VB, Binder A *et al.* Fasting lipids and anticardiolipin antibodies as risk factors for vascular disease in systemic lupus erythematosus. *Ann Rheum Dis* 1992; **51**: 152–155.
- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). *JAMA* 2001; **285**(19): 2486–2497.
- Hodis HN, Quismorio FP, Wickham E *et al.* The lipid, lipoprotein, and apolipoprotein effects of hydroxychloroquine in patients with systemic lupus erythematosus. *J Rheumatol* 1993; **20**: 661–665.
- Kavanaugh A, Adams-Huet B, Jain R *et al.* Hydroxychloroquine effects on lipoprotein profiles (the HELP trial): a double-blind, randomized, placebo-controlled, pilot study in patients with systemic lupus erythematosus. *J Clin Rheumatol* 1997; **3**: 3–8.
- Rahman P, Gladman DD, Urowitz MB *et al.* The cholesterol lowering effect of antimalarial drugs is enhanced in patients with lupus taking corticosteroid drugs. *J Rheumatol* 1999; **26**: 325–330.
- Tam LS, Li EK, Lam CW *et al.* Hydroxychloroquine has no significant effect on lipids and apolipoproteins in Chinese systemic lupus erythematosus patients with mild or inactive disease. *Lupus* 2000; **9**: 413–416.
- Tam LS, Gladman DD, Hallett D *et al.* Effect of antimalarial agents on the fasting lipid profile in systemic lupus erythematosus. *J Rheumatol* 2000; **27**: 2142–2145.
- Borba EF, Bonfá E. Longterm beneficial effect of chloroquine diphosphate on lipoprotein profile in lupus patients with and without steroid therapy. *J Rheumatol* 2001; **28**: 780–785.
- Brown MS, Goldstein JL. A receptor-mediated pathway for cholesterol homeostasis. *Science* 1986; **232**: 34–47.
- Maranhão RC, Cesar TB, Pedroso-Mariani SR *et al.* Metabolic behavior in rats of a nonprotein microemulsion resembling low density lipoprotein. *Lipids* 1993; **28**: 691–696.
- Maranhão RC, Roland IA, Toffoletto O *et al.* Plasma kinetic behavior in hyperlipidemic subjects of a lipidic microemulsion that binds to LDL receptors. *Lipids* 1997; **32**: 627–632.
- Hirata RDC, Hirata MH, Mesquita CH *et al.* Effects of apolipoprotein B-100 on the metabolism of a lipid microemulsion model in rats. *Biochim Biophys Acta* 1999; **1437**: 53–62.
- Pinto LB, Wajgarten M, Silva EL *et al.* Plasma kinetics of a cholesterol-rich emulsion in young, middle-aged and elderly subjects. *Lipids* 2001; **36**: 1307–1311.
- Santos RD, Chacra AP, Morikawa A *et al.* Plasma kinetics of free and esterified cholesterol in familial hypercholesterolemia: effects of simvastatin. *Lipids* 2005; **40**: 737–743.
- Santos RD, Hueb W, Oliveira AA *et al.* Plasma kinetics of a cholesterol-rich emulsion in subjects with or without coronary artery disease. *J Lipid Res* 2003; **44**: 464–469.
- Tan EM, Cohen AS, Fries JF *et al.* The 1982 revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1982; **25**: 1271–1277.
- Bombardier C, Gladman DD, Urowitz MB *et al.*, and the committee on prognosis studies in SLE. Derivation of the SLEDAI. A disease activity index for lupus patients. *Arthritis Rheum* 1992; **35**: 630–640.
- Ginsburg GS, Small DM, Atkinson D. Microemulsion of phospholipids and cholesterol esters: protein-free models of low density lipoprotein. *J Biol Chem* 1982; **257**: 8216–8227.
- Marchese SRM, Mesquita CH, Cunha ILL. Anacomp program application to calculate ¹³⁷C transfer rates in marine organisms and dose in man. *J Radioanal Nucl Chem* 1998; **232**: 233–236.
- Maranhão RC, Feres MC, Martins MT *et al.* Plasma kinetics of a chylomicron-like emulsion in patients with coronary artery disease. *Atherosclerosis* 1996; **126**: 15–25.
- Siedel J, Hägele EO, Ziegenhorn J *et al.* Reagent for the enzymatic determination of serum total cholesterol with improved lipolytic efficiency. *Clin Chem* 1983; **29**: 1075–1080.

- 45 Fossati P, Prencipe L. Serum triglycerides determined colorimetrically with an enzyme that produces hydrogen peroxide. *Clin Chem* 1982; **28**: 2077–2080.
- 46 Warnick GR, Cheung MC, Albers JJ. Comparison of current methods for high-density lipoprotein cholesterol quantification. *Clin Chem* 1979; **25**: 596–604.
- 47 Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972; **18**: 499–502.
- 48 Brunzell JD, Albers JJ, Haas LB *et al.* Prevalence of serum lipid abnormalities in chronic hemodialysis. *Metabolism* 1977; **26**: 903–910.
- 49 Attman PO, Alaupovic P. Lipid and apolipoprotein profiles of uremic dyslipoproteinemia. Relation to renal function and dialysis. *Nephron* 1991; **57**: 401–410.
- 50 Appel GB, Blum CB, Chien S *et al.* The hyperlipidemia of the nephrotic syndrome. Relation to plasma albumin concentration, oncotic pressure and viscosity. *N Engl J Med* 1985; **312**: 1544–1548.
- 51 Thompson GR, Soutar AK, Spengel FA *et al.* Defects of receptor-mediated low density lipoprotein catabolism in homozygous familial hypercholesterolemia and hypothyroidism. *Proc Soc Natl Acad Sci (USA)* 1981; **78**: 2591–2595.
- 52 Brown WV. Lipoprotein disorders in diabetes mellitus. *Med Clin North Am* 1994; **78**: 143–161.
- 53 Heiss G, Tamir I, Davis CE *et al.* Lipoprotein-cholesterol distributions in selected North American populations: the Lipid Research Clinics Program Prevalence Study. *Circulation* 1980; **61**: 302–315.
- 54 Kannel WB, Hjortland MC, McNamara PM *et al.* Menopause and risk of cardiovascular disease. The Framingham Study. *Ann Intern Med* 1976; **85**: 447–452.
- 55 Henkin Y, Como JA, Oberman A. Secondary dyslipidemia. Inadvertent effects of drugs in clinical practice. *JAMA* 1992; **267**: 961–968.
- 56 Munro R, Morrison E, McDonald AG *et al.* Effect of disease modifying agents on the lipid profiles of patients with rheumatoid arthritis. *Ann Rheum Dis* 1997; **56**: 374–377.
- 57 Innerarity TL, Mahley RW. Enhanced binding by cultured human fibroblasts of apo-E-containing lipoproteins as compared with low density lipoproteins. *Biochemistry* 1978; **17**: 1440–1447.
- 58 Capell WH, Zambon A, Austin MA *et al.* Compositional differences of LDL particles in normal subjects with LDL subclass phenotype A and LDL subclass phenotype B. *Arterioscler Thromb Vasc Biol* 1996; **16**: 1040–1046.
- 59 Ohmura H, Mokuno H, Sawano M *et al.* Lipid compositional differences of small, dense low-density lipoprotein particle influence its oxidative susceptibility: possible implication of increased risk of coronary artery disease in subjects with phenotype B. *Metabolism* 2002; **51**: 1081–1087.
- 60 Chapman MJ, Guerin M, Bruckert E. Atherogenic, dense low-density lipoproteins. Pathophysiology and new therapeutic approaches. *Eur Heart J* 1998; **19** Suppl A: A24–A30.
- 61 Maranhao RC, Garicochea B, Silva EL *et al.* Increased plasma removal of microemulsions resembling the lipid phase of low-density lipoproteins (LDL) in patients with acute myeloid leukemia. *Braz J Med Biol Res* 1992; **25**: 1003–1007.
- 62 Maranhao RC, Garicochea B, Silva EL *et al.* Plasma kinetics and biodistribution of a lipid emulsion resembling low-density lipoprotein in patients with acute leukemia. *Cancer Res* 1994; **54**: 4660–4666.
- 63 Middleton B, Middleton A. Cyclic AMP stimulates the synthesis and function of the low-density lipoprotein receptor in human vascular smooth-muscle cells and fibroblasts. *Biochem J* 1992; **282**(3): 853–861.
- 64 Middleton A, Middleton B. Agents which increase cyclic AMP have diverse effects on low-density-lipoprotein-receptor function in human vascular smooth-muscle cells and skin fibroblasts. *Biochem J* 1990; **267**(3): 607–614.
- 65 Powrie JK, Shojaae-Moradie F, Watts GF *et al.* Effects of chloroquine on the dyslipidemia of non-insulin-dependent diabetes mellitus. *Metabolism* 1993; **42**: 415–419.
- 66 Gerstein HC, Thorpe KE, Taylor DW *et al.* The effectiveness of hydroxychloroquine in patients with type 2 diabetes mellitus who are refractory to sulfonyleureas – a randomized trial. *Diabetes Res Clin Pract* 2002; **55**: 209–219.
- 67 Petri M, Spence D, Bone LR *et al.* Coronary artery disease risk factors in the Johns Hopkins Lupus Cohort: prevalence, recognition by patients, and preventive practices. *Medicine* 1992; **71**: 291–302.
- 68 Bruce IN, Urowitz MB, Gladman DD *et al.* Natural history of hypercholesterolemia in systemic lupus erythematosus. *J Rheumatol* 1999; **26**: 2137–2143.
- 69 Svenungsson E, Jensen-Ustad K, Heimbürger M *et al.* Risk factors for cardiovascular disease in systemic lupus erythematosus. *Circulation* 2001; **104**: 1887–1893.
- 70 Abeles AM, Pillinger MH. Statins as antiinflammatory and immunomodulatory agents: a future in rheumatologic therapy? *Arthritis Rheum* 2006; **54**: 393–407.