

```


In [4]: import numpy as np
import matplotlib.pyplot as plt
# specify size of region:
Lx=10.0
Ly=10.0
# introduce two point charges:
x1=2; y1=3; z1= 3; q1= 2 #position of charge #1
x2=3; y2=2; z2= -1; q2=-2 #position of charge #2
x3=6; y3=8; z3= 2; q3= 0.5

# create a 2D array in NumPy for the potential with N elements in each
# direction and set its elements initially to zero
Nx = 101 #number of points on our x-axis.
Ny = 1010 #number of points on our y-axis.
phia = np.zeros((Ny,Nx))
phi2 = np.zeros((Ny,Nx))
phi3 = np.zeros((Ny,Nx))
X = np.zeros(Nx)
Y = np.zeros(Ny)
Dx=Lx/(Nx-1); Dy=Ly/(Ny-1) # find spacing between the points in x and y
z=1 # choose to work at fixed z=1
for nx in range (0,Nx):
    X[nx]= Dx*nx
    for ny in range (0,Ny):
        Y[ny]= Dy*ny
        phia[ny,nx]= q1/((Dx*nx-x1)**2+(Dy*ny-y1)**2+(z-z1)**2)**0.5
        phi2[ny,nx]= phia[ny,nx] + q2/((Dx*nx-x2)**2+(Dy*ny-y2)**2+(z-z2)**2)
        phi3[ny,nx]= phi2[ny,nx] + q3/((Dx*nx-x3)**2+(Dy*ny-y3)**2+(z-z3)**2)

Nl = 30 # the number of contour levels to be plotted.
clevel = np.zeros(Nl)
for n in range(0,Nl):
    clevel[n]=0.02*n; # select the values of phi for which
                      # equipotential lines will be plotted

fig = plt.figure(figsize=(10,10))
ax = fig.add_subplot(1,1,1)
ax.contour(X,Y,phi3,levels=clevel, colors='b', linewidths=1)
ax.contour(X,Y,-phi3,levels=clevel, colors='b', linewidths=1, linestyle='d')
plt.show()

```



In []:

In []: