Lecture 1 Practice Problems

Michelle Wachs

(1) Give the cycle type of each of the following permutations.

- (a) [397468152]
- (b) [234561]
- (c) [1234]

(2) For $\lambda \vdash n$, let $z_{\lambda} = 1^{m_1} m_1 ! 2^{m_2} m_2 ! \cdots$, where m_i is the number of occurrences of i in λ . Show that the number of permutations of cycle type λ is $\frac{n!}{z_{\lambda}}$.

(3) In this problem we show that the various characterizations of the Eulerian polynomial $A_n(t)$ are consistent with each other and with Euler's original definition:

$$\sum_{i>1} i^n t^i = \frac{t A_n(t)}{(1-t)^{n+1}}.$$

(a) Using Euler's original definition of $A_n(t)$, prove his exponential generating function formula

$$\sum_{n>0} A_n(t) \frac{z^n}{n!} = \frac{1-t}{e^{(t-1)z} - t}.$$

(b) Using Euler's original definition of $A_n(t)$, prove that the coefficients of $A_n(t)$ satisfy the recurrence

$$\left\langle {n\atop j}\right\rangle = (n-j)\left\langle {n-1\atop j-1}\right\rangle + (j+1)\left\langle {n-1\atop j}\right\rangle$$

(c) Using the combinatorial characterization of $A_n(t)$ involving des, prove that the coefficients of $A_n(t)$ satisfy the above recurrence.

(d) Using the combinatorial characterization of $A_n(t)$ involving exc, prove that the coefficients of $A_n(t)$ satisfy the above recurrence.

(4) Give a bijective proof that des and exc are equidistributed on \mathfrak{S}_n .

(5) Let C_n be the set of permutations in \mathfrak{S}_n of cycle type (n). Find a formula for

$$\sum_{\sigma \in C_n} t^{\operatorname{exc}(\sigma)}.$$

(6) A binary tree is a rooted tree in which each node has at most two children and each child is either a left child or a right child. An increasing binary tree is a binary tree with n nodes labeled with $1, 2, \ldots, n$ so that every node has a label that is bigger than its parent. Let \mathcal{T}_n be the set of increasing binary trees with n nodes. For $T \in \mathcal{T}_n$, let l(T) be the number of left children in T. Prove

(a)
$$\sum_{T \in \mathcal{T}_n} t^{l(T)} = A_n(t).$$

(b)
$$A_n(t) = \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} \gamma_{n,k} t^k (1+t)^{n-1-2k},$$

where $\gamma_{n,k} = |\{\sigma \in \mathfrak{S}_n : \sigma \text{ has no double descents, no final descent, and } \operatorname{des}(\sigma) = k\}|.$

- (7) Show that γ -positivity implies palindromicity and unimodality, but the converse is false.
- (8) Prove:

(a)

$$\sum_{\sigma \in \mathfrak{S}_n} q^{\mathrm{inv}(\sigma)} = [n]_q!.$$

(b) Let $\mathfrak{S}_{k_1,k_2,...,k_m}$ be the set of permutations of the multiset $\{1^{k_1},2^{k_2},\ldots,m^{k_m}\}$. In other words $\mathfrak{S}_{k_1,k_2,...,k_m}$ is the set of words with k_i i's for each i. Prove that

$$\sum_{\sigma \in \mathfrak{S}_{k_1, k_2, \dots, k_m}} q^{\mathrm{inv}(\sigma)} = \left[\begin{array}{c} n \\ k_1, k_2, \dots, k_m \end{array} \right]_q.$$

(9) Let

$$A_n(q,t) := \sum_{\sigma \in \mathfrak{S}} q^{\operatorname{maj}(\sigma) - \operatorname{exc}(\sigma)} t^{\operatorname{exc}(\sigma)}$$

and let Alt_n be the set of permutations σ in \mathfrak{S}_n such that

$$\sigma(1) > \sigma(2) < \sigma(3) > \sigma(4) < \sigma(5) > \dots < \sigma(n).$$

Prove that

$$A_n(q,-1) = \begin{cases} (-1)^{\frac{n-1}{2}} \sum_{\sigma \in Alt_n} q^{\operatorname{inv}(\sigma)} & \text{if } n \text{ is odd} \\ 0 & \text{if } n \text{ is even} \end{cases}.$$

(10) Prove the closed form formula

$$A_n(q,t) = \sum_{m=1}^{\lfloor \frac{n+1}{2} \rfloor} \sum_{k_1,\dots,k_m \ge 2} \begin{bmatrix} n \\ k_1 - 1, k_2, \dots, k_m \end{bmatrix}_q t^{m-1} \prod_{i=1}^m [k_i - 1]_t$$