


# Some families of reduced Kronecker coefficients, plane partitions and quasipolynomials

Laura Colmenarejo

Universidad de Sevilla

**ECCO 2016**

**June 20, 2016**



# CONTENTS

Framework

Reduced Kronecker coefficients

Generating functions

Plane partitions

Quasipolynomials

# CONTENTS

Framework

Reduced Kronecker coefficients

Generating functions

Plane partitions

Quasipolynomials

## SYMMETRIC FUNCTIONS

Let  $X$  be an (infinite) alphabet and  $\lambda \vdash n$ . The **Schur function** is defined as

$$s_\lambda = \sum_{T \in SSYT} x^T$$

## SYMMETRIC FUNCTIONS

Let  $X$  be an (infinite) alphabet and  $\lambda \vdash n$ . The **Schur function** is defined as

$$s_\lambda = \sum_{T \in SSYT} x^T$$

We define the **ordinary product**

$$\begin{aligned} Sym \otimes Sym &\longrightarrow Sym \\ (f, g) &\longrightarrow f \cdot g \end{aligned}$$

## SYMMETRIC FUNCTIONS

Let  $X$  be an (infinite) alphabet and  $\lambda \vdash n$ . The **Schur function** is define as

$$s_\lambda = \sum_{T \in SSYT} x^T$$

We define the **ordinary product**

$$\begin{aligned} Sym \otimes Sym &\longrightarrow Sym \\ (f, g) &\longrightarrow f \cdot g \end{aligned}$$

Consider the following case

$$s_\lambda s_\mu = \sum_{\nu} c_{\lambda\mu}^\nu s_\nu$$

## SYMMETRIC FUNCTIONS

Let  $X$  be an (infinite) alphabet and  $\lambda \vdash n$ . The **Schur function** is defined as

$$s_\lambda = \sum_{T \in \text{SSYT}} x^T$$

We define the **ordinary product**

$$\begin{aligned} \text{Sym} \otimes \text{Sym} &\longrightarrow \text{Sym} \\ (f, g) &\longrightarrow f \cdot g \end{aligned}$$

Consider the following case

$$s_\lambda s_\mu = \sum_{\nu} c_{\lambda\mu}^\nu s_\nu$$

Littlewood-Richardson coefficients:  $c_{\lambda\mu}^\nu$

# LITTLEWOOD-RICHARDSON COEFFICIENTS

**Littlewood-Richardson rule** The Littlewood-Richardson coefficient  $c_{\lambda\mu}^{\nu}$  counts the number of Littlewood-Richardson tableaux of shape  $\nu/\lambda$  and weight  $\mu$ .



# LITTLEWOOD-RICHARDSON COEFFICIENTS

**Littlewood-Richardson rule** The Littlewood-Richardson coefficient  $c_{\lambda/\mu}^{\nu}$  counts the number of Littlewood-Richardson tableaux of shape  $\nu/\lambda$  and weight  $\mu$ .

Example:

$$s_{21} \cdot s_{21} = s_{42} + s_{411} + s_{33} + 2s_{321} + s_{3111} + s_{222} + s_{2211}$$

# LITTLEWOOD-RICHARDSON COEFFICIENTS

**Littlewood-Richardson rule** The Littlewood-Richardson coefficient  $c_{\lambda/\mu}^{\nu}$  counts the number of Littlewood-Richardson tableaux of shape  $\nu/\lambda$  and weight  $\mu$ .

Example:

$$s_{21} \cdot s_{21} = s_{42} + s_{411} + s_{33} + 2s_{321} + s_{3111} + s_{222} + s_{2211}$$

- The Littlewood-Richardson rule was stated in 1934.

# LITTLEWOOD-RICHARDSON COEFFICIENTS

**Littlewood-Richardson rule** The Littlewood-Richardson coefficient  $c_{\lambda/\mu}^{\nu}$  counts the number of Littlewood-Richardson tableaux of shape  $\nu/\lambda$  and weight  $\mu$ .

Example:

$$s_{21} \cdot s_{21} = s_{42} + s_{411} + s_{33} + 2s_{321} + s_{3111} + s_{222} + s_{2211}$$

- ▶ The Littlewood-Richardson rule was stated in 1934.
- ▶ The first rigorous proofs of the rule were given by Schützenberger (1977) and Thomas (1974). They use the Robinson-Schensted correspondence.

# LITTLEWOOD-RICHARDSON COEFFICIENTS

**Littlewood-Richardson rule** The Littlewood-Richardson coefficient  $c_{\lambda/\mu}^{\nu}$  counts the number of Littlewood-Richardson tableaux of shape  $\nu/\lambda$  and weight  $\mu$ .

Example:

$$s_{21} \cdot s_{21} = s_{42} + s_{411} + s_{33} + 2s_{321} + s_{3111} + s_{222} + s_{2211}$$

- ▶ The Littlewood-Richardson rule was stated in 1934.
- ▶ The first rigorous proofs of the rule were given by Schützenberger (1977) and Thomas (1974). They use the Robinson-Schensted correspondence.
- ▶ Description in terms of *integral hives* and *honeycombs*, by T. Tao and A. Knutson (1999).

# KRONECKER COEFFICIENTS

# KRONECKER COEFFICIENTS

Let  $X$  and  $Y$  be two alphabets.

- ▶ Sum:  $X + Y = x_1 + y_1 + x_2 + y_2 + \dots$
- ▶ Product:  $XY = x_1y_1 + x_1y_2 + \dots + x_2y_1 + x_2y_2 + \dots$

## KRONECKER COEFFICIENTS

Let  $X$  and  $Y$  be two alphabets.

- ▶ Sum:  $X + Y = x_1 + y_1 + x_2 + y_2 + \dots$
- ▶ Product:  $XY = x_1y_1 + x_1y_2 + \dots + x_2y_1 + x_2y_2 + \dots$

The Littlewood-Richardson coefficients can be obtained as

$$s_\nu[X + Y] = \sum_{\lambda, \mu} c_{\lambda\mu}^\nu s_\lambda[X] \cdot s_\mu[Y]$$

where we sum over all the partitions such that  $|\lambda| + |\mu| = |\nu|$ .

# KRONECKER COEFFICIENTS

Let  $X$  and  $Y$  be two alphabets.

- ▶ Sum:  $X + Y = x_1 + y_1 + x_2 + y_2 + \dots$
- ▶ Product:  $XY = x_1y_1 + x_1y_2 + \dots + x_2y_1 + x_2y_2 + \dots$

The Littlewood-Richardson coefficients can be obtained as

$$s_\nu[X + Y] = \sum_{\lambda, \mu} c_{\lambda\mu}^\nu s_\lambda[X] \cdot s_\mu[Y]$$

where we sum over all the partitions such that  $|\lambda| + |\mu| = |\nu|$ .

$$s_\nu[XY] = \sum_{\lambda, \mu} g_{\lambda\mu}^\nu s_\lambda[X] s_\mu[Y]$$

where we sum over all the partitions such that  $|\lambda| = |\mu| = |\nu|$ .



# KRONECKER COEFFICIENTS

Let  $X$  and  $Y$  be two alphabets.

- ▶ Sum:  $X + Y = x_1 + y_1 + x_2 + y_2 + \dots$
- ▶ Product:  $XY = x_1y_1 + x_1y_2 + \dots + x_2y_1 + x_2y_2 + \dots$

The Littlewood-Richardson coefficients can be obtained as

$$s_\nu[X + Y] = \sum_{\lambda, \mu} c_{\lambda\mu}^\nu s_\lambda[X] \cdot s_\mu[Y]$$

where we sum over all the partitions such that  $|\lambda| + |\mu| = |\nu|$ .

$$s_\nu[XY] = \sum_{\lambda, \mu} g_{\lambda\mu}^\nu s_\lambda[X] s_\mu[Y]$$

where we sum over all the partitions such that  $|\lambda| = |\mu| = |\nu|$ .

Kronecker coefficients:  $g_{\lambda\mu}^\nu$

# KRONECKER COEFFICIENTS

- There is **NO** description in general.

# KRONECKER COEFFICIENTS

- ▶ There is **NO** description in general.
- ▶ Case of two rows partitions: Thibon, R. Orellana, E. Briand and M. H. Rosas.

# KRONECKER COEFFICIENTS

- ▶ There is **NO** description in general.
- ▶ Case of two rows partitions: Thibon, R. Orellana, E. Briand and M. H. Rosas.
- ▶ Case of hooks: Blasiak, Remmel.

## KRONECKER COEFFICIENTS

- ▶ There is **NO** description in general.
- ▶ Case of two rows partitions: Thibon, R. Orellana, E. Briand and M. H. Rosas.
- ▶ Case of hooks: Blasiak, Remmel.
- ▶ Case of one partition with two parts: R. Orellana and C. Ballantine.

## KRONECKER COEFFICIENTS

- ▶ There is **NO** description in general.
- ▶ Case of two rows partitions: Thibon, R. Orellana, E. Briand and M. H. Rosas.
- ▶ Case of hooks: Blasiak, Remmel.
- ▶ Case of one partition with two parts: R. Orellana and C. Ballantine.
- ▶ Stability properties and complexity: G. Panova and I. Pak.

# KRONECKER COEFFICIENTS

- ▶ There is **NO** description in general.
- ▶ Case of two rows partitions: Thibon, R. Orellana, E. Briand and M. H. Rosas.
- ▶ Case of hooks: Blasiak, Remmel.
- ▶ Case of one partition with two parts: R. Orellana and C. Ballantine.
- ▶ Stability properties and complexity: G. Panova and I. Pak.

The Kronecker coefficients define a *new* product on  $Sym$

$$\begin{aligned} Sym \otimes Sym &\longrightarrow Sym \\ (s_\lambda, s_\mu) &\longrightarrow s_\lambda * s_\mu \end{aligned}$$

where

$$s_\lambda * s_\mu := \sum_{\nu} g_{\lambda\mu}^{\nu} s_{\nu}$$

# CONTENTS

Framework

Reduced Kronecker coefficients

Generating functions

Plane partitions

Quasipolynomials



Reduced Kronecker coefficients:  $\overline{g}_{\lambda\mu}^\nu$

$$\begin{aligned}
 s_{(10,2)} * s_{(10,2)} &= s_{(12)} + s_{(11,1)} + 2 \cdot s_{(10,2)} + s_{(10,1,1)} + \\
 &+ s_{(9,3)} + 2 \cdot s_{(9,2,1)} + s_{(9,1,1,1)} + \\
 &+ s_{(8,4)} + s_{(8,3,1)} + s_{(8,2,2)}
 \end{aligned}$$

# Reduced Kronecker coefficients: $\bar{g}_{\lambda\mu}^\nu$

$$\begin{aligned}
 s_{(10,2)} * s_{(10,2)} &= s_{(12)} + s_{(11,1)} + 2 \cdot s_{(10,2)} + s_{(10,1,1)} + \\
 &+ s_{(9,3)} + 2 \cdot s_{(9,2,1)} + s_{(9,1,1,1)} + \\
 &+ s_{(8,4)} + s_{(8,3,1)} + s_{(8,2,2)}
 \end{aligned}$$

$$\begin{aligned}
 s_{(\bullet,2)} * s_{(\bullet,2)} &= s_{(\bullet)} + s_{(\bullet,1)} + 2 \cdot s_{(\bullet,2)} + s_{(\bullet,1,1)} + \\
 &+ s_{(\bullet,3)} + 2 \cdot s_{(\bullet,2,1)} + s_{(\bullet,1,1,1)} + \\
 &+ s_{(\bullet,4)} + s_{(\bullet,3,1)} + s_{(\bullet,2,2)}
 \end{aligned}$$

Reduced Kronecker coefficients:  $\bar{g}_{\lambda\mu}^\nu$

$$\begin{aligned} s_{(\bullet,2)} * s_{(\bullet,2)} &= s_{(\bullet)} + s_{(\bullet,1)} + 2 \cdot s_{(\bullet,2)} + s_{(\bullet,1,1)} + \\ &+ s_{(\bullet,3)} + 2 \cdot s_{(\bullet,2,1)} + s_{(\bullet,1,1,1)} + \\ &+ s_{(\bullet,4)} + s_{(\bullet,3,1)} + s_{(\bullet,2,2)} \end{aligned}$$

The **reduced Kronecker coefficients**  $\bar{g}_{\lambda\mu}^\nu$  are the stable values of the sequence of Kronecker coefficients, after we disregard the first part.

# RELATION WITH KRONECKER COEFFICIENTS

## RELATION WITH KRONECKER COEFFICIENTS

- Kronecker coefficients can be recovered from reduced Kronecker coefficients.

$$g_{\lambda\mu}^{\nu} = \sum_{i=1}^{\ell(\lambda)\ell(\mu)} (-1)^{i+1} \bar{g}_{\bar{\lambda}\bar{\mu}}^{\nu^{\dagger i}}$$

## RELATION WITH KRONECKER COEFFICIENTS

- Kronecker coefficients can be recovered from reduced Kronecker coefficients.

$$g_{\lambda\mu}^{\nu} = \sum_{i=1}^{\ell(\lambda)\ell(\mu)} (-1)^{i+1} \bar{g}_{\bar{\lambda}\bar{\mu}}^{\nu^{\dagger i}}$$

- Reduced Kronecker coefficients are Kronecker coefficients.

# RELATION WITH KRONECKER COEFFICIENTS

- Kronecker coefficients can be recovered from reduced Kronecker coefficients.

$$g_{\lambda\mu}^{\nu} = \sum_{i=1}^{\ell(\lambda)\ell(\mu)} (-1)^{i+1} \bar{g}_{\bar{\lambda}\bar{\mu}}^{\nu^{\dagger i}}$$

- Reduced Kronecker coefficients are Kronecker coefficients.

Proposition (E. Briand - R. Orellana - M. Rosas, 2011)

$$\bar{g}_{\lambda\mu}^{\nu} = g_{\lambda[n]\mu[n]}^{\nu[n]}$$

with  $\lambda[n] = (n - |\lambda|, \lambda_1, \lambda_2, \dots)$  and  $n$  sufficiently large.

# FAMILIES

- ▶ **Family 1**  $\bar{g}_{(k^a), (k^a)}^{(k)}$
- ▶ **Family 2**  $\bar{g}_{((k+i)^a), (k^a)}^{(k)}$
- ▶ **Family 3**  $\bar{g}_{(k^{a+1}), (k+i, k^a)}^{(k)}$



# CONTENTS

Framework

Reduced Kronecker coefficients

Generating functions

Plane partitions

Quasipolynomials

# FAMILY 1

## Theorem (L. C. - M. Rosas, 2015)

*Consider the reduced Kronecker coefficients  $\left\{ \bar{g}_{(k^a), (k^a)}^{(k)} \right\}_{k \geq 0}$ .*

*Their generating functions is*

# FAMILY 1

## Theorem (L. C. - M. Rosas, 2015)

Consider the reduced Kronecker coefficients  $\left\{ \bar{g}_{(k^a), (k^a)}^{(k)} \right\}_{k \geq 0}$ .

Their generating functions is

$$\mathcal{F}_a = \frac{1}{(1-x)(1-x^2)^2 \cdots (1-x^a)^2(1-x^{a+1})}$$

# FAMILY 1

## Theorem (L. C. - M. Rosas, 2015)


Consider the reduced Kronecker coefficients  $\left\{ \bar{g}_{(k^a), (k^a)}^{(k)} \right\}_{k \geq 0}$ .

Their generating functions is


$$\mathcal{F}_a = \frac{1}{(1-x)(1-x^2)^2 \cdots (1-x^a)^2(1-x^{a+1})}$$

# KRONECKER TABLEAU

# KRONECKER TABLEAU



# KRONECKER TABLEAU




## ► $\alpha$ -condition:

- If  $\alpha_1 = \alpha_2$ , there is no condition.
- If  $\alpha_1 > \alpha_2$ , then

$$(\#1)_{2nd \text{ Row}} = \alpha_1 - \alpha_2$$

$$(\#2)_{1st \text{ Row}} = \alpha_1 - \alpha_2$$

# KRONECKER TABLEAU



►  $\alpha$ -condition:

- If  $\alpha_1 = \alpha_2$ , there is no condition.
- If  $\alpha_1 > \alpha_2$ , then


$$(\#1)_{2nd \text{ Row}} = \alpha_1 - \alpha_2$$

$$(\#2)_{1st \text{ Row}} = \alpha_1 - \alpha_2$$

- The reverse reading word is an  $\alpha$ -lattice permutation.



# KRONECKER TABLEAU



EXAMPLE FOR  $a = 4$ ,  $k = 7$ ,  $\alpha = (3, 2, 1, 1)$ ,  
 $sh = (21, 7, 7, 7, 7)/\alpha$ ,  $ty = (18, 5, 6, 6, 7)$

							1	1	1	1	1	1	1	1	1	1	1	1	2	3	5
							1	2	2	2	2										
							1	3	3	3	3	3									
							4	4	4	4	4	4									
							1	5	5	5	5	5	5								

# SKETCH OF THE PROOF FOR CASE $a = b$

Using a result of R. Orellana and C. Ballantine,

$$\overline{g}_{(k^a)(k^a)}^{(k)} = \# \left\{ \begin{array}{l} \text{Kronecker tableau with} \\ sh = (3k, k^a)/\alpha \\ ty = (3k, k^a)/\alpha \\ \alpha \vdash k \end{array} \right\}$$

# SKETCH OF THE PROOF FOR CASE $a = b$

Using a result of R. Orellana and C. Ballantine,

$$\overline{g}_{(k^a)(k^a)}^{(k)} = \# \left\{ \begin{array}{l} \text{Kronecker tableau with} \\ sh = (3k, k^a)/\alpha \\ ty = (3k, k^a)/\alpha \\ \alpha \vdash k \end{array} \right\}$$

Looking at the generating function  $\mathcal{F}_a$ ,

$$\mathcal{F}_a = \frac{1}{(1-x)(1-x^2)^2 \cdots (1-x^a)^2(1-x^{a+1})}$$

$$\text{Coefficient of } x^k \text{ in } \mathcal{F}_a = \# \left\{ \begin{array}{l} \text{colored sequences of } k \\ \text{with parts in} \\ \{1, 2, \overline{2}, \dots, a, \overline{a}, a+1\} \end{array} \right\}$$

# SKETCH OF THE PROOF FOR CASE $a = b$

There exists a bijection


$$\left\{ \begin{array}{l} \text{colored sequences of } k \\ \text{with parts in} \\ \{1, 2, \bar{2}, \dots, a, \bar{a}, a+1\} \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{Kronecker tableau} \\ sh = (3k, k^a)/\alpha \\ ty = (3k, k^a)/\alpha \\ \alpha \vdash k \end{array} \right\}$$

# SKETCH OF THE PROOF FOR CASE $a = b$

There exists a bijection

$$\left\{ \begin{array}{l} \text{colored sequences of } k \\ \text{with parts in} \\ \{1, 2, \bar{2}, \dots, a, \bar{a}, a+1\} \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{Kronecker tableau} \\ sh = (3k, k^a)/\alpha \\ ty = (3k, k^a)/\alpha \\ \alpha \vdash k \end{array} \right\}$$

It is based on the following identification



EXAMPLE:  $a = 4$  AND  $\beta = (1, \bar{2}, \bar{4})$

EXAMPLE:  $a = 4$  AND  $\beta = (1, \bar{2}, \bar{4})$

STEP 0. We know that  $\lambda = \nu = (21, 7, 7, 7, 7)$ .

**EXAMPLE:**  $a = 4$  AND  $\beta = (1, \bar{2}, \bar{4})$

STEP 0. We know that  $\lambda = \nu = (21, 7, 7, 7, 7)$ .

STEP 1. Column identification

$\bar{4}$	$\bar{2}$	$1$
$\downarrow$	$\downarrow$	$\downarrow$
		1
	1	3
	4	4
1	5	5

STEP 2. Identify  $\alpha$ :

Looking at the picture, we have that  $\alpha = (3, 2, 1, 1)$ , which is a partition of 7.



EXAMPLE:  $a = 4$  AND  $\beta = (1, \bar{2}, \bar{4})$

STEP 0. We know that  $\lambda = \nu = (21, 7, 7, 7, 7)$ .

### STEP 1. Column identification

STEP 2. Identify  $\alpha$ :

Looking at the picture, we have that  $\alpha = (3, 2, 1, 1)$ , which is a partition of 7.

		1
	1	3
	4	4
1	5	5

STEP 3. Complete rest of the tableau

		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	3	5
		1	2	2	2	2													
		1	3	3	3	3	3												
		4	4	4	4	4	4												
1	5	5	5	5	5	5													

Family 2:  $\bar{g}_{((k+i)^a), (k^a)}^{(k)}$

Family 2:  $\bar{g}_{((k+i)^a), (k^a)}^{(k)}$

Table: Case  $a = 2$

k	0	1	2	3	4	5	6	7	8	9	10	11	12	13
i= 0	1	1	3	4	7	9	14	17	24	29	38	45	57	66
i= 1	0	0	0	1	1	3	4	7	9	14	17	24	29	38
i= 2	0	0	0	0	0	0	1	1	3	4	7	9	14	17
i= 3	0	0	0	0	0	0	0	0	0	1	1	3	4	7

Family 2:  $\bar{g}_{((k+i)^a), (k^a)}^{(k)}$

Table: Case  $a = 2$

k	0	1	2	3	4	5	6	7	8	9	10	11	12	13
i= 0	1	1	3	4	7	9	14	17	24	29	38	45	57	66
i= 1	0	0	0	1	1	3	4	7	9	14	17	24	29	38
i= 2	0	0	0	0	0	0	1	1	3	4	7	9	14	17
i= 3	0	0	0	0	0	0	0	0	0	1	1	3	4	7

### Theorem (L. C., 2015)

After some initial zeros, the generating function for the reduced Kronecker coefficients  $\bar{g}_{((k+i)^a), (k^a)}^{(k)}$  is exactly  $\mathcal{F}_a$ .

Family 3:  $\overline{g}^{(k)}_{((k+i), k^{a-1}), (k^a)}$

Family 3:  $\bar{g}_{((k+i), k^{a-1}), (k^a)}^{(k)}$

Table: Case  $a = 2$ .

k	0	1	2	3	4	5	6	7	8	9	10	11	12
i= 0	<b>1</b>	1	3	4	7	9	14	17	24	29	38	45	57
i= 1	0	<b>1</b>	<b>2</b>	4	7	11	16	23	31	41	53	67	83
i= 2	0	0	<b>1</b>	<b>2</b>	<b>5</b>	8	14	20	30	40	55	70	91
i= 3	0	0	0	<b>1</b>	<b>2</b>	<b>5</b>	<b>9</b>	15	23	34	47	64	84
i= 4	0	0	0	0	<b>1</b>	<b>2</b>	<b>5</b>	<b>9</b>	<b>16</b>	24	37	51	71
i= 5	0	0	0	0	0	<b>1</b>	<b>2</b>	<b>5</b>	<b>9</b>	<b>16</b>	<b>25</b>	38	54
i= 6	0	0	0	0	0	0	<b>1</b>	<b>2</b>	<b>5</b>	<b>9</b>	<b>16</b>	<b>25</b>	<b>39</b>

### Family 3: $\bar{g}_{((k+i), k^{a-1}), (k^a)}^{(k)}$

k	0	1	2	3	4	5	6	7	8	9	10	11	12
i= 0	<b>1</b>	1	3	4	7	9	14	17	24	29	38	45	57
i= 1	0	<b>1</b>	<b>2</b>	4	7	11	16	23	31	41	53	67	83
i= 2	0	0	<b>1</b>	<b>2</b>	<b>5</b>	8	14	20	30	40	55	70	91
i= 3	0	0	0	<b>1</b>	<b>2</b>	<b>5</b>	<b>9</b>	15	23	34	47	64	84

### Theorem (L. C., 2015)

*The stable value of the  $j^{\text{th}}$  diagonal corresponds to the reduced Kronecker coefficients  $\bar{g}_{(k^a), (2k-j, k^{a-1})}^{(k)}$ , when  $k \geq 2j$  and their generating function is*

$$\mathcal{G}_a = \frac{1}{(1-x)^2(1-x^2)^3 \dots (1-x^{a-1})^3(1-x^a)^2(1-x^{a+1})}$$

# CONTENTS

Framework

Reduced Kronecker coefficients

Generating functions

**Plane partitions**

Quasipolynomials



# PLANE PARTITIONS

A **plane partition** is a two-dimensional array of non-negative integers  $n_{i,j}$  (with positive integer indices  $i$  and  $j$ ) that is non-increasing in both indices and for which only finitely many of the  $n_{i,j}$  are nonzero.

# PLANE PARTITIONS


A **plane partition** is a two-dimensional array of non-negative integers  $n_{i,j}$  (with positive integer indices  $i$  and  $j$ ) that is non-increasing in both indices and for which only finitely many of the  $n_{i,j}$  are nonzero.

5	3	2	1
4	2	2	
2	2		
2	1		

# PLANE PARTITIONS

A **plane partition** is a two-dimensional array of non-negative integers  $n_{i,j}$  (with positive integer indices  $i$  and  $j$ ) that is non-increasing in both indices and for which only finitely many of the  $n_{i,j}$  are nonzero.

5	3	2	1
4	2	2	
2	2		
2	1		



# PLANE PARTITIONS

A **plane partition** is a two-dimensional array of non-negative integers  $n_{i,j}$  (with positive integer indices  $i$  and  $j$ ) that is non-increasing in both indices and for which only finitely many of the  $n_{i,j}$  are nonzero.

## Theorem (MacMahon, 1915)

*Let  $r = \min(a, l)$  and  $s = \max(a, l)$ . Then, the generating function for the plane partitions fitting inside an  $l \times a$  rectangle is*

$$\prod_{j=r}^s \left( \frac{1}{1-x^j} \right)^r \cdot \prod_{i=1}^{r-1} \left( \frac{1}{1-x^i} \right)^i \left( \frac{1}{1-x^{s+i}} \right)^{r-i}$$

# RELATION WITH FAMILIES 1 AND 3

## Theorem (L. C. - M. Rosas, 2015)


*The reduced Kronecker coefficient  $\bar{g}_{(k^a), (k^a)}^{(k)}$  counts the number of plane partitions of  $k$  fitting inside a  $2 \times a$  rectangle.*

# RELATION WITH FAMILIES 1 AND 3

## Theorem (L. C. - M. Rosas, 2015)

*The reduced Kronecker coefficient  $\bar{g}_{(k^a), (k^a)}^{(k)}$  counts the number of plane partitions of  $k$  fitting inside a  $2 \times a$  rectangle.*

The reduced Kronecker coefficient for  $a = 4$  and  $k = 3$  is  $\bar{g}_{(3,3,3,3), (3,3,3,3)}^{(3)} = 5$ .



# RELATION WITH FAMILIES 1 AND 3

## Theorem (L. C. - M. Rosas, 2015)

*The reduced Kronecker coefficient  $\bar{g}_{(k^a), (k^a)}^{(k)}$  counts the number of plane partitions of  $k$  fitting inside a  $2 \times a$  rectangle.*

## Theorem (L. C. , 2015)

*For the stable values of the  $j^{\text{th}}$  diagonal in Family 3,*

$$\bar{\bar{g}}_a(j) = \sum_{l=0}^j \# \left\{ \begin{array}{c} \text{plane partitions} \\ \text{of } l \\ \text{in } 3 \times (a-1) \end{array} \right\} \# \left\{ \begin{array}{c} \text{plane partitions} \\ \text{of } j-l \\ \text{in } 2 \times 1 \end{array} \right\}$$

# CONTENTS

Framework

Reduced Kronecker coefficients

Generating functions

Plane partitions

Quasipolynomials



## Theorem (L. C. - M. Rosas, 2015)

Let  $\ell$  be the least common multiple of  $1, 2, \dots, a, a + 1$ .

The coefficients  $\bar{g}_{(k^a), (k^a)}^{(k)}$  are described by a quasipolynomial of degree  $2a - 1$  and period dividing  $\ell$ . In fact, we have checked that the period is exactly  $\ell$  for  $a \leq 10$ .

For  $a = 2$ , the coefficients are given by the quasipolynomial of degree 3 and period 6:

$$\left\{ \begin{array}{ll} 1/72k^3 + 1/6k^2 + 2/3k + 1 & \text{if } k \equiv 0 \pmod{6} \\ 1/72k^3 + 1/6k^2 + 13/24k + 5/18 & \text{if } k \equiv 1 \pmod{6} \\ 1/72k^3 + 1/6k^2 + 2/3k + 8/9 & \text{if } k \equiv 2 \pmod{6} \\ 1/72k^3 + 1/6k^2 + 13/24k + 1/2 & \text{if } k \equiv 3 \pmod{6} \\ 1/72k^3 + 1/6k^2 + 2/3k + 7/9 & \text{if } k \equiv 4 \pmod{6} \\ 1/72k^3 + 1/6k^2 + 13/24k + 7/18 & \text{if } k \equiv 5 \pmod{6} \end{array} \right.$$

## DESCRIPTION FOR FAMILY 3

### Theorem (L. C., 2015)

*Let  $\ell$  be the least common multiple of  $1, 2, \dots, a, a + 1$ .*

*The coefficients  $\overline{g}_a(j)$  are described by a quasipolynomial of degree  $3a - 2$  and period dividing  $\ell$ . In fact, we have checked that the period is exactly  $\ell$  for  $a \leq 7$ .*

For  $a = 2$ , the coefficients are given by the quasipolynomial of degree 4 and period 6:

$$\left\{ \begin{array}{ll} 1/288j^4 + 1/16j^3 + 7/18j^2 + j + 1 & j \equiv 0 \pmod{6} \\ 1/288j^4 + 1/16j^3 + 7/18j^2 + 15/16j + 175/288 & j \equiv 1 \pmod{6} \\ 1/288j^4 + 1/16j^3 + 7/18j^2 + j + 8/9 & j \equiv 2 \pmod{6} \\ 1/288j^4 + 1/16j^3 + 7/18j^2 + 15/16j + 23/32 & j \equiv 3 \pmod{6} \\ 1/288j^4 + 1/16j^3 + 7/18j^2 + j + 8/9 & j \equiv 4 \pmod{6} \\ 1/288j^4 + 1/16j^3 + 7/18j^2 + 15/16j + 175/288 & j \equiv 5 \pmod{6} \end{array} \right.$$

# Thank you very much!



# ¡Muchas gracias!

