Exercises: Algebraic Structures on Combinatorial Species

Marcelo Aguiar

ECCO 2016

1. Enumeration via Species

- (1) (a) * Convince yourself that the species Π is indeed a set species.
 - (b) * For any finite set I, consider the collection $P[I] = I \cup \{1, 2\}$. Is P a set species?
- (2) Let I be the species of *involutions* (i.e., I[J] is the set of bijections $\sigma: J \to J$ such that $\sigma^2 = \mathrm{id}$) and P the species of *perfect matchings* (i.e., P[J] is the set of partitions of J into 2-element sets). Let E_{1,2} be the species characteristic of singletons and doubletons:

$$\mathbf{E}_{1,2}[J] = \begin{cases} \{*_J\} & \text{if } |J| = 1 \text{ or } 2, \\ \emptyset & \text{otherwise.} \end{cases}$$

(a) Show that $I = E \circ E_{1,2}$ and deduce that

$$I(x) = e^{x + \frac{x^2}{2}}.$$

(b) Show that $I = E \cdot P$. Deduce that

$$P(x) = e^{\frac{x^2}{2}}$$

and that the number of perfect matchings on [2n] is $(2n-1)!! := (2n-1) \cdot (2n-3) \cdot \cdot \cdot 3 \cdot 1$.

(3) Let P and Q be finite set species. Show that

$$(P \cdot Q)(x) = P(x)Q(x).$$

On the right hand side we have the usual product of formal power series, on the left hand side we have the Cauchy product of the two species.

(4) Show that $L \circ E_+ = \Sigma$, and deduce that

$$\Sigma(x) = \frac{1}{2 - e^x}.$$

This is the generating function for the number of set compositions (the ballot numbers).

(5) (a) Let P be a set species. Show that the symmetric group S_n acts on the set P[n] by

$$\sigma \cdot x = P[\sigma](x).$$

(b) Let $f: P \to Q$ be a morphism of set species. Show that the map $f_{[n]}: P[n] \to Q[n]$ is a morphism of S_n -sets; that is,

$$f_{[n]}(\sigma \cdot x) = \sigma \cdot f_{[n]}(x)$$

1

for all $x \in P[n]$ and $\sigma \in S_n$.

- (c) Show that the action of S_n on L[n] has one orbit, while the action on B[n] has as many orbits as partitions of n.
- (d) Deduce that L and B are not isomorphic.

2. Monoids and Hopf monoids

- (1) * Convince yourself that concatenation turns the species Σ of set compositions into a monoid, and L into a submonoid.
- (2) * Verify that the species Π of set partitions satisfies the compatibility axiom of a bimonoid (with the structure given in the lecture).
- (3) (a) Let P and Q be monoid species. Define a monoid structure on the species $P \cdot Q$.
 - (b) Deduce monoid structures on $E \cdot E$ and on $E^{\cdot k}$, and describe them explicitly in terms of subsets and functions.
- (4) Show that the antipode of the exponential Hopf monoid E is given by

$$S_I(*_I) = (-1)^{|I|} *_I.$$

(5) Let H be a bimonoid. Let End(**H**) denote the set of all morphisms of vector species $f: \mathbf{H} \to \mathbf{H}$. It is a vector space under pointwise addition and scalar multiplication. The convolution f*g of two morphisms f and g is defined by

$$(f * g)_I(x) = \sum_{S \sqcup T = I} f_S(x|_S) \cdot g_T(x/_S).$$

for all finite sets I and all $x \in H[I]$ (and then extended linearly to $\mathbf{H}[I]$). Let $u : \mathbf{H} \to \mathbf{H}$ be the linear extension of $\iota \epsilon : \mathbf{H} \to \mathbf{H}$.

- (a) Show that convolution is associative and unital, with u being the unit. In this manner $\operatorname{End}(\mathbf{H})$ is an algebra.
- (b) Show that H is a Hopf monoid if and only if id: $\mathbf{H} \to \mathbf{H}$ is invertible in the algebra $\operatorname{End}(\mathbf{H})$, and that in this case S is the convolution inverse of id.
- (c) Deduce the uniqueness of the antipode.
- (6) Show that a morphism $f: H \to K$ of connected bimonoids preserves the antipodes. In other words,

$$f_I(s_I(x)) = s_I(f_I(x))$$

for all I and $x \in H[I]$.