

Enumerating lattice 3-polytopes

Mónica Blanco, Francisco Santos


Departamento de Matemáticas Estadística y Computación
Universidad de Cantabria, SPAIN

June 17, 2016

5 Encuentro Colombiano de Combinatoria,
Universidad Nacional Sede Medellín


Definition

- **Lattice polytope** $P :=$
convex hull of a finite set of points in \mathbb{Z}^d
(or in a d -dimensional lattice).




Definition

- **Lattice polytope** $P :=$
convex hull of a finite set of points in \mathbb{Z}^d
(or in a d -dimensional lattice).




Definition

- **Lattice polytope** $P :=$
convex hull of a finite set of points in \mathbb{Z}^d
(or in a d -dimensional lattice).




Definition

- ▶ **Lattice polytope** $P :=$
convex hull of a finite set of points in \mathbb{Z}^d
(or in a d -dimensional lattice).
- ▶ **Size** of $P :=$
number of lattice points in P : $|P \cap \mathbb{Z}^d|$




Definition

- ▶ **Lattice polytope** $P :=$
convex hull of a finite set of points in \mathbb{Z}^d
(or in a d -dimensional lattice).
- ▶ **Size** of $P :=$
number of lattice points in P : $|P \cap \mathbb{Z}^d|$



Definition


- ▶ **Lattice polytope** $P :=$
convex hull of a finite set of points in \mathbb{Z}^d
(or in a d -dimensional lattice).
- ▶ **Size** of $P :=$
number of lattice points in P : $|P \cap \mathbb{Z}^d|$
- ▶ **Volume** of $P :=$
volume “normalized to the lattice” =
 $d! \times$ Euclidean volume.



Definition

- ▶ **Lattice polytope** $P :=$
convex hull of a finite set of points in \mathbb{Z}^d
(or in a d -dimensional lattice).
- ▶ **Size** of $P :=$
number of lattice points in P : $|P \cap \mathbb{Z}^d|$
- ▶ **Volume** of $P :=$
volume “normalized to the lattice” =
 $d! \times$ Euclidean volume.


The *smaller* lattice d -polytopes have
volume 1, and the volume is always an
integer number.



Definition

- ▶ **Lattice polytope** $P :=$
convex hull of a finite set of points in \mathbb{Z}^d
(or in a d -dimensional lattice).
- ▶ **Size** of $P :=$
number of lattice points in P : $|P \cap \mathbb{Z}^d|$
- ▶ **Volume** of $P :=$
volume “normalized to the lattice” =
 $d! \times$ Euclidean volume.


The *smaller* lattice d -polytopes have volume 1, and the volume is always an integer number.



Definition

- ▶ **Lattice polytope** $P :=$
convex hull of a finite set of points in \mathbb{Z}^d
(or in a d -dimensional lattice).
- ▶ **Size** of $P :=$
number of lattice points in P : $|P \cap \mathbb{Z}^d|$
- ▶ **Volume** of $P :=$
volume “normalized to the lattice” =
 $d! \times$ Euclidean volume.

The *smaller* lattice d -polytopes have volume 1, and the volume is always an integer number.



Unimodular equivalence

A **unimodular transformation** is a linear integer map $t : \mathbb{R}^d \rightarrow \mathbb{R}^d$ that preserves the lattice. That is,

$$t(x) = A \cdot x + b, \quad x \in \mathbb{R}^d$$

for $A \in \mathbb{Z}^{d \times d}$, $\det(A) = \pm 1$ and $b \in \mathbb{Z}^d$.

($t \in GL(n, \mathbb{Z})$ + translations).

Two lattice polytopes P and Q are said **equivalent** or **unimodularly equivalent** if there is an affine unimodular transformation t such that $t(P) = Q$.

Unimodular equivalence

A **unimodular transformation** is a linear integer map $t : \mathbb{R}^d \rightarrow \mathbb{R}^d$ that preserves the lattice. That is,

$$t(x) = A \cdot x + b, \quad x \in \mathbb{R}^d$$

for $A \in \mathbb{Z}^{d \times d}$, $\det(A) = \pm 1$ and $b \in \mathbb{Z}^d$.


($t \in GL(n, \mathbb{Z})$ + translations).

Two lattice polytopes P and Q are said **equivalent** or **unimodularly equivalent** if there is an affine unimodular transformation t such that $t(P) = Q$.


Remark

Size and volume are invariant modulo unimodular equivalence.

Examples of unimodular transformations




Examples of unimodular transformations



► $\vec{x} \mapsto \vec{x} + \vec{b}$


Examples of unimodular transformations

$$\vec{x} \mapsto \vec{x} + \vec{b}$$




Examples of unimodular transformations


► $\vec{x} \mapsto \vec{x} + \vec{b}$




Examples of unimodular transformations




► $\vec{x} \mapsto \vec{x} + \vec{b}$



► $\vec{x} \mapsto -\vec{x}$




Examples of unimodular transformations



► $\vec{x} \mapsto \vec{x} + \vec{b}$

► $\vec{x} \mapsto -\vec{x}$


Examples of unimodular transformations



► $\vec{x} \mapsto \vec{x} + \vec{b}$

► $\vec{x} \mapsto -\vec{x}$

Examples of unimodular transformations




► $\vec{x} \mapsto \vec{x} + \vec{b}$

► $\vec{x} \mapsto -\vec{x}$

► $(x, y) \mapsto (-x, y)$

Examples of unimodular transformations




► $\vec{x} \mapsto \vec{x} + \vec{b}$

► $\vec{x} \mapsto -\vec{x}$

► $(x, y) \mapsto (-x, y)$

Examples of unimodular transformations




► $\vec{x} \mapsto \vec{x} + \vec{b}$


► $\vec{x} \mapsto -\vec{x}$

► $(x, y) \mapsto (-x, y)$


Examples of unimodular transformations



► $\vec{x} \mapsto \vec{x} + \vec{b}$




► $\vec{x} \mapsto -\vec{x}$




► $(x, y) \mapsto (-x, y)$




► $(x, y) \mapsto$
 $(x + (-1)y, y)$




Examples of unimodular transformations




► $\vec{x} \mapsto \vec{x} + \vec{b}$





► $\vec{x} \mapsto -\vec{x}$




► $(x, y) \mapsto (-x, y)$




► $(x, y) \mapsto$
 $(x + (-1)y, y)$




Examples of unimodular transformations



► $\vec{x} \mapsto \vec{x} + \vec{b}$



► $\vec{x} \mapsto -\vec{x}$



► $(x, y) \mapsto (-x, y)$



► $(x, y) \mapsto$
 $(x + (-1)y, y)$



Examples of unimodular transformations




$$\vec{x} \mapsto \vec{x} + \vec{b}$$



$$\vec{x} \mapsto -\vec{x}$$




$$(x, y) \mapsto (-x, y)$$



$$(x, y) \mapsto (x + (-1)y, y)$$



Examples of unimodular transformations



► $\vec{x} \mapsto \vec{x} + \vec{b}$


► $\vec{x} \mapsto -\vec{x}$

► $(x, y) \mapsto (-x, y)$

► $(x, y) \mapsto$
 $(x + (-1)y, y)$

► $(x, y) \mapsto$
 $(x + 3y, y)$

Examples of unimodular transformations



► $\vec{x} \mapsto \vec{x} + \vec{b}$


► $\vec{x} \mapsto -\vec{x}$

► $(x, y) \mapsto (-x, y)$

► $(x, y) \mapsto$
 $(x + (-1)y, y)$

► $(x, y) \mapsto$
 $(x + 3y, y)$

Examples of unimodular transformations



► $\vec{x} \mapsto \vec{x} + \vec{b}$

► $\vec{x} \mapsto -\vec{x}$

► $(x, y) \mapsto (-x, y)$

► $(x, y) \mapsto$
 $(x + (-1)y, y)$

► $(x, y) \mapsto$
 $(x + 3y, y)$

Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES

Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES

For this, we separate by size: let $n \geq d + 1$

$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES

For this, we separate by size: let $n \geq d + 1$

$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$

Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES

For this, we separate by size: let $n \geq d + 1$

$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$



Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES

For this, we separate by size: let $n \geq d + 1$

$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$



- **Dimension 2:** for each n , the cardinal of $\mathcal{P}_2(n)$ is **finite** $|\mathcal{P}_2(n)| < \infty$


Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES

For this, we separate by size: let $n \geq d + 1$


$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$



- **Dimension 2:** for each n , the cardinal of $\mathcal{P}_2(n)$ is **finite** $|\mathcal{P}_2(n)| < \infty$

All polygons of sizes 3, 4 and 5:




Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES


For this, we separate by size: let $n \geq d + 1$

$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$



- **Dimension 2:** for each n , the cardinal of $\mathcal{P}_2(n)$ is **finite** $|\mathcal{P}_2(n)| < \infty$
All polygons of sizes 3, 4 and 5:




Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES


For this, we separate by size: let $n \geq d + 1$

$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$



- **Dimension 2:** for each n , the cardinal of $\mathcal{P}_2(n)$ is **finite** $|\mathcal{P}_2(n)| < \infty$
All polygons of sizes 3, 4 and 5:




Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES


For this, we separate by size: let $n \geq d + 1$

$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$



- **Dimension 2:** for each n , the cardinal of $\mathcal{P}_2(n)$ is **finite** $|\mathcal{P}_2(n)| < \infty$
All polygons of sizes 3, 4 and 5:




Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES

For this, we separate by size: let $n \geq d + 1$


$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$



- **Dimension 2:** for each n , the cardinal of $\mathcal{P}_2(n)$ is **finite** $|\mathcal{P}_2(n)| < \infty$

All polygons of sizes 3, 4 and 5:




Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES

For this, we separate by size: let $n \geq d + 1$



$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$



- **Dimension 2:** for each n , the cardinal of $\mathcal{P}_2(n)$ is **finite** $|\mathcal{P}_2(n)| < \infty$

All polygons of sizes 3, 4 and 5:





Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES

For this, we separate by size: let $n \geq d + 1$


$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$



- **Dimension 2:** for each n , the cardinal of $\mathcal{P}_2(n)$ is **finite** $|\mathcal{P}_2(n)| < \infty$

All polygons of sizes 3, 4 and 5:




Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES

For this, we separate by size: let $n \geq d + 1$


$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$



- **Dimension 2:** for each n , the cardinal of $\mathcal{P}_2(n)$ is **finite** $|\mathcal{P}_2(n)| < \infty$

All polygons of sizes 3, 4 and 5:




Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES

For this, we separate by size: let $n \geq d + 1$



$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$



- **Dimension 2:** for each n , the cardinal of $\mathcal{P}_2(n)$ is **finite** $|\mathcal{P}_2(n)| < \infty$

All polygons of sizes 3, 4 and 5:



Goal

WE WANT TO CLASSIFY (classes of) LATTICE d -POLYTOPES

For this, we separate by size: let $n \geq d + 1$



$$\mathcal{P}_d(n) := \{(\text{classes of}) \text{ lattice } d\text{-polytopes of size } n\}$$

- **Dimension 1:** for each $n \geq 2$, there is one lattice 1-polytope of size n , a segment of length $n - 1$: $|\mathcal{P}_1(n)| = 1$



- **Dimension 2:** for each n , the cardinal of $\mathcal{P}_2(n)$ is **finite** $|\mathcal{P}_2(n)| < \infty$

All polygons of sizes 3, 4 and 5:



Dimension 3, $n = 4$ (empty tetrahedra)


In dimension 3, Reeve tetrahedra are
infinitely many lattice 3-polytopes with
4 lattice points

$$|\mathcal{P}_3(4)| = \infty$$

Dimension 3, $n = 4$ (empty tetrahedra)

In dimension 3, Reeve tetrahedra are
infinitely many lattice 3-polytopes with
4 lattice points


$$|\mathcal{P}_3(4)| = \infty$$



Dimension 3, $n = 4$ (empty tetrahedra)

In dimension 3, Reeve tetrahedra are
infinitely many lattice 3-polytopes with
4 lattice points


$$|\mathcal{P}_3(4)| = \infty$$



Dimension 3, $n = 4$ (empty tetrahedra)

In dimension 3, Reeve tetrahedra are
infinitely many lattice 3-polytopes with
4 lattice points


$$|\mathcal{P}_3(4)| = \infty$$



Dimension 3, $n = 4$ (empty tetrahedra)

In dimension 3, Reeve tetrahedra are
infinitely many lattice 3-polytopes with
4 lattice points


$$|\mathcal{P}_3(4)| = \infty$$



Dimension 3, $n = 4$ (empty tetrahedra)

In dimension 3, Reeve tetrahedra are
infinitely many lattice 3-polytopes with
4 lattice points


$$|\mathcal{P}_3(4)| = \infty$$



Dimension 3, $n = 4$ (empty tetrahedra)

In dimension 3, Reeve tetrahedra are infinitely many lattice 3-polytopes with 4 lattice points $|\mathcal{P}_3(4)| = \infty$

Elements in $\mathcal{P}_3(4)$ are called **empty tetrahedra**: tetrahedra in which the only lattice points are the four vertices. Their classification is classical (White 1964):



$$\mathcal{P}_3(4) = \{T(p, q) \mid p, q \in \mathbb{Z}, 0 < p \leq q, \gcd(p, q) = 1\},$$

where $T(p, q) := \text{conv} \{(0, 0, 0), (1, 0, 0), (0, 0, 1), (p, q, 1)\}$.

Dimension 3, $n = 5, 6$

- For sizes $n = 5, 6$, Santos and B. (2014) classified all lattice 3-polytopes of those sizes. In terms of cardinality,

$$|\mathcal{P}_3(5)| = \infty, \quad |\mathcal{P}_3(6)| = \infty$$

Dimension 3, $n = 5, 6$

- ▶ For sizes $n = 5, 6$, Santos and B. (2014) classified all lattice 3-polytopes of those sizes. In terms of cardinality,

$$|\mathcal{P}_3(5)| = \infty, \quad |\mathcal{P}_3(6)| = \infty$$

- ▶ The classification was done via oriented matroids (information on the position of the set of points in the space). For 5 points, there are such possible oriented matroids, for 6 points there are 55.

Dimension 3, $n = 5, 6$

- ▶ For sizes $n = 5, 6$, Santos and B. (2014) classified all lattice 3-polytopes of those sizes. In terms of cardinality,

$$|\mathcal{P}_3(5)| = \infty, \quad |\mathcal{P}_3(6)| = \infty$$

- ▶ The classification was done via oriented matroids (information on the position of the set of points in the space). For 5 points, there are such possible oriented matroids, for 6 points there are 55. But for 7 points there are already 5000...

Dimension 3, $n = 5, 6$

- ▶ For sizes $n = 5, 6$, Santos and B. (2014) classified all lattice 3-polytopes of those sizes. In terms of cardinality,

$$|\mathcal{P}_3(5)| = \infty, \quad |\mathcal{P}_3(6)| = \infty$$

- ▶ The classification was done via oriented matroids (information on the position of the set of points in the space). For 5 points, there are such possible oriented matroids, for 6 points there are 55.
But for 7 points there are already 5000... and for 8 points is around the 10 millions!!!!

Dimension 3, $n = 5, 6$

- For sizes $n = 5, 6$, Santos and B. (2014) classified all lattice 3-polytopes of those sizes. In terms of cardinality,

$$|\mathcal{P}_3(5)| = \infty, \quad |\mathcal{P}_3(6)| = \infty$$

- The classification was done via oriented matroids (information on the position of the set of points in the space). For 5 points, there are such possible oriented matroids, for 6 points there are 55. But for 7 points there are already 5000... and for 8 points is around the 10 millions!!!!
So another approach is required. We notice that the number of lattice points is significantly affected by another property of lattice polytopes.

Dimension 3, $n = 5, 6$

- For sizes $n = 5, 6$, Santos and B. (2014) classified all lattice 3-polytopes of those sizes. In terms of cardinality,

$$|\mathcal{P}_3(5)| = \infty, \quad |\mathcal{P}_3(6)| = \infty$$

- The classification was done via oriented matroids (information on the position of the set of points in the space). For 5 points, there are such possible oriented matroids, for 6 points there are 55.
But for 7 points there are already 5000... and for 8 points is around the 10 millions!!!!
So another approach is required. We notice that the number of lattice points is significantly affected by another property of lattice polytopes.

... If we look at the **WIDTH** of 3-polytopes of sizes 4, 5 and 6, we see that only finitely many of them have width > 1

Dimension 3, $n = 5, 6$

- For sizes $n = 5, 6$, Santos and B. (2014) classified all lattice 3-polytopes of those sizes. In terms of cardinality,

$$|\mathcal{P}_3(5)| = \infty, \quad |\mathcal{P}_3(6)| = \infty$$

- The classification was done via oriented matroids (information on the position of the set of points in the space). For 5 points, there are such possible oriented matroids, for 6 points there are 55.

But for 7 points there are already 5000... and for 8 points is around the 10 millions!!!!

So another approach is required. We notice that the number of lattice points is significantly affected by another property of lattice polytopes.

... If we look at the **WIDTH** of 3-polytopes of sizes 4, 5 and 6, we see that only finitely many of them have width > 1

Size	4	5	6
width 1	∞	∞	∞
width 2	—	9	74
width 3	—	—	2

(Lattice) Width


Definition

- ▶ **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
= length of the interval $f(P)$

(Lattice) Width

Definition


- **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
= length of the interval $f(P)$



(Lattice) Width

Definition


- ▶ **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
= length of the interval $f(P)$
- ▶ **Width of P :** = Minimum width of P with respect to a linear NON-CONSTANT, INTEGER functional



(Lattice) Width

Definition

- ▶ **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
= length of the interval $f(P)$




- ▶ **Width of P :** = Minimum width of P with respect to a linear NON-CONSTANT, INTEGER functional = minimum lattice distance between two parallel lattice hyperplanes enclosing P


(Lattice) Width

Definition


- ▶ **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
= length of the interval $f(P)$




- ▶ **Width of P :** = Minimum width of P with respect to a linear NON-CONSTANT, INTEGER functional = minimum lattice distance between two parallel lattice hyperplanes enclosing P



Width: 2



Width: 1




Width: 2


(Lattice) Width

Definition


- **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
= length of the interval $f(P)$




- **Width of P :** = Minimum width of P with respect to a linear NON-CONSTANT, INTEGER functional = minimum lattice distance between two parallel lattice hyperplanes enclosing P



Width: 2



Width: 1




Width: 2


(Lattice) Width

Definition


- **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
= length of the interval $f(P)$




- **Width of P :** = Minimum width of P with respect to a linear NON-CONSTANT, INTEGER functional = minimum lattice distance between two parallel lattice hyperplanes enclosing P



Width: 2



Width: 1




Width: 2


(Lattice) Width

Definition


- **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
= length of the interval $f(P)$




- **Width of P :** = Minimum width of P with respect to a linear NON-CONSTANT, INTEGER functional = minimum lattice distance between two parallel lattice hyperplanes enclosing P



Width: 2



Width: 1




Width: 2


(Lattice) Width

Definition


- **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
= length of the interval $f(P)$




- **Width of P :** = Minimum width of P with respect to a linear NON-CONSTANT, INTEGER functional = minimum lattice distance between two parallel lattice hyperplanes enclosing P



Width: 2



Width: 1




Width: 2


(Lattice) Width

Definition


- **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
 $=$ length of the interval $f(P)$




- **Width of P :** Minimum width of P with respect to a linear NON-CONSTANT, INTEGER functional $=$ minimum lattice distance between two parallel lattice hyperplanes enclosing P



Width: 2



Width: 1




Width: 2


(Lattice) Width

Definition


- ▶ **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
= length of the interval $f(P)$




- ▶ **Width of P :** = Minimum width of P with respect to a linear NON-CONSTANT, INTEGER functional = minimum lattice distance between two parallel lattice hyperplanes enclosing P



Width: 2



Width: 1




Width: 2


(Lattice) Width

Definition


- ▶ **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
= length of the interval $f(P)$




- ▶ **Width of P :** = Minimum width of P with respect to a linear NON-CONSTANT, INTEGER functional = minimum lattice distance between two parallel lattice hyperplanes enclosing P



Width: 2



Width: 1




Width: 2


(Lattice) Width

Definition


- **Width of P with respect to f ,**
for a linear functional $f : \mathbb{R}^d \rightarrow \mathbb{R}$
= length of the interval $f(P)$




- **Width of P :** = Minimum width of P with respect to a linear NON-CONSTANT, INTEGER functional = minimum lattice distance between two parallel lattice hyperplanes enclosing P



Width: 2



Width: 1



Width: 2

Width $> 1 \implies$ **finite** number of classes (for $d = 3$)

$$\mathcal{P}_d^*(n) := \{P \in \mathcal{P}_d(n) \mid \text{width}(P) > 1\}$$

Width $> 1 \implies$ **finite** number of classes (for $d = 3$)

$$\mathcal{P}_d^*(n) := \{P \in \mathcal{P}_d(n) \mid \text{width}(P) > 1\}$$

For each $n \geq 4$:

- There are **infinitely** many equivalence classes of **width 1**:

$$|\mathcal{P}_d(n) \setminus \mathcal{P}_d^*(n)| = \infty.$$


Width $> 1 \implies$ **finite** number of classes (for $d = 3$)

$$\mathcal{P}_d^*(n) := \{P \in \mathcal{P}_d(n) \mid \text{width}(P) > 1\}$$

For each $n \geq 4$:

- There are **infinitely** many equivalence classes of **width 1**:

$$|\mathcal{P}_d(n) \setminus \mathcal{P}_d^*(n)| = \infty.$$




Width $> 1 \implies$ **finite** number of classes (for $d = 3$)

$$\mathcal{P}_d^*(n) := \{P \in \mathcal{P}_d(n) \mid \text{width}(P) > 1\}$$

For each $n \geq 4$:

- There are **infinitely** many equivalence classes of **width 1**:

$$|\mathcal{P}_d(n) \setminus \mathcal{P}_d^*(n)| = \infty.$$



Width $> 1 \implies$ **finite** number of classes (for $d = 3$)

$$\mathcal{P}_d^*(n) := \{P \in \mathcal{P}_d(n) \mid \text{width}(P) > 1\}$$

For each $n \geq 4$:

- There are **infinitely** many equivalence classes of **width 1**:


$$|\mathcal{P}_d(n) \setminus \mathcal{P}_d^*(n)| = \infty.$$

- If the width is > 1 :

Lemma (B.-Santos, 2014+)

*For each $n \geq 4$, there are **finitely** many lattice 3-polytopes of width greater than one and size n . That is,*

$$|\mathcal{P}_3^*(n)| < \infty, \text{ for each } n \geq 4$$



Width $> 1 \implies$ **finite** number of classes (for $d = 3$)

$$\mathcal{P}_d^*(n) := \{P \in \mathcal{P}_d(n) \mid \text{width}(P) > 1\}$$

For each $n \geq 4$:

- There are **infinitely** many equivalence classes of **width 1**:

$$|\mathcal{P}_d(n) \setminus \mathcal{P}_d^*(n)| = \infty.$$


- If the width is > 1 :

Lemma (B.-Santos, 2014+)

*For each $n \geq 4$, there are **finitely** many lattice 3-polytopes of width greater than one and size n . That is,*

$$|\mathcal{P}_3^*(n)| < \infty, \text{ for each } n \geq 4$$

WE CAN ENUMERATE the complete list $\mathcal{P}_3^*(n)$ of lattice 3-polytopes of size n AND WIDTH > 1 , for each n



Essential vertices

From now on, let $P \in \mathcal{P}_d^*(n)$. For each vertex $v \in \text{vert}(P)$, we denote by P^v the polytope $\text{conv}(P \setminus \{v\} \cap \mathbb{Z}^d) \subset \mathbb{R}^d$. This polytope has size $n - 1$ but it is not necessarily full-dimensional.

Essential vertices

From now on, let $P \in \mathcal{P}_d^*(n)$. For each vertex $v \in \text{vert}(P)$, we denote by P^v the polytope $\text{conv}(P \setminus \{v\} \cap \mathbb{Z}^d) \subset \mathbb{R}^d$. This polytope has size $n - 1$ but it is not necessarily full-dimensional.

Definition (Essential vertex)


We say that v is an **essential vertex** of P if $P^v \notin \mathcal{P}_d^*(n - 1)$. That is, if P^v is either $(d - 1)$ -dimensional or has width one.

Essential vertices

From now on, let $P \in \mathcal{P}_d^*(n)$. For each vertex $v \in \text{vert}(P)$, we denote by P^v the polytope $\text{conv}(P \setminus \{v\} \cap \mathbb{Z}^d) \subset \mathbb{R}^d$. This polytope has size $n - 1$ but it is not necessarily full-dimensional.

Definition (Essential vertex)

We say that v is an **essential vertex** of P if $P^v \notin \mathcal{P}_d^*(n-1)$. That is, if P^v is either $(d-1)$ -dimensional or has width one.



Essential vertices


From now on, let $P \in \mathcal{P}_d^*(n)$. For each vertex $v \in \text{vert}(P)$, we denote by P^v the polytope $\text{conv}(P \setminus \{v\} \cap \mathbb{Z}^d) \subset \mathbb{R}^d$. This polytope has size $n - 1$ but it is not necessarily full-dimensional.

Definition (Essential vertex)

We say that v is an **essential vertex** of P if $P^v \notin \mathcal{P}_d^*(n - 1)$. That is, if P^v is either $(d - 1)$ -dimensional or has width one.

$$\bullet (-1, 1, 2)$$

$P^{(-1,1,2)}$ is 2-dimensional




Essential vertices

From now on, let $P \in \mathcal{P}_d^*(n)$. For each vertex $v \in \text{vert}(P)$, we denote by P^v the polytope $\text{conv}(P \setminus \{v\} \cap \mathbb{Z}^d) \subset \mathbb{R}^d$. This polytope has size $n - 1$ but it is not necessarily full-dimensional.

Definition (Essential vertex)

We say that v is an **essential vertex** of P if $P^v \notin \mathcal{P}_d^*(n - 1)$. That is, if P^v is either $(d - 1)$ -dimensional or has width one.




$P^{(1,0,0)}$ has width one

Essential vertices

From now on, let $P \in \mathcal{P}_d^*(n)$. For each vertex $v \in \text{vert}(P)$, we denote by P^v the polytope $\text{conv}(P \setminus \{v\} \cap \mathbb{Z}^d) \subset \mathbb{R}^d$. This polytope has size $n - 1$ but it is not necessarily full-dimensional.

Definition (Essential vertex)

We say that v is an **essential vertex** of P if $P^v \notin \mathcal{P}_d^*(n - 1)$. That is, if P^v is either $(d - 1)$ -dimensional or has width one.




Essential vertices

From now on, let $P \in \mathcal{P}_d^*(n)$. For each vertex $v \in \text{vert}(P)$, we denote by P^v the polytope $\text{conv}(P \setminus \{v\} \cap \mathbb{Z}^d) \subset \mathbb{R}^d$. This polytope has size $n - 1$ but it is not necessarily full-dimensional.

Definition (Essential vertex)

We say that v is an **essential vertex** of P if $P^v \notin \mathcal{P}_d^*(n - 1)$. That is, if P^v is either $(d - 1)$ -dimensional or has width one.



Quasiminimal vs. Merged

Definition

Let $P \in \mathcal{P}_d^*(n)$.

- ▶ We say that P is **quasiminimal** if it has ≤ 1 NON-essential vertices.
That is, if there is **at most** one vertex v such that $P^v \in \mathcal{P}_d^*(n-1)$.

Quasiminimal vs. Merged

Definition

Let $P \in \mathcal{P}_d^*(n)$.

- ▶ We say that P is **quasiminimal** if it has ≤ 1 NON-essential vertices. That is, if there is **at most** one vertex v such that $P^v \in \mathcal{P}_d^*(n-1)$.
- ▶ We say that P is **merged** if there are ≥ 2 NON-essential vertices (i.e. $u, v \in \text{vert}(P)$ with $P^u, P^v \in \mathcal{P}_d^*(n-1)$) AND the polytope $P^{u,v} := \text{conv}(P^u \cap P^v \cap \mathbb{Z}^d)$ is still d -dimensional.

Quasiminimal vs. Merged

Definition

Let $P \in \mathcal{P}_d^*(n)$.


- ▶ We say that P is **quasiminimal** if it has ≤ 1 NON-essential vertices. That is, if there is **at most** one vertex v such that $P^v \in \mathcal{P}_d^*(n-1)$.
- ▶ We say that P is **merged** if there are ≥ 2 NON-essential vertices (i.e. $u, v \in \text{vert}(P)$ with $P^u, P^v \in \mathcal{P}_d^*(n-1)$) AND the polytope $P^{u,v} := \text{conv}(P^u \cap P^v \cap \mathbb{Z}^d)$ is still d -dimensional.
- ▶ We say that $P \in \mathcal{P}_d^*(n)$ is an **exception** if it is neither *quasiminimal* nor *merged*. That is, if it has ≥ 2 NON-essential vertices, AND **for all pairs** $u, v \in \text{vert}(P)$ of non-essential vertices, $P^u, P^v \in \mathcal{P}_d^*(n-1)$ are such that $P^{u,v} := \text{conv}(P^u \cap P^v \cap \mathbb{Z}^d)$ is $(d-1)$ -dimensional.

Quasiminimals

- ▶ We say that P is **quasiminimal** if it has ≤ 1 NON-essential vertices. That is, if there is **at most** one vertex v such that $P^v \in \mathcal{P}_d^*(n-1)$.

Quasiminimals

- We say that P is **quasiminimal** if it has ≤ 1 NON-essential vertices.
That is, if there is **at most** one vertex v such that $P^v \in \mathcal{P}_d^*(n-1)$.




$P^{(-1,1,2)}$ is 2-dimensional
 $P^{(1,0,0)}$ has width one
 $P^{(-1,0,0)}$ has width one
 $P^{(0,5,0)}$ has width larger than one

essential vertices

Quasiminimals

- We say that P is **quasiminimal** if it has ≤ 1 NON-essential vertices. That is, if there is **at most** one vertex v such that $P^v \in \mathcal{P}_d^*(n-1)$.



$P^{(-1,1,2)}$ is 2-dimensional
 $P^{(1,0,0)}$ has width one
 $P^{(-1,0,0)}$ has width one
 $P^{(0,5,0)}$ has width larger than one

essential vertices


We will denote by $\mathcal{Q}_d(n)$ the set of quasiminimal d -polytopes of size n .

Merged

- ▶ We say that P is **merged** if there are ≥ 2 NON-essential vertices (i.e. $u, v \in \text{vert}(P)$ with $P^u, P^v \in \mathcal{P}_d^*(n-1)$) AND the polytope $P^{u,v} := \text{conv}(P^u \cap P^v \cap \mathbb{Z}^d)$ is still d -dimensional.


Merged

- We say that P is **merged** if there are ≥ 2 NON-essential vertices (i.e. $u, v \in \text{vert}(P)$ with $P^u, P^v \in \mathcal{P}_d^*(n-1)$) AND the polytope $P^{u,v} := \text{conv}(P^u \cap P^v \cap \mathbb{Z}^d)$ is still d -dimensional.




Merged

- We say that P is **merged** if there are ≥ 2 NON-essential vertices (i.e. $u, v \in \text{vert}(P)$ with $P^u, P^v \in \mathcal{P}_d^*(n-1)$) AND the polytope $P^{u,v} := \text{conv}(P^u \cap P^v \cap \mathbb{Z}^d)$ is still d -dimensional.




Merged

- We say that P is **merged** if there are ≥ 2 NON-essential vertices (i.e. $u, v \in \text{vert}(P)$ with $P^u, P^v \in \mathcal{P}_d^*(n-1)$) AND the polytope $P^{u,v} := \text{conv}(P^u \cap P^v \cap \mathbb{Z}^d)$ is still d -dimensional.




Merged

- We say that P is **merged** if there are ≥ 2 NON-essential vertices (i.e. $u, v \in \text{vert}(P)$ with $P^u, P^v \in \mathcal{P}_d^*(n-1)$) AND the polytope $P^{u,v} := \text{conv}(P^u \cap P^v \cap \mathbb{Z}^d)$ is still d -dimensional.



Merged


- We say that P is **merged** if there are ≥ 2 NON-essential vertices (i.e. $u, v \in \text{vert}(P)$ with $P^u, P^v \in \mathcal{P}_d^*(n-1)$) AND the polytope $P^{u,v} := \text{conv}(P^u \cap P^v \cap \mathbb{Z}^d)$ is still d -dimensional.



We will denote by $\mathcal{M}_d(n)$ the set of merged d -polytopes of size n .


Exceptions

- We say that $P \in \mathcal{P}_d^*(n)$ is an **exception** if it is neither *quasiminimal* nor *merged*. That is, if it has ≥ 2 NON-essential vertices, AND for all pairs $u, v \in \text{vert}(P)$ of non-essential vertices, $P^u, P^v \in \mathcal{P}_d^*(n-1)$ are such that $P^{u,v} := \text{conv}(P^u \cap P^v \cap \mathbb{Z}^d)$ is $(d-1)$ -dimensional.



Exceptions

- We say that $P \in \mathcal{P}_d^*(n)$ is an **exception** if it is neither *quasiminimal* nor *merged*. That is, if it has ≥ 2 NON-essential vertices, AND for all pairs $u, v \in \text{vert}(P)$ of non-essential vertices, $P^u, P^v \in \mathcal{P}_d^*(n-1)$ are such that $P^{u,v} := \text{conv}(P^u \cap P^v \cap \mathbb{Z}^d)$ is $(d-1)$ -dimensional.




Theorem (Blanco and Santos)

There is a single lattice 3-dimensional exception, and it is of size $n = 6$:

$$|\mathcal{P}_3^*(6) \setminus (\mathcal{Q}_3(6) \cup \mathcal{M}_3(6))| = 1, \quad \mathcal{P}_3^*(n) = \mathcal{Q}_3(n) \cup \mathcal{M}_3(n), \text{ for all } n \geq 7.$$

Notice that $\mathcal{P}_3^*(4) = \emptyset$ and, hence, $\mathcal{P}_3^*(5) = \mathcal{Q}_3(5)$.

Classifying merged polytopes: size $n \rightarrow n - 1$



Size $n - 1 \rightarrow n$: Merging algorithm

INPUT: a finite list L of lattice d -polytopes of size $n - 1$ and width > 1 .

Size $n - 1 \rightarrow n$: Merging algorithm

INPUT: a finite list L of lattice d -polytopes of size $n - 1$ and width > 1 .


OUTPUT: the list $L' = \text{Merging}(L)$ of all lattice d -polytopes of size n obtained merging polytopes of L .

Size $n - 1 \rightarrow n$: Merging algorithm

INPUT: a finite list L of lattice d -polytopes of size $n - 1$ and width > 1 .

OUTPUT: the list $L' = \text{Merging}(L)$ of all lattice d -polytopes of size n obtained merging polytopes of L .

For each $P_1, P_2 \in L$, and for each vertex v_1 of P_1 and v_2 of P_2 :




Size $n - 1 \rightarrow n$: Merging algorithm

INPUT: a finite list L of lattice d -polytopes of size $n - 1$ and width > 1 .

OUTPUT: the list $L' = \text{Merging}(L)$ of all lattice d -polytopes of size n obtained merging polytopes of L .

For each $P_1, P_2 \in L$, and for each vertex v_1 of P_1 and v_2 of P_2 :

1. Let $P'_1 = \text{conv}(\mathbb{Z}^d \cap P_1 \setminus \{v_1\})$ and $P'_2 = \text{conv}(\mathbb{Z}^d \cap P_2 \setminus \{v_2\})$.




Size $n - 1 \rightarrow n$: Merging algorithm

INPUT: a finite list L of lattice d -polytopes of size $n - 1$ and width > 1 .

OUTPUT: the list $L' = \text{Merging}(L)$ of all lattice d -polytopes of size n obtained merging polytopes of L .

For each $P_1, P_2 \in L$, and for each vertex v_1 of P_1 and v_2 of P_2 :

1. Let $P'_1 = \text{conv}(\mathbb{Z}^d \cap P_1 \setminus \{v_1\})$ and $P'_2 = \text{conv}(\mathbb{Z}^d \cap P_2 \setminus \{v_2\})$.
2. Check if P'_1 and P'_2 are d -dimensional and equivalent. If they are, let $t : \mathbb{Z}^d \rightarrow \mathbb{Z}^d$ be an equivalence sending P'_1 to P'_2 (there are finitely many possibilities for t ; do step 3 for each).




Size $n - 1 \rightarrow n$: Merging algorithm

INPUT: a finite list L of lattice d -polytopes of size $n - 1$ and width > 1 .

OUTPUT: the list $L' = \text{Merging}(L)$ of all lattice d -polytopes of size n obtained merging polytopes of L .

For each $P_1, P_2 \in L$, and for each vertex v_1 of P_1 and v_2 of P_2 :

1. Let $P'_1 = \text{conv}(\mathbb{Z}^d \cap P_1 \setminus \{v_1\})$ and $P'_2 = \text{conv}(\mathbb{Z}^d \cap P_2 \setminus \{v_2\})$.
2. Check if P'_1 and P'_2 are d -dimensional and equivalent. If they are, let $t : \mathbb{Z}^d \rightarrow \mathbb{Z}^d$ be an equivalence sending P'_1 to P'_2 (there are finitely many possibilities for t ; do step 3 for each).
3. If the size of $P := \text{conv}(t(P_1) \cup P_2)$ equals n , add P to L' .




Size $n - 1 \rightarrow n$: Merging algorithm

INPUT: a finite list L of lattice d -polytopes of size $n - 1$ and width > 1 .

OUTPUT: the list $L' = \text{Merging}(L)$ of all lattice d -polytopes of size n obtained merging polytopes of L .

For each $P_1, P_2 \in L$, and for each vertex v_1 of P_1 and v_2 of P_2 :

1. Let $P'_1 = \text{conv}(\mathbb{Z}^d \cap P_1 \setminus \{v_1\})$ and $P'_2 = \text{conv}(\mathbb{Z}^d \cap P_2 \setminus \{v_2\})$.
2. Check if P'_1 and P'_2 are d -dimensional and equivalent. If they are, let $t : \mathbb{Z}^d \rightarrow \mathbb{Z}^d$ be an equivalence sending P'_1 to P'_2 (there are finitely many possibilities for t ; do step 3 for each).
3. If the size of $P := \text{conv}(t(P_1) \cup P_2)$ equals n , add P to L' .




Dimension 3: By definition, and since $\mathcal{P}_3^*(n - 1)$ is a finite list:

$\mathcal{M}_3(n) = \text{Merging}(\mathcal{P}_3^*(n - 1))$, for all n .


Quasiminimal polytopes

Let $P \in \mathcal{Q}_d(n)$ and, for each essential vertex $v \in \text{vert}(P)$, let $f_v : \mathbb{R}^d \rightarrow \mathbb{R}$ be an integer linear functional that gives width one (or zero) to P^v .



Quasiminimal polytopes

Let $P \in \mathcal{Q}_d(n)$ and, for each essential vertex $v \in \text{vert}(P)$, let $f_v : \mathbb{R}^d \rightarrow \mathbb{R}$ be an integer linear functional that gives width one (or zero) to P^v .




We distinguish two cases:

Definition (Boxed vs. spiked)

- If the set $\{f_v : v \text{ is essential vertex of } P\}$ linearly spans $(\mathbb{R}^d)^*$, then we can find d linearly independent f_v . We call these polytopes **boxed**, because *most of their lattice points lie in the vertices of a d -parallelepiped*.

Quasiminimal polytopes

Let $P \in \mathcal{Q}_d(n)$ and, for each essential vertex $v \in \text{vert}(P)$, let $f_v : \mathbb{R}^d \rightarrow \mathbb{R}$ be an integer linear functional that gives width one (or zero) to P^v .




We distinguish two cases:

Definition (Boxed vs. spiked)

- ▶ If the set $\{f_v : v \text{ is essential vertex of } P\}$ linearly spans $(\mathbb{R}^d)^*$, then we can find d linearly independent f_v . We call these polytopes **boxed**, because *most of their lattice points lie in the vertices of a d -parallelepiped*.
- ▶ If the set $\{f_v : v \text{ is essential vertex of } P\}$ does not linearly span $(\mathbb{R}^d)^*$, then there is a projection that respects all f_v . We call these polytopes **spiked**, because *most of their lattice points lie in a lattice segment*.

Quasiminimal polytopes


Let $P \in \mathcal{Q}_d(n)$ and, for each essential vertex $v \in \text{vert}(P)$, let $f_v : \mathbb{R}^d \rightarrow \mathbb{R}$ be an integer linear functional that gives width one (or zero) to P^v .



We distinguish two cases:


Definition (Boxed vs. spiked)

- ▶ If the set $\{f_v : v \text{ is essential vertex of } P\}$ linearly spans $(\mathbb{R}^d)^*$, then we can find d linearly independent f_v . We call these polytopes **boxed**, because *most of their lattice points lie in the vertices of a d -parallelepiped*.
- ▶ If the set $\{f_v : v \text{ is essential vertex of } P\}$ does not linearly span $(\mathbb{R}^d)^*$, then there is a projection that respects all f_v . We call these polytopes **spiked**, because *most of their lattice points lie in a lattice segment*.



Quasiminimal polytopes


Let $P \in \mathcal{Q}_d(n)$ and, for each essential vertex $v \in \text{vert}(P)$, let $f_v : \mathbb{R}^d \rightarrow \mathbb{R}$ be an integer linear functional that gives width one (or zero) to P^v .



We distinguish two cases:


Definition (Boxed vs. spiked)

- ▶ If the set $\{f_v : v \text{ is essential vertex of } P\}$ linearly spans $(\mathbb{R}^d)^*$, then we can find d linearly independent f_v . We call these polytopes **boxed**, because *most of their lattice points lie in the vertices of a d -parallelepiped*.
- ▶ If the set $\{f_v : v \text{ is essential vertex of } P\}$ does not linearly span $(\mathbb{R}^d)^*$, then there is a projection that respects all f_v . We call these polytopes **spiked**, because *most of their lattice points lie in a lattice segment*.



Quasiminimal polytopes


Let $P \in \mathcal{Q}_d(n)$ and, for each essential vertex $v \in \text{vert}(P)$, let $f_v : \mathbb{R}^d \rightarrow \mathbb{R}$ be an integer linear functional that gives width one (or zero) to P^v .



We distinguish two cases:


Definition (Boxed vs. spiked)

- ▶ If the set $\{f_v : v \text{ is essential vertex of } P\}$ linearly spans $(\mathbb{R}^d)^*$, then we can find d linearly independent f_v . We call these polytopes **boxed**, because *most of their lattice points lie in the vertices of a d -parallelepiped*.
- ▶ If the set $\{f_v : v \text{ is essential vertex of } P\}$ does not linearly span $(\mathbb{R}^d)^*$, then there is a projection that respects all f_v . We call these polytopes **spiked**, because *most of their lattice points lie in a lattice segment*.



Quasiminimal polytopes


Let $P \in \mathcal{Q}_d(n)$ and, for each essential vertex $v \in \text{vert}(P)$, let $f_v : \mathbb{R}^d \rightarrow \mathbb{R}$ be an integer linear functional that gives width one (or zero) to P^v .



We distinguish two cases:

Definition (Boxed vs. spiked)

- ▶ If the set $\{f_v : v \text{ is essential vertex of } P\}$ linearly spans $(\mathbb{R}^d)^*$, then we can find d linearly independent f_v . We call these polytopes **boxed**, because *most of their lattice points lie in the vertices of a d -parallelepiped*.
- ▶ If the set $\{f_v : v \text{ is essential vertex of } P\}$ does not linearly span $(\mathbb{R}^d)^*$, then there is a projection that respects all f_v . We call these polytopes **spiked**, because *most of their lattice points lie in a lattice segment*.



Boxed polytopes

It is easy to see that boxed d -polytopes have bounded size: apart from d vertices, the only possible lattice points are the 2^d vertices of *the* d -parallelepiped.

Boxed polytopes

It is easy to see that boxed d -polytopes have bounded size: apart from d vertices, the only possible lattice points are the 2^d vertices of *the* d -parallelepiped.

In dimension three this implies that there are finitely many.

We have enumerated those of dimension 3 with computer help. Let the list of them, for each size $n \in \{7, \dots, 11\}$, be denoted $\text{Boxed}_3(n)$.

Spiked polytopes


Spiked d -polytopes can, however, have arbitrarily large size. Still, it turns out that the projection of spiked d -polytopes in the direction of *the* lattice segment has very specific properties:

Spiked polytopes

Spiked d -polytopes can, however, have arbitrarily large size. Still, it turns out that the projection of spiked d -polytopes in the direction of *the* lattice segment has very specific properties:

Theorem (Blanco and Santos)

Every spiked 3-polytope of size $n \geq 7$ projects to one of the following 2-polytopes in such a way that all the vertices in the projection have a unique element in the preimage.



This allows us to explicitly list spiked 3-polytopes for each given size $n \geq 7$. We denote this list by $\text{Spiked}_3(n)$.

Quasiminimal polytopes

Putting these things together, we present the full classification of quasiminimal 3-polytopes:

Theorem (Blanco and Santos)

For $7 \leq n \leq 11$, $\mathcal{Q}_3(n) = \text{Boxed}_3(n) \cup \text{Spiked}_3(n)$, and it has 50, 42, 44, 46 and 49 elements, respectively.

For $n > 11$, $\mathcal{Q}_3(n) = \text{Spiked}_3(n)$ and it has $4n + 7$ elements if $n \equiv 0 \pmod{3}$, and $4n + 5$ otherwise.

SUMMARY & RESULTS

- ▶ $\mathcal{P}_3^*(5)$ and $\mathcal{P}_3^*(6)$ (classified by B-Santos).
- ▶ $\mathcal{Q}_3(n)$, for $n \geq 7$, can be computed explicitly.
- ▶ $\mathcal{M}_3(n) = \text{Merging}(\mathcal{P}_3^*(n-1))$, for $n \geq 7$.

SUMMARY & RESULTS

- ▶ $\mathcal{P}_3^*(5)$ and $\mathcal{P}_3^*(6)$ (classified by B-Santos).
- ▶ $\mathcal{Q}_3(n)$, for $n \geq 7$, can be computed explicitly.
- ▶ $\mathcal{M}_3(n) = \text{Merging}(\mathcal{P}_3^*(n-1))$, for $n \geq 7$.

Assume $\mathcal{P}_3^*(n-1)$ previously computed.

$$\mathcal{P}_3^*(n) = \text{Merging}(\mathcal{P}_3^*(n-1)) \cup \mathcal{Q}_3(n).$$

SUMMARY & RESULTS

- ▶ $\mathcal{P}_3^*(5)$ and $\mathcal{P}_3^*(6)$ (classified by B-Santos).
- ▶ $\mathcal{Q}_3(n)$, for $n \geq 7$, can be computed explicitly.
- ▶ $\mathcal{M}_3(n) = \text{Merging}(\mathcal{P}_3^*(n-1))$, for $n \geq 7$.

Assume $\mathcal{P}_3^*(n-1)$ previously computed.

$$\mathcal{P}_3^*(n) = \text{Merging}(\mathcal{P}_3^*(n-1)) \cup \mathcal{Q}_3(n).$$

Size	4	5	6	7	8	9	10	11
width 2	0	9	74	477	2524	10862	40885	137803
width 3	0	0	2	19	151	836	4148	18635
width 4	0	0	0	0	0	0	2	26
quasiminimal	0	9	35	50	42	44	46	49
merged	0	0	40	446	2633	11654	44989	156415
exceptions	0	0	1	0	0	0	0	0
total	0	9	76	496	2675	11698	45035	156464

Thank you for your patience!!