

Weight classification by Body Mass Index (BMI) (variable calculated from one or more BRFSS questions) (Age-adjusted Prevalence)**View by: Overall****Response: All**

Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
--------------------------------	------------------------------------	---------------------------------------	-------------------------------------

Alabama

Percent (%)	36.2	33.4	29.1	1.2
95% CI	34.5 - 37.9	31.7 - 35.1	27.4 - 30.8	0.9 - 1.6
n	2260	2152	1748	85

Birmingham-Hoover, AL Metropolitan Statistical Area

Percent (%)	37.8	29.5	31.3	*
95% CI	34.0 - 41.5	26.1 - 32.9	27.6 - 35.0	
n	438	391	348	

Tuscaloosa, AL Metropolitan Statistical Area

Percent (%)	36.9	31.6	30.4	*
95% CI	31.4 - 42.4	26.2 - 37.0	24.8 - 36.0	
n	192	155	137	

Alaska

Percent (%)	34.1	32.5	31.9	1.5
95% CI	31.2 - 37.0	29.7 - 35.3	29.0 - 34.8	0.8 - 2.2
n	1027	1063	909	32

Anchorage, AK Metropolitan Statistical Area

Percent (%)	36.3	28.1	34.0	*
95% CI	31.8 - 40.7	24.1 - 32.0	29.5 - 38.4	
n	339	309	303	

Arizona

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	29.5	34.7	33.3	2.4
95% CI	28.5 - 30.6	33.6 - 35.8	32.2 - 34.4	2.0 - 2.8
n	4031	5188	4662	283

Phoenix-Mesa-Scottsdale, AZ Metropolitan Statistical Area

Percent (%)	29.3	34.8	33.7	2.3
95% CI	28.0 - 30.6	33.4 - 36.1	32.3 - 35.1	1.8 - 2.8
n	2383	3068	2726	155

Arkansas

Percent (%)	34.9	35.1	28.3	1.7
95% CI	32.2 - 37.6	32.3 - 37.8	25.7 - 30.9	1.1 - 2.4
n	1627	1739	1423	95

Fayetteville-Springdale-Rogers, AR-MO Metropolitan Statistical Area

Percent (%)	32.4	34.8	32.2	*
95% CI	25.7 - 39.1	28.1 - 41.6	25.3 - 39.2	
n	211	273	243	

Little Rock-North Little Rock-Conway, AR Metropolitan Statistical Area

Percent (%)	35.4	30.7	31.6	*
95% CI	30.2 - 40.7	25.9 - 35.4	26.2 - 36.9	
n	355	395	349	

California

Percent (%)	24.7	35.7	37.1	2.5
95% CI	23.4 - 26.0	34.3 - 37.1	35.6 - 38.5	1.9 - 3.1
n	2118	3063	3133	161

Los Angeles-Long Beach-Anaheim, CA Metropolitan Statistical Area

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
--	--------------------------------	------------------------------------	---------------------------------------	-------------------------------------

Percent (%)	23.6	35.6	37.8	3.0
95% CI	21.0 - 26.3	32.7 - 38.4	34.7 - 40.8	1.6 - 4.4
n	418	716	725	37

Oakland-Hayward-Berkeley, CA Metropolitan Division

Percent (%)	20.4	32.7	44.7	2.2
95% CI	16.5 - 24.2	27.9 - 37.5	39.6 - 49.8	1.0 - 3.5
n	141	202	273	16

Riverside-San Bernardino-Ontario, CA Metropolitan Statistical Area

Percent (%)	31.6	36.0	31.5	*
95% CI	27.7 - 35.6	32.0 - 39.9	27.8 - 35.2	
n	273	328	306	

Sacramento--Roseville--Arden-Arcade, CA Metropolitan Statistical Area

Percent (%)	27.9	35.6	34.3	*
95% CI	24.0 - 31.8	31.4 - 39.8	30.2 - 38.5	
n	240	315	294	

Colorado

Percent (%)	22.4	35.8	39.5	2.3
95% CI	21.3 - 23.5	34.5 - 37.0	38.2 - 40.8	1.9 - 2.8
n	2070	3348	3463	182

Colorado Springs, CO Metropolitan Statistical Area

Percent (%)	22.7	37.4	37.5	*
95% CI	19.7 - 25.7	33.9 - 40.9	33.9 - 41.1	
n	251	436	394	

Denver-Aurora-Lakewood, CO Metropolitan Statistical Area

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
Percent (%)	21.7	37.1	39.2	2.1
95% CI	20.2 - 23.3	35.3 - 38.8	37.3 - 41.0	1.5 - 2.6
n	962	1628	1642	82
Connecticut				
Percent (%)	26.4	35.6	36.3	1.8
95% CI	25.0 - 27.7	34.1 - 37.0	34.8 - 37.9	1.4 - 2.2
n	2555	3582	3203	165
Hartford-West Hartford-East Hartford, CT Metropolitan Statistical Area				
Percent (%)	26.4	35.5	36.1	2.0
95% CI	24.1 - 28.6	33.0 - 37.9	33.5 - 38.7	1.3 - 2.8
n	909	1261	1076	54
Delaware				
Percent (%)	31.0	36.4	30.5	2.2
95% CI	28.7 - 33.2	33.9 - 38.8	28.1 - 32.8	1.4 - 3.0
n	1224	1329	1039	56
District of Columbia				
Percent (%)	23.3	31.4	43.0	2.4
95% CI	21.5 - 25.0	29.4 - 33.4	40.8 - 45.2	1.7 - 3.1
n	985	1140	1320	77
Washington-Arlington-Alexandria, DC-VA-MD-WV Metropolitan Division				
Percent (%)	27.3	35.4	34.8	2.5
95% CI	25.6 - 29.0	33.4 - 37.3	32.9 - 36.8	1.7 - 3.4
n	2258	2702	2630	141
Florida				

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	28.2	34.9	34.7	2.1
95% CI	26.6 - 29.9	33.2 - 36.6	33.0 - 36.5	1.6 - 2.6
n	6150	6976	6381	406

Crestview-Fort Walton Beach-Destin, FL Metropolitan Statistical Area

Percent (%)	27.1	38.4	32.5	*
95% CI	21.0 - 33.1	31.1 - 45.6	25.7 - 39.3	
n	147	197	166	

Deltona-Daytona Beach-Ormond Beach, FL Metropolitan Statistical Area

Percent (%)	31.3	30.5	33.3	*
95% CI	25.1 - 37.4	24.1 - 36.8	27.1 - 39.5	
n	194	228	238	

Gainesville, FL Metropolitan Statistical Area

Percent (%)	31.5	27.4	37.7	*
95% CI	24.1 - 38.9	20.2 - 34.5	30.0 - 45.3	
n	162	143	163	

Jacksonville, FL Metropolitan Statistical Area

Percent (%)	33.1	33.1	31.2	*
95% CI	27.7 - 38.4	28.2 - 38.0	26.3 - 36.2	
n	508	571	539	

Miami-Fort Lauderdale-West Palm Beach, FL Metropolitan Statistical Area

Percent (%)	27.6	33.1	37.1	*
95% CI	23.0 - 32.1	28.9 - 37.4	32.5 - 41.7	
n	284	426	475	

North Port-Sarasota-Bradenton, FL Metropolitan Statistical Area

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	24.4	37.1	37.2	*
95% CI	19.1 - 29.7	30.9 - 43.3	30.9 - 43.5	
n	170	270	250	

Orlando-Kissimmee-Sanford, FL Metropolitan Statistical Area

Percent (%)	25.8	38.9	32.1	3.1
95% CI	22.1 - 29.5	34.4 - 43.5	27.9 - 36.4	1.5 - 4.7
n	361	471	430	28

Panama City, FL Metropolitan Statistical Area

Percent (%)	26.3	36.1	33.5	*
95% CI	20.3 - 32.2	29.4 - 42.8	26.4 - 40.6	
n	197	264	220	

Pensacola-Ferry Pass-Brent, FL Metropolitan Statistical Area

Percent (%)	28.9	39.0	29.5	*
95% CI	23.1 - 34.8	32.5 - 45.5	23.7 - 35.3	
n	192	227	207	

Port St. Lucie, FL Metropolitan Statistical Area

Percent (%)	23.4	38.9	37.2	*
95% CI	18.4 - 28.3	31.4 - 46.3	29.6 - 44.8	
n	155	218	215	

Tallahassee, FL Metropolitan Statistical Area

Percent (%)	32.2	25.3	38.7	*
95% CI	26.1 - 38.3	20.0 - 30.5	31.9 - 45.5	
n	284	270	277	

Tampa-St. Petersburg-Clearwater, FL Metropolitan Statistical Area

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)	
Percent (%)	24.2	36.5	38.1		*
95% CI	20.7 - 27.8	32.0 - 41.0	33.4 - 42.7		
n	397	496	483		
Georgia					
Percent (%)	31.3	33.3	33.5		1.9
95% CI	29.6 - 32.9	31.6 - 35.0	31.8 - 35.2		1.4 - 2.5
n	1724	1860	1678		107
Atlanta-Sandy Springs-Roswell, GA Metropolitan Statistical Area					
Percent (%)	29.4	33.3	35.5		1.8
95% CI	27.1 - 31.7	31.0 - 35.6	33.2 - 37.9		1.1 - 2.5
n	767	904	879		47
Augusta-Richmond County, GA-SC Metropolitan Statistical Area					
Percent (%)	37.7	32.4	28.2		*
95% CI	31.9 - 43.4	26.8 - 38.0	22.8 - 33.6		
n	280	287	198		
Hawaii					
Percent (%)	24.1	34.8	38.7		2.4
95% CI	22.6 - 25.5	33.3 - 36.4	37.0 - 40.3		1.9 - 3.0
n	1739	2514	2886		180
Kahului-Wailuku-Lahaina, HI Metropolitan Statistical Area					
Percent (%)	24.0	35.0	38.6		2.4
95% CI	20.9 - 27.1	31.4 - 38.6	35.0 - 42.3		1.4 - 3.4
n	310	437	504		36
Idaho					

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
--	--------------------------------	------------------------------------	---------------------------------------	-------------------------------------

Percent (%)	29.1	36.1	33.3	1.4
95% CI	27.3 - 31.0	34.1 - 38.2	31.2 - 35.3	1.0 - 1.9
n	1360	1729	1458	76

Boise City, ID Metropolitan Statistical Area

Percent (%)	29.6	35.9	33.1	1.4
95% CI	26.1 - 33.1	32.2 - 39.7	29.3 - 36.9	0.6 - 2.2
n	392	433	377	18

Illinois

Percent (%)	30.5	34.2	33.5	1.8
95% CI	28.8 - 32.1	32.5 - 36.0	31.7 - 35.3	1.2 - 2.4
n	1633	1897	1645	72

Chicago-Naperville-Elgin, IL-IN-WI Metropolitan Statistical Area

Percent (%)	29.5	33.8	34.9	1.8
95% CI	27.7 - 31.4	32.0 - 35.7	33.0 - 36.9	1.2 - 2.4
n	1485	1739	1508	68

Indiana

Percent (%)	33.1	33.9	31.1	1.9
95% CI	32.0 - 34.3	32.7 - 35.0	30.0 - 32.3	1.5 - 2.3
n	4434	4639	3694	206

Evansville, IN-KY Metropolitan Statistical Area

Percent (%)	30.4	33.0	35.2	*
95% CI	25.7 - 35.0	28.1 - 38.0	30.0 - 40.4	
n	202	202	187	

Fort Wayne, IN Metropolitan Statistical Area

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
--	--------------------------------	------------------------------------	---------------------------------------	-------------------------------------

Percent (%)	33.4	30.0	34.3	*
95% CI	29.1 - 37.8	25.9 - 34.1	29.7 - 38.9	
n	269	275	249	

Indianapolis-Carmel-Anderson, IN Metropolitan Statistical Area

Percent (%)	31.5	33.8	32.3	2.4
95% CI	29.5 - 33.4	31.7 - 35.9	30.2 - 34.4	1.6 - 3.3
n	1310	1401	1164	65

South Bend-Mishawaka, IN-MI Metropolitan Statistical Area

Percent (%)	33.2	35.9	28.6	*
95% CI	27.0 - 39.5	29.6 - 42.3	22.5 - 34.8	
n	174	199	143	

Iowa

Percent (%)	36.1	33.2	29.3	1.4
95% CI	34.7 - 37.5	31.8 - 34.5	27.9 - 30.7	0.9 - 1.8
n	2603	2511	1920	79

Cedar Rapids, IA Metropolitan Statistical Area

Percent (%)	34.5	35.0	29.7	*
95% CI	29.9 - 39.2	30.3 - 39.6	25.2 - 34.1	
n	185	199	167	

Des Moines-West Des Moines, IA Metropolitan Statistical Area

Percent (%)	34.0	34.4	29.9	1.7
95% CI	30.9 - 37.1	31.2 - 37.5	26.8 - 33.1	0.7 - 2.6
n	448	444	358	17

Omaha-Council Bluffs, NE-IA Metropolitan Statistical Area

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
Percent (%)	33.0	35.0	30.0	2.0
95% CI	30.7 - 35.2	32.7 - 37.3	27.7 - 32.2	1.2 - 2.9
n	1061	1114	872	46
Sioux City, IA-NE-SD Metropolitan Statistical Area				
Percent (%)	29.4	40.4	28.4	*
95% CI	24.3 - 34.5	34.9 - 46.0	22.9 - 33.8	
n	319	365	256	
Kansas				
Percent (%)	32.2	34.5	31.3	1.9
95% CI	31.3 - 33.1	33.6 - 35.4	30.5 - 32.2	1.7 - 2.2
n	6551	7155	5732	345
Kansas City, MO-KS Metropolitan Statistical Area				
Percent (%)	30.7	35.9	32.0	1.3
95% CI	28.8 - 32.7	33.9 - 38.0	30.1 - 34.0	0.8 - 1.7
n	2198	2567	2203	104
Manhattan, KS Metropolitan Statistical Area				
Percent (%)	25.5	34.5	37.3	2.8
95% CI	22.1 - 28.9	30.5 - 38.4	33.0 - 41.5	1.4 - 4.1
n	226	296	243	18
Salina, KS Micropolitan Statistical Area				
Percent (%)	33.3	33.6	31.4	*
95% CI	28.4 - 38.2	28.8 - 38.3	26.5 - 36.3	
n	196	206	156	
Topeka, KS Metropolitan Statistical Area				

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	37.3	31.2	30.0	1.5
95% CI	34.0 - 40.7	28.0 - 34.4	26.7 - 33.3	0.6 - 2.3
n	555	491	425	25

Wichita, KS Metropolitan Statistical Area

Percent (%)	34.0	34.9	29.7	1.5
95% CI	31.9 - 36.0	32.8 - 36.9	27.7 - 31.6	1.0 - 2.0
n	1200	1328	1069	56

Kentucky

Percent (%)	34.1	33.1	31.3	1.5
95% CI	32.3 - 35.9	31.3 - 34.8	29.5 - 33.1	1.1 - 2.0
n	2923	2782	2184	129

Huntington-Ashland, WV-KY-OH Metropolitan Statistical Area

Percent (%)	38.3	35.8	24.5	*
95% CI	34.2 - 42.4	31.5 - 40.1	21.0 - 28.1	
n	405	384	278	

Lexington-Fayette, KY Metropolitan Statistical Area

Percent (%)	31.5	32.7	34.6	*
95% CI	25.9 - 37.1	27.3 - 38.1	29.1 - 40.1	
n	138	162	173	

Louisville/Jefferson County, KY-IN Metropolitan Statistical Area

Percent (%)	29.8	34.2	34.5	*
95% CI	25.7 - 33.8	30.0 - 38.4	30.1 - 38.9	
n	669	639	542	

Louisiana

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	35.9	33.6	28.5	2.0
95% CI	34.0 - 37.7	31.8 - 35.5	26.7 - 30.4	1.5 - 2.5
n	1622	1532	1201	84

Baton Rouge, LA Metropolitan Statistical Area

Percent (%)	32.9	34.3	30.3	*
95% CI	28.7 - 37.0	30.0 - 38.6	25.8 - 34.7	
n	249	251	205	

New Orleans-Metairie, LA Metropolitan Statistical Area

Percent (%)	35.4	34.8	27.8	2.0
95% CI	31.1 - 39.8	30.3 - 39.2	23.7 - 32.0	0.9 - 3.1
n	275	289	231	19

Maine

Percent (%)	28.3	35.1	34.9	1.6
95% CI	26.7 - 29.9	33.4 - 36.9	33.1 - 36.7	1.1 - 2.1
n	2745	3387	3026	137

Portland-South Portland, ME Metropolitan Statistical Area

Percent (%)	23.7	37.1	37.4	1.7
95% CI	20.9 - 26.6	33.9 - 40.3	34.2 - 40.6	0.9 - 2.6
n	600	942	887	40

Maryland

Percent (%)	30.8	34.6	32.7	1.9
95% CI	29.4 - 32.2	33.1 - 36.1	31.3 - 34.2	1.4 - 2.4
n	3964	4452	3784	186

Baltimore-Columbia-Towson, MD Metropolitan Statistical Area

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
--	--------------------------------	------------------------------------	---------------------------------------	-------------------------------------

Percent (%)	30.5	35.0	32.8	1.7
95% CI	28.4 - 32.6	32.7 - 37.3	30.5 - 35.0	1.0 - 2.4
n	1455	1661	1444	68

Hagerstown-Martinsburg, MD-WV Metropolitan Statistical Area

Percent (%)	37.9	33.8	27.0	*
95% CI	31.9 - 43.8	28.0 - 39.5	21.3 - 32.6	
n	266	254	199	

Salisbury, MD-DE Metropolitan Statistical Area

Percent (%)	31.7	36.2	30.4	*
95% CI	28.2 - 35.2	32.3 - 40.1	26.4 - 34.5	
n	636	727	516	

Silver Spring-Frederick-Rockville, MD Metropolitan Division

Percent (%)	20.7	37.2	39.8	2.3
95% CI	18.1 - 23.3	33.9 - 40.5	36.5 - 43.1	1.0 - 3.5
n	532	802	840	40

Massachusetts

Percent (%)	25.3	34.6	37.7	2.5
95% CI	23.5 - 27.1	32.6 - 36.5	35.6 - 39.7	1.5 - 3.4
n	1591	2298	2269	114

Boston, MA Metropolitan Division

Percent (%)	25.2	34.3	37.3	*
95% CI	21.7 - 28.7	30.6 - 38.0	33.5 - 41.1	
n	446	631	645	

Cambridge-Newton-Framingham, MA Metropolitan Division

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	23.1	34.2	40.8	*
95% CI	20.2 - 25.9	31.0 - 37.3	37.4 - 44.1	
n	505	754	833	

Springfield, MA Metropolitan Statistical Area

Percent (%)	23.9	30.3	42.9	*
95% CI	18.5 - 29.3	24.1 - 36.5	36.1 - 49.6	
n	152	195	209	

Worcester, MA-CT Metropolitan Statistical Area

Percent (%)	27.0	37.4	33.0	*
95% CI	22.6 - 31.5	32.6 - 42.3	28.0 - 38.0	
n	374	497	391	

Michigan

Percent (%)	31.6	34.3	31.8	2.3
95% CI	30.4 - 32.8	33.0 - 35.5	30.5 - 33.1	1.9 - 2.8
n	3275	3649	3042	163

Grand Rapids-Wyoming, MI Metropolitan Statistical Area

Percent (%)	31.8	31.9	35.6	*
95% CI	27.9 - 35.7	28.1 - 35.7	31.6 - 39.6	
n	288	312	309	

Lansing-East Lansing, MI Metropolitan Statistical Area

Percent (%)	31.5	32.3	34.3	*
95% CI	26.2 - 36.8	26.8 - 37.8	28.9 - 39.8	
n	144	164	158	

Warren-Troy-Farmington Hills, MI Metropolitan Division

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
Percent (%)	27.9	37.3	32.1	2.7
95% CI	25.3 - 30.5	34.6 - 40.0	29.5 - 34.8	1.5 - 3.8
n	664	923	737	40
Minnesota				
Percent (%)	28.0	36.0	34.2	1.8
95% CI	27.0 - 28.9	34.9 - 37.0	33.2 - 35.3	1.5 - 2.2
n	4622	5753	4836	234
Duluth, MN-WI Metropolitan Statistical Area				
Percent (%)	26.3	41.7	31.2	*
95% CI	21.6 - 31.0	36.8 - 46.6	26.9 - 35.6	
n	274	391	315	
Minneapolis-St. Paul-Bloomington, MN-WI Metropolitan Statistical Area				
Percent (%)	25.6	35.1	37.4	1.9
95% CI	24.4 - 26.9	33.7 - 36.5	35.9 - 38.9	1.4 - 2.3
n	1992	2673	2559	111
Rochester, MN Metropolitan Statistical Area				
Percent (%)	29.6	37.5	31.8	*
95% CI	25.2 - 34.0	32.1 - 42.8	26.8 - 36.7	
n	203	223	184	
St. Cloud, MN Metropolitan Statistical Area				
Percent (%)	29.5	37.4	30.2	*
95% CI	24.6 - 34.4	32.8 - 42.0	25.4 - 35.0	
n	191	246	185	
Mississippi				

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	37.2	32.3	28.1	2.5
95% CI	35.0 - 39.3	30.2 - 34.4	25.9 - 30.2	1.7 - 3.2
n	1718	1651	1284	97

Jackson, MS Metropolitan Statistical Area

Percent (%)	34.9	34.1	28.4	*
95% CI	28.9 - 40.8	28.1 - 40.1	22.6 - 34.1	
n	174	173	149	

Missouri

Percent (%)	32.1	34.8	30.8	2.4
95% CI	30.4 - 33.7	33.1 - 36.5	29.1 - 32.4	1.7 - 3.0
n	2317	2494	2077	122

St. Louis, MO-IL Metropolitan Statistical Area

Percent (%)	30.6	34.7	32.6	2.0
95% CI	27.5 - 33.8	31.5 - 37.9	29.3 - 35.9	1.1 - 3.0
n	638	750	659	32

Montana

Percent (%)	24.9	36.1	37.1	1.8
95% CI	23.3 - 26.6	34.3 - 38.0	35.2 - 39.1	1.3 - 2.3
n	1575	2071	1778	88

Billings, MT Metropolitan Statistical Area

Percent (%)	26.0	38.8	33.1	*
95% CI	21.7 - 30.4	34.1 - 43.5	28.5 - 37.8	
n	140	223	195	

Nebraska

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
Percent (%)	32.4	36.0	29.7	1.9
95% CI	31.1 - 33.6	34.7 - 37.3	28.5 - 31.0	1.4 - 2.4
n	4963	5272	3916	179
Grand Island, NE Metropolitan Statistical Area				
Percent (%)	35.5	34.4	29.3	*
95% CI	31.1 - 39.9	29.8 - 39.0	24.7 - 34.0	
n	289	247	191	
Lincoln, NE Metropolitan Statistical Area				
Percent (%)	28.9	38.0	30.8	2.3
95% CI	26.0 - 31.8	34.8 - 41.2	27.7 - 33.8	1.1 - 3.6
n	422	545	451	22
North Platte, NE Micropolitan Statistical Area				
Percent (%)	34.1	36.9	27.3	*
95% CI	28.8 - 39.4	31.4 - 42.4	22.1 - 32.5	
n	194	196	138	
Scottsbluff, NE Micropolitan Statistical Area				
Percent (%)	35.7	34.3	28.9	*
95% CI	30.7 - 40.8	29.1 - 39.4	23.8 - 34.0	
n	203	216	157	
Nevada				
Percent (%)	26.5	38.8	32.7	2.0
95% CI	24.2 - 28.8	36.2 - 41.4	30.2 - 35.2	1.3 - 2.7
n	901	1335	1191	69
Reno, NV Metropolitan Statistical Area				

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
Percent (%)	23.7	37.0	36.6	2.7
95% CI	20.5 - 26.9	33.5 - 40.5	33.0 - 40.3	1.3 - 4.1
n	270	444	433	27
New Hampshire				
Percent (%)	27.6	36.3	34.6	1.5
95% CI	25.5 - 29.6	34.0 - 38.7	32.3 - 36.9	0.9 - 2.1
n	1492	1892	1737	70
Claremont-Lebanon, NH-VT Micropolitan Statistical Area				
Percent (%)	26.8	36.3	34.9	*
95% CI	22.9 - 30.7	31.8 - 40.8	30.5 - 39.3	
n	435	589	562	
Rockingham County-Strafford County, NH Metropolitan Division				
Percent (%)	24.9	38.1	36.0	*
95% CI	21.2 - 28.6	33.8 - 42.4	31.6 - 40.4	
n	340	475	388	
New Jersey				
Percent (%)	26.7	34.4	37.5	1.5
95% CI	25.0 - 28.3	32.8 - 36.1	35.7 - 39.3	1.0 - 1.9
n	3002	3894	3361	137
Camden, NJ Metropolitan Division				
Percent (%)	31.9	34.2	32.4	*
95% CI	27.7 - 36.1	29.9 - 38.5	28.0 - 36.8	
n	483	542	437	
Newark, NJ-PA Metropolitan Division				

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
Percent (%)	24.5	36.9	37.1	1.5
95% CI	21.6 - 27.4	33.6 - 40.1	33.8 - 40.4	0.7 - 2.4
n	834	1249	1123	44
Wilmington, DE-MD-NJ Metropolitan Division				
Percent (%)	31.0	35.4	31.1	2.5
95% CI	28.0 - 34.0	32.2 - 38.5	28.1 - 34.2	1.5 - 3.6
n	731	778	598	46
New Mexico				
Percent (%)	28.6	36.2	33.1	2.1
95% CI	26.9 - 30.3	34.4 - 38.0	31.2 - 35.0	1.5 - 2.7
n	1758	2273	1887	128
Albuquerque, NM Metropolitan Statistical Area				
Percent (%)	26.8	36.6	34.8	1.8
95% CI	23.8 - 29.7	33.4 - 39.9	31.5 - 38.0	0.9 - 2.7
n	412	596	511	30
New York				
Percent (%)	25.2	35.0	37.4	2.4
95% CI	24.0 - 26.3	33.7 - 36.3	36.1 - 38.8	1.9 - 2.9
n	2994	4106	3871	209
Albany-Schenectady-Troy, NY Metropolitan Statistical Area				
Percent (%)	27.8	34.7	35.5	*
95% CI	22.6 - 33.1	29.2 - 40.3	29.4 - 41.6	
n	163	216	197	
Buffalo-Cheektowaga-Niagara Falls, NY Metropolitan Statistical Area				

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	28.4	35.7	33.2	*
95% CI	22.9 - 34.0	29.1 - 42.3	27.0 - 39.4	
n	140	176	157	

Nassau County-Suffolk County, NY Metropolitan Division

Percent (%)	26.2	35.5	37.2	*
95% CI	22.6 - 29.7	31.8 - 39.3	33.3 - 41.0	
n	294	425	385	

New York-Jersey City-White Plains, NY-NJ Metropolitan Division

Percent (%)	23.1	35.0	39.6	2.3
95% CI	21.8 - 24.5	33.6 - 36.5	38.1 - 41.1	1.7 - 2.8
n	1879	2916	2873	135

Rochester, NY Metropolitan Statistical Area

Percent (%)	27.2	31.2	39.9	*
95% CI	21.4 - 33.0	25.0 - 37.3	32.9 - 46.8	
n	144	179	167	

North Carolina

Percent (%)	31.6	34.4	32.2	1.8
95% CI	29.8 - 33.5	32.5 - 36.3	30.3 - 34.1	1.2 - 2.3
n	1442	1602	1369	70

Charlotte-Concord-Gastonia, NC-SC Metropolitan Statistical Area

Percent (%)	33.3	33.5	31.7	*
95% CI	29.2 - 37.3	29.5 - 37.5	27.7 - 35.6	
n	428	477	421	

North Dakota

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	33.1	35.9	29.4	1.6
95% CI	31.5 - 34.7	34.3 - 37.6	27.7 - 31.0	1.1 - 2.1
n	2241	2461	1749	81

Bismarck, ND Metropolitan Statistical Area

Percent (%)	30.2	37.4	30.4	*
95% CI	26.4 - 34.0	33.5 - 41.4	26.5 - 34.2	
n	329	403	295	

Fargo, ND-MN Metropolitan Statistical Area

Percent (%)	29.2	39.9	29.7	*
95% CI	26.0 - 32.5	36.2 - 43.5	26.4 - 33.1	
n	358	503	350	

Grand Forks, ND-MN Metropolitan Statistical Area

Percent (%)	35.4	31.3	31.9	*
95% CI	30.3 - 40.5	26.4 - 36.2	26.8 - 37.0	
n	200	198	160	

Minot, ND Micropolitan Statistical Area

Percent (%)	34.9	35.4	27.8	*
95% CI	29.5 - 40.3	29.9 - 40.8	22.3 - 33.4	
n	217	236	162	

Ohio

Percent (%)	33.4	33.4	31.4	1.8
95% CI	32.0 - 34.8	32.0 - 34.8	30.0 - 32.8	1.3 - 2.2
n	4001	4023	3189	185

Cincinnati, OH-KY-IN Metropolitan Statistical Area

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
Percent (%)	29.7	32.3	35.8	2.3
95% CI	26.6 - 32.8	29.2 - 35.4	32.4 - 39.1	1.2 - 3.4
n	491	573	519	31
Cleveland-Elyria, OH Metropolitan Statistical Area				
Percent (%)	32.4	35.0	30.7	1.8
95% CI	28.3 - 36.5	30.9 - 39.1	26.7 - 34.7	0.8 - 2.9
n	321	388	304	22
Columbus, OH Metropolitan Statistical Area				
Percent (%)	32.1	35.5	30.8	1.6
95% CI	29.3 - 34.9	32.6 - 38.5	27.9 - 33.7	0.9 - 2.3
n	638	679	547	30
Dayton, OH Metropolitan Statistical Area				
Percent (%)	34.8	32.4	31.1	*
95% CI	29.0 - 40.6	26.7 - 38.1	25.5 - 36.7	
n	202	183	158	
Toledo, OH Metropolitan Statistical Area				
Percent (%)	41.5	30.5	26.8	*
95% CI	36.3 - 46.8	25.8 - 35.1	22.2 - 31.5	
n	264	245	189	
Oklahoma				
Percent (%)	36.4	33.8	28.3	1.4
95% CI	34.7 - 38.1	32.2 - 35.5	26.7 - 29.9	1.0 - 1.9
n	2137	2116	1766	94
Oklahoma City, OK Metropolitan Statistical Area				

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
--	--------------------------------	------------------------------------	---------------------------------------	-------------------------------------

Percent (%)	34.3	33.3	30.4	2.0
95% CI	31.5 - 37.2	30.5 - 36.2	27.6 - 33.1	0.9 - 3.0
n	661	681	609	32

Tulsa, OK Metropolitan Statistical Area

Percent (%)	36.0	34.0	29.3	*
95% CI	32.0 - 39.9	30.4 - 37.6	25.6 - 33.1	
n	395	409	342	

Oregon

Percent (%)	28.9	34.9	33.9	2.3
95% CI	27.3 - 30.5	33.2 - 36.6	32.3 - 35.6	1.7 - 3.0
n	1390	1734	1694	91

Portland-Vancouver-Hillsboro, OR-WA Metropolitan Statistical Area

Percent (%)	26.8	33.7	37.1	2.4
95% CI	24.8 - 28.8	31.5 - 35.9	34.8 - 39.4	1.4 - 3.4
n	809	970	1046	49

Pennsylvania

Percent (%)	30.8	35.0	32.0	2.2
95% CI	29.2 - 32.5	33.3 - 36.7	30.3 - 33.6	1.7 - 2.8
n	1950	2177	1904	104

Allentown-Bethlehem-Easton, PA-NJ Metropolitan Statistical Area

Percent (%)	29.8	31.7	37.5	*
95% CI	24.4 - 35.2	25.7 - 37.7	30.7 - 44.3	
n	267	277	243	

Montgomery County-Bucks County-Chester County, PA Metropolitan Division

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
Percent (%)	24.2	36.3	37.1	*
95% CI	19.7 - 28.8	31.2 - 41.4	32.1 - 42.2	
n	142	215	214	
Philadelphia, PA Metropolitan Division				
Percent (%)	31.7	35.5	31.1	1.7
95% CI	27.8 - 35.6	31.5 - 39.5	27.4 - 34.8	0.7 - 2.6
n	270	321	303	16
Pittsburgh, PA Metropolitan Statistical Area				
Percent (%)	30.5	34.8	32.7	2.0
95% CI	27.4 - 33.6	31.5 - 38.1	29.5 - 36.0	1.1 - 3.0
n	417	476	436	24
Rhode Island				
Percent (%)	29.6	34.4	34.4	1.7
95% CI	27.6 - 31.6	32.3 - 36.4	32.2 - 36.5	1.0 - 2.3
n	1528	1892	1643	78
Providence-Warwick, RI-MA Metropolitan Statistical Area				
Percent (%)	31.3	34.5	32.4	1.8
95% CI	28.4 - 34.3	31.6 - 37.3	29.7 - 35.2	0.9 - 2.7
n	1674	2059	1786	85
South Carolina				
Percent (%)	34.1	33.4	30.9	1.6
95% CI	32.7 - 35.5	32.0 - 34.8	29.5 - 32.3	1.2 - 1.9
n	3577	3713	3017	168
Charleston-North Charleston, SC Metropolitan Statistical Area				

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	31.8	34.9	31.8	1.5
95% CI	28.5 - 35.1	31.4 - 38.4	28.4 - 35.2	0.7 - 2.4
n	454	489	425	21

Columbia, SC Metropolitan Statistical Area

Percent (%)	35.5	34.6	29.1	*
95% CI	31.5 - 39.6	30.6 - 38.5	25.1 - 33.0	
n	412	406	320	

Florence, SC Metropolitan Statistical Area

Percent (%)	37.3	25.7	35.6	*
95% CI	31.4 - 43.1	20.4 - 31.1	29.3 - 41.9	
n	180	143	151	

Greenville-Anderson-Mauldin, SC Metropolitan Statistical Area

Percent (%)	31.9	34.0	32.1	2.0
95% CI	28.3 - 35.5	30.2 - 37.7	28.4 - 35.8	0.9 - 3.1
n	436	478	439	26

Hilton Head Island-Bluffton-Beaufort, SC Metropolitan Statistical Area

Percent (%)	24.6	36.7	37.2	*
95% CI	19.3 - 29.9	29.9 - 43.5	30.6 - 43.9	
n	152	198	208	

Myrtle Beach-Conway-North Myrtle Beach, SC-NC Metropolitan Statistical Area

Percent (%)	30.1	36.0	31.9	*
95% CI	25.3 - 34.9	31.1 - 40.9	26.6 - 37.2	
n	267	335	244	

Spartanburg, SC Metropolitan Statistical Area

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)	
Percent (%)	35.6	29.6	33.3		*
95% CI	28.1 - 43.1	24.0 - 35.1	25.4 - 41.3		
n	149	174	132		
South Dakota					
Percent (%)	31.5	35.6	31.4	1.5	
95% CI	29.2 - 33.7	33.2 - 37.9	29.1 - 33.8	0.8 - 2.3	
n	2258	2420	1779	76	
Aberdeen, SD Micropolitan Statistical Area					
Percent (%)	28.3	35.9	33.7		*
95% CI	21.5 - 35.0	27.8 - 44.1	25.8 - 41.5		
n	140	183	135		
Rapid City, SD Metropolitan Statistical Area					
Percent (%)	35.3	36.9	26.7		*
95% CI	29.7 - 40.8	31.3 - 42.5	21.9 - 31.6		
n	325	362	292		
Sioux Falls, SD Metropolitan Statistical Area					
Percent (%)	28.8	35.3	35.4		*
95% CI	23.8 - 33.9	30.2 - 40.4	30.0 - 40.8		
n	330	370	313		
Tennessee					
Percent (%)	32.5	35.3	30.2	2.0	
95% CI	30.7 - 34.4	33.3 - 37.2	28.3 - 32.1	1.4 - 2.5	
n	1810	1890	1551	103	
Kingsport-Bristol-Bristol, TN-VA Metropolitan Statistical Area					

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	29.4	35.4	32.8	*
95% CI	23.8 - 35.0	29.0 - 41.7	26.3 - 39.3	
n	171	216	155	

Knoxville, TN Metropolitan Statistical Area

Percent (%)	31.3	31.4	35.3	*
95% CI	25.8 - 36.8	26.3 - 36.6	29.6 - 41.1	
n	156	198	180	

Memphis, TN-MS-AR Metropolitan Statistical Area

Percent (%)	37.0	35.1	26.1	*
95% CI	32.2 - 41.7	30.5 - 39.7	22.0 - 30.3	
n	317	290	245	

Nashville-Davidson--Murfreesboro--Franklin, TN Metropolitan Statistical Area

Percent (%)	27.6	37.3	33.7	*
95% CI	23.9 - 31.3	32.9 - 41.7	29.3 - 38.1	
n	255	320	253	

Texas

Percent (%)	32.8	36.3	29.1	1.8
95% CI	30.9 - 34.6	34.4 - 38.2	27.3 - 30.9	1.2 - 2.3
n	3761	3960	3323	194

Austin-Round Rock, TX Metropolitan Statistical Area

Percent (%)	29.0	32.9	37.2	*
95% CI	25.1 - 32.9	28.8 - 37.0	33.0 - 41.4	
n	403	525	568	

College Station-Bryan, TX Metropolitan Statistical Area

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	32.9	33.4	32.7	*
95% CI	24.1 - 41.7	23.4 - 43.5	23.1 - 42.3	
n	142	168	172	

Corpus Christi, TX Metropolitan Statistical Area

Percent (%)	37.0	33.2	27.1	*
95% CI	30.3 - 43.7	25.8 - 40.5	20.4 - 33.8	
n	225	238	201	

Dallas-Plano-Irving, TX Metropolitan Division

Percent (%)	28.2	37.0	33.2	*
95% CI	23.4 - 32.9	31.7 - 42.3	28.2 - 38.3	
n	342	407	336	

El Paso, TX Metropolitan Statistical Area

Percent (%)	34.4	33.8	30.1	*
95% CI	28.8 - 40.1	28.3 - 39.2	24.4 - 35.8	
n	165	171	134	

Fort Worth-Arlington, TX Metropolitan Division

Percent (%)	36.7	33.7	28.0	*
95% CI	31.0 - 42.3	28.3 - 39.2	22.5 - 33.5	
n	179	197	136	

Houston-The Woodlands-Sugar Land, TX Metropolitan Statistical Area

Percent (%)	30.5	42.7	24.9	*
95% CI	26.2 - 34.7	38.0 - 47.4	20.9 - 28.8	
n	679	744	622	

San Antonio-New Braunfels, TX Metropolitan Statistical Area

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
--	--------------------------------	------------------------------------	---------------------------------------	-------------------------------------

Percent (%)	30.6	37.5	30.4	*
95% CI	24.1 - 37.0	30.4 - 44.5	23.6 - 37.1	
n	185	205	186	

Wichita Falls, TX Metropolitan Statistical Area

Percent (%)	43.2	28.3	20.6	*
95% CI	32.4 - 54.0	17.5 - 39.0	13.0 - 28.3	
n	173	147	133	

Utah

Percent (%)	25.5	35.5	36.9	2.2
95% CI	24.4 - 26.6	34.3 - 36.7	35.6 - 38.1	1.7 - 2.6
n	2517	3384	3248	176

Ogden-Clearfield, UT Metropolitan Statistical Area

Percent (%)	29.0	35.4	33.6	1.9
95% CI	26.5 - 31.5	32.8 - 38.1	31.0 - 36.2	1.1 - 2.7
n	474	591	540	29

Provo-Orem, UT Metropolitan Statistical Area

Percent (%)	24.1	35.5	38.4	2.1
95% CI	21.7 - 26.4	32.8 - 38.2	35.6 - 41.1	1.1 - 3.1
n	403	590	599	28

Salt Lake City, UT Metropolitan Statistical Area

Percent (%)	25.0	35.0	37.6	2.4
95% CI	23.2 - 26.8	32.9 - 37.1	35.5 - 39.8	1.7 - 3.1
n	840	1067	1047	59

Vermont

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	27.0	33.9	37.6	1.5
95% CI	25.2 - 28.7	32.1 - 35.7	35.6 - 39.6	1.1 - 2.0
n	1615	2188	2076	101

Burlington-South Burlington, VT Metropolitan Statistical Area

Percent (%)	27.5	31.9	39.5	1.2
95% CI	24.0 - 30.9	28.7 - 35.1	35.8 - 43.2	0.5 - 1.8
n	406	509	501	23

Virginia

Percent (%)	29.6	35.8	32.7	1.8
95% CI	28.2 - 31.1	34.3 - 37.4	31.2 - 34.2	1.3 - 2.3
n	2744	3205	2724	136

Richmond, VA Metropolitan Statistical Area

Percent (%)	30.9	32.6	35.6	*
95% CI	27.2 - 34.5	28.5 - 36.7	31.4 - 39.7	
n	466	489	428	

Virginia Beach-Norfolk-Newport News, VA-NC Metropolitan Statistical Area

Percent (%)	32.4	37.0	29.6	1.1
95% CI	29.4 - 35.4	33.8 - 40.2	26.5 - 32.6	0.6 - 1.6
n	632	699	560	33

Washington

Percent (%)	27.4	34.1	36.5	2.0
95% CI	26.3 - 28.5	33.0 - 35.3	35.3 - 37.7	1.6 - 2.4
n	3422	4311	4126	203

Seattle-Bellevue-Everett, WA Metropolitan Division

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0-29.9)	Normal Weight (BMI 18.5-24.9)	Underweight (BMI 12.0-18.4)
--	--------------------------------	-----------------------------------	--------------------------------------	------------------------------------

Percent (%)	22.9	33.2	41.4	2.4
95% CI	21.3 - 24.5	31.4 - 35.0	39.5 - 43.3	1.8 - 3.1
n	875	1347	1515	79

Spokane-Spokane Valley, WA Metropolitan Statistical Area

Percent (%)	28.0	33.6	36.1	*
95% CI	24.0 - 32.0	29.5 - 37.7	31.7 - 40.5	
n	296	373	317	

West Virginia

Percent (%)	37.7	32.7	28.3	1.3
95% CI	35.9 - 39.6	30.9 - 34.5	26.5 - 30.1	0.9 - 1.7
n	1980	1786	1279	71

Charleston, WV Metropolitan Statistical Area

Percent (%)	39.1	34.6	25.2	*
95% CI	34.3 - 43.9	29.8 - 39.4	20.6 - 29.8	
n	305	267	170	

Wisconsin

Percent (%)	31.6	34.6	32.5	1.3
95% CI	29.7 - 33.5	32.7 - 36.4	30.6 - 34.5	0.8 - 1.8
n	1706	1973	1600	69

Milwaukee-Waukesha-West Allis, WI Metropolitan Statistical Area

Percent (%)	28.7	38.0	32.3	*
95% CI	24.8 - 32.6	33.9 - 42.1	28.2 - 36.3	
n	308	379	331	

Wyoming

	Obese (BMI 30.0 - 99.8)	Overweight (BMI 25.0- 29.9)	Normal Weight (BMI 18.5- 24.9)	Underweight (BMI 12.0- 18.4)
--	--------------------------------	------------------------------------	---------------------------------------	-------------------------------------

Percent (%)	28.3	35.5	34.1	2.0
95% CI	26.5 - 30.1	33.5 - 37.5	32.1 - 36.1	1.4 - 2.7
n	1223	1511	1350	74

Guam

Percent (%)	33.2	32.9	31.3	2.6
95% CI	30.1 - 36.3	29.8 - 36.1	28.1 - 34.4	1.6 - 3.7
n	487	467	438	35

Puerto Rico

Percent (%)	32.6	34.8	31.0	1.6
95% CI	30.6 - 34.6	32.8 - 36.8	29.1 - 32.9	1.1 - 2.0
n	1299	1609	1293	63

San Juan-Carolina-Caguas, PR Metropolitan Statistical Area

Percent (%)	32.1	34.7	31.6	1.5
95% CI	29.7 - 34.5	32.2 - 37.2	29.2 - 34.0	1.0 - 2.1
n	811	1007	827	40

Footnote

* Age-adjusted prevalence estimate not available if the unweighted sample size for the denominator was < 50 or the Relative Standard Error (RSE) is > 0.3 or if the state did not collect data for that calendar year.

Data Source: Behavioral Risk Factor Surveillance System (BRFSS)

Note

Use caution in interpreting cell sizes less than 50.

If you see that a question is not available for a particular year, it is because the question was dropped or changed. Check the category of interest for that year to find similar items.

Denominator includes all respondents except those with missing, don't know, and refused answers.

** Median value reported with no confidence intervals.