

Predicting pitcher ERAs in a given year using conventional stats

Evan Chung, Krystal Liu

December 13th, 2025

Executive Summary

- **Purpose**

- This project is exploring which conventional pitching statistic can predict a pitcher's ERA in a given year and develops a statistical model predicting ERA, while discussing if using conventional stats is a good fit.

- **Data and Methods**

- The data set for this project was obtained from kaggle and has data from 2015-2024.
- Predictors examined: WHIP (Walks and Hits per Inning Pitched), HR.9 (Home Runs allowed per 9 Innings Pitched), K.9 (Strikeouts per 9 innings pitched), age, team, and season fixed effects.
- A fixed effects model was used to control for season-to-season variation.
- The model performance is evaluated by R-Squared, significance testing, and through comparison

- **Key Findings**

- WHIP is the strongest predictor of ERA, while HR.9, K.9, and team all have a statistically and practically significant impact on ERA as well.
- Age was statistically significant but had very small practical significance.
- Season fixed effects captured some unobserved variation.
- Team dichotomous variables were used for 4 statistically significant teams to account for more variation in ballpark and defense differences between teams.
- The final model explains a strong share of ERA variation, with an r-Squared score of .7888.

- **Implications**

- MLB teams can predict pitching performance reliably by using a small set of conventional metrics (WHIP, HR.9, K.9).
- The results can be compared with more advanced stats to see which is a better measure of predicting pitcher performance.

- **Conclusion**

- ERA can be predicted by using a small set of conventional statistics and fixed-effects modeling. The findings can provide guidance for player evaluation and resource allocation when signing players within professional baseball.

Introduction

Baseball is a very popular sport not only in America but in many Asian and Latin American countries as well. Its impact has spanned over 100 years and influences popular culture and linguistics. As America's past time, it is undoubtedly an important facet when looking at American culture throughout the years.

As the world has moved more digital and relied on computers more, all aspects of life have changed. Sports is no different in this regard. New types of stats and way of predicting performance have come along and have impacted sports front offices to re-evaluate how they evaluate players when signing and paying them. Of the 4 most popular sport leagues that exist in the US, Baseball is one of the pioneers in these analytics used by sports front offices.

Publications on analytics in baseball have existed since 1964 and sabermetrics in baseball have existed since the 1980s. These analyses and sabermetrics provide an empirical analysis into performance of each player and teams overall. Many of these saber metrics are not the typical statistics that people have historically used when looking at player performance like earned run average or batting average. The goal of these sabermetrics is to quantify more about the sport and provide more unbiased, mathematical ways to evaluate players. There are 4 main aspects that are looked at in sabermetrics, including hitting, pitching, base running, and fielding. However, despite existing for many years, there is still push back between if sabermetrics are a good tool to be using and if old school "eye tests" and traditional statistics are still sufficient in predicting player performance.

This project will focus on the metrics of pitching and attempt to explore the conventional statistics that can be used to predict earned run average (ERA), a main statistic for evaluating a pitchers performance, used by MLB front offices and other sports analysts. It will then discuss if using traditional statistics to predict pitchers performance is still a good tool, or if potentially using more advanced sabermetrics is better. We hypothesize that WHIP will have the most impact on ERA with age, team, strikeouts per 9 innings, and home runs per 9 innings all having a significant effect on ERA as well.

Literature Review

Since the idea of using regression and statistics has existed in baseball for many years, there are many other sources that have explored a similar topics. To narrow down the statistics to look at as the explanatory variables for ERA, the paper *Predicting Earned Run Average in Major League Baseball* by Paul C. Brendel was a valuable tool to look at. This paper goes into predicting ERA of pitchers using simple linear regression.

It looks at 4 different predictors of individual metrics for a pitcher, including strikeout to walk ratio, fastball velocity, first pitch strike percentage, and groundball to flyball percentage. These statistics are more about the individual pitcher's ability in a vacuum rather than their effect on opponent results. First pitch strike percentage is the pitcher's ability to throw a strike the first pitch of an at-bat and fastball velocity reflects how hard a given pitcher is able to throw. In this paper, it was discovered that of these 4 statistics, strikeout to walk ratio had the largest explanatory power of ERA with an R-Squared value of .32. The other three variables had very low R-Squared values, and even when added all together, the model's R-Squared value was only .36. This is an indication that model is not a very good fit for what it is attempting to do. Learning from this, instead of focusing on pitcher's individual abilities, the outcomes of opponents was noted as more impactful explanatory variables. Brendel explains that "K:BB may have been the best predictor since it was the only independent variable directly correlated with opponent on base results. Future studies may want to examine predictors that also have this quality, such as left on base percentage" (Brendel, n.d).

To address outcome based statistics, the next paper that was looked at was Beneventano, et al.'s paper *Predicting Run Production and Run Prevention in Baseball: The Impact of Sabermetrics*. This paper is attempting to compare whether sabermetrics are a good predictor of runs scored and run prevention in comparison to conventional statistics. The statistics for the explanatory variables were based more on opponents results and are as follows: Walks and Hits allowed per Inning Pitched (WHIP), Strikeouts per 9 innings, Home runs allowed per 9 innings, fielding percentage, and double plays turned. They also ran their model on sabermetric statistics, or statistics that are more advanced in calculation and may not make as much intuitive sense without the mathematical reasoning behind them. For their model, a stepwise-multiple-regression was used. The results of this model were promising, as their final model was able to get a very high R-Squared score of .988. They found that WHIP was the most important variable when predicting ERA, with the other variables still having significant impact. This paper was much more promising in its results as they had a very high R-Squared value. This led us to believe that opponent result statistics like WHIP and home runs per 9 innings were better predictors than individual pitching ability metrics. In this paper, they also examined the effect of defensive metrics like fielding percentage and double plays turned. These are stats that the pitcher is not directly responsible for, but rather the other 8 players on the field with them. This is an important thing to take note of and should not be ignored when making a model predicting ERA. One interesting note by the researchers is that none of the sabermetric statistics made it into their final model except WHIP, which was essentially just the combination of two conventional statistics, and is not as math based as some of the other sabermetric statistics. This meant that based on their findings that conventional statistics are still the best for determining pitcher performance.

The paper *Using the James-Stein Estimator to Predict Pitcher Performance* takes a very different approach to predicting ERA for a specific pitcher based on the previous year. The paper used a James-Stein estimator, which is a biased predictor that accounts for the pitcher's previous year performance and shrinks them back towards the group mean ERA. This brought the potential of biased results to the forefront and made us re-evaluate if this is something that needs to be accounted for in this project. While this is a valid concern, our project is aiming to predict a specific pitcher's ERA in a singular year and should not be affected by previous years in our calculation, so we did not adjust for it. However, the importance of normality testing was also discussed and made us make sure our errors were normally distributed.

Data and Identification Strategy

Data

The data for this project was obtained from a data set on kaggle. It has pitching data from 2015-2024 in both the hitting and pitching categories and got its data from publicly available sites like ESPN and fangraphs that have MLB data. Our variable of interest is earned run average and we are looking at the pitcher-year level.

The sample obtained for the project has 3141 observations. Each observation is a pitcher for one specific year that has at least 50 innings pitched. By limiting the data sets to pitchers with at least 50 innings pitched in a year, it allows for the pitchers who have very little innings and very outlier stats to be filtered out and not skew the results.

Results

% Table created by stargazer v.5.2.3 by Marek Hlavac, Social Policy Institute. E-mail: marek.hlavac at gmail.com % Date and time: Mon, Dec 08, 2025 - 10:14:33 PM

Table 1: Measuring the Relationship Between ERA and Conventional Pitching Statistics

	<i>Dependent variable:</i>	
	ERA	
	(1)	(2)
WHIP	4.713*** (0.063)	3.684*** (0.057)
HR.9		1.029*** (0.023)
K.9		-0.028*** (0.005)
CHC		0.196*** (0.057)
COL		-0.160*** (0.060)
MIL		-0.120** (0.056)
OAK		0.165*** (0.058)
Constant	-2.004*** (0.080)	-1.629*** (0.097)
Observations	3,141	3,141
R ²	0.644	0.788
Adjusted R ²	0.644	0.787
Residual Std. Error	0.707 (df = 3139)	0.546 (df = 3133)
F Statistic	5,671.500*** (df = 1; 3139)	1,662.744*** (df = 7; 3133)

Note:

*p<0.1; **p<0.05; ***p<0.01

Table 2: Measuring the Relationship Between ERA and Conventional Pitching Statistics with Season Fixed Effects

Dependent Variable:	ERA
Model:	Time FE Only (1)
<i>Variables</i>	
WHIP	3.677*** (0.0568)
HR.9	1.026*** (0.0234)
K.9	-0.0316*** (0.0054)
Age	-0.0075*** (0.0027)
COL	0.1905*** (0.0568)
CHC	-0.1508** (0.0598)
MIL	-0.1200** (0.0558)
OAK	0.1606*** (0.0582)
<i>Fixed-effects</i>	
Season	Yes
<i>Fit statistics</i>	
Observations	3,141
R ²	0.78995
Within R ²	0.78563
<i>IID standard-errors in parentheses</i>	
<i>Signif. Codes: ***: 0.01, **: 0.05, *: 0.1</i>	

Discussion Conclusions

References

Beneventano , P., Berger, P., & Weinberg, B. (2012). Predicting Run Production and Run Prevention in Baseball: The Impact of Sabermetrics. *International Journal of Business, Humanities and Technology* , 2, 67–75.

Brendel, P. C. (n.d.). Predicting Earned Run Average in Major League Baseball.

Using the James-Stein Estimator to Predict Pitcher Performance. (n.d.).

Appendix