

# Stats 202C: Homework #1

*Professor S.C. Zhu*

Assignment: 1-4

**Eric Chuu**

UID: 604406828

April 28, 2017

## Problem 1

(1) The period of a state  $i$  is given by

$$d_i = \gcd\{n \geq 1 : K_{ii}^n > 0\} \quad (1)$$

We also know that a state  $i$  is aperiodic if  $d_i = 1$ .

For  $K_0$ , we draw the transition graph diagram and see that all states can be reached in finite time with nonzero probability, so it is irreducible. Since the Markov chain is irreducible, then all states have the same period. Thus, it suffices to find the period of any state. If we begin in state 1, we can return to state 1 in two steps by going to state 2 and then back to state 1. We can also return to state 1 in three steps by going to state 2 then state 3, then returning back to state 1 directly. Then,  $d_1 = 1$ , and state 1 is aperiodic, and we conclude that  $K_0$  is aperiodic.

For  $K_1$ , we can see from the transition graph diagram,  $d_i = 1$  for  $i = 1, \dots, 5$ , so  $K_1$  is aperiodic. We can see that  $K_1$  is reducible by considering states 5 and 4. If we are in states 4 or 5, the probability of going into states 1, 2, or 3 is 0.

For  $K_2$ , we draw the transition graph diagram and see that all states can be reached in finite time with nonzero probability, so it is irreducible. For each state, we can see that  $d_i = 2$  for  $i = 1, \dots, 5$ , so  $K_2$  is periodic.

(2) The eigenvalues and left eigen-vectors for each matrix are given below.

(3) For each matrix, there are (insert number here) invariant probabilities.

(4) The eigen-values for  $K_0, K_0^{50}, K_0^{200}$  are plotted below.




Figure 1: Left: Eigenvalues for  $K_0$ . Center: Eigenvalues for  $K_0^{50}$ . Right: Eigenvalues for  $K_0^{200}$

(5) We compute the matrix  $K_0^{200}$ , and we see that it becomes the “ideal” matrix, where each row is  $\pi$ .

$$K_0^{200} = \begin{bmatrix} 0.809 & 0.2512 & 0.3071 & 0.2389 & 0.1219 \\ 0.809 & 0.2512 & 0.3071 & 0.2389 & 0.1219 \\ 0.809 & 0.2512 & 0.3071 & 0.2389 & 0.1219 \\ 0.809 & 0.2512 & 0.3071 & 0.2389 & 0.1219 \\ 0.809 & 0.2512 & 0.3071 & 0.2389 & 0.1219 \end{bmatrix}$$

□

## Problem 2

### 2.1

The plot of  $d_{\text{TV}(n), d_{\text{KL}(n)}}$  for the first 200 steps is shown below.

(2) We calculate the contraction coefficient  $K_0$  using

$$C(K_0) = \max_{x,y} \|K_0(x, \cdot) - K_0(y, \cdot)\|_{\text{TV}} \quad (2)$$

For two initial probabilities  $\nu_1, \nu_2$ ,

$$\|\nu_1 \cdot K_0 - \nu_2 \cdot K_0\|_{\text{TV}} \leq C(K_0) \|\nu_1 - \nu_2\|_{\text{TV}} \quad (3)$$

As  $\|\nu_1 - \nu_2\|_{\text{TV}} \leq 1$ , if  $C(K) < 1$ , then the convergence rate can be upper bounded by

$$A(n) = \|\nu_1 \cdot K_0^n - \nu_2 \cdot K_0^n\|_{\text{TV}} \leq C^n(K_0) \|\nu_1 - \nu_2\|_{\text{TV}} \leq C^n(K_0), \quad \forall \nu_1, \nu_2 \quad (4)$$

The plot for the bound  $C^n(K_0)$  over  $n = 1, \dots, 100$  is shown below.

(3) The Diaconis-Hanlon bound is given by

$$B(n) = \|\pi - \nu K_0^n\|_{\text{TV}} \leq \sqrt{\frac{1 - \pi(x_0)}{4\pi(x_0)}} \lambda_{\text{slem}}^n \quad (5)$$

where  $x_0 = 1$  is the initial state and  $\pi(x_0)$  is the target probability at  $x_0 = 1$  and  $\lambda_{\text{slem}}^n$  is the second largest eigen-value modulus. The plot of the real convergence  $d_{\text{TV}}$  in comparison with  $A(n), B(n)$  is given below.

**Problem 3**

Show that the Kullback-Leibler divergence decreases monotonically,

$$KL(\pi||\nu) - KL(\pi||\mu) = \mathbb{E}[KL(P(y, x) || Q(y, x))] \geq 0 \quad (6)$$

**Solution**