

## Copper oxide nanoparticles biosynthesized improve germination and bioactive compounds in wheat sprouts

Hortensia ORTEGA-ORTIZ<sup>1</sup>, Jazmín M. GAUCIN-DELGADO<sup>2\*</sup>,  
Pablo PRECIADO-RANGEL<sup>2</sup>, Manuel FORTIS-HERNANDEZ<sup>2</sup>,  
Luis G. HERNANDEZ-MONTIEL<sup>3</sup>, Efraín DE La CRUZ-LAZARO<sup>4</sup>,  
Liliana LARA-CAPISTRAN<sup>5</sup>

<sup>1</sup>Applied Chemistry Research Center, Department of Advanced Materials, Enrique Reyna H. No. 140, San José of the Cerritos, 25294, Saltillo, Coahuila, Mexico; [hortensia.ortega@cqua.edu.mx](mailto:hortensia.ortega@cqua.edu.mx)

<sup>2</sup>National Technological Institute of Mexico/ Torreon Technological Institute, Carretera Torreón-San Pedro km 7.5, 27170, Torreón, Coahuila, México; [fortismanuel@hotmail.com](mailto:fortismanuel@hotmail.com); [jazmontse@hotmail.com](mailto:jazmontse@hotmail.com); [ppreciador@yahoo.com.mx](mailto:ppreciador@yahoo.com.mx)

<sup>3</sup>Biological Research Center of the Northwest, Av. Politecnico Nacional 195, Col. Playa Palo Santa Rita, 23090, La Paz, Baja California Sur, Mexico; [lherandez@cibnor.mx](mailto:lherandez@cibnor.mx)

<sup>4</sup>Academic Division of Agricultural Sciences-Juarez Autonomous University of Tabasco, Km. 25, Villahermosa-Teapa Highway, Ra. The Huasteca, 2nd. Section. Center, Tabasco 86280, Mexico; [efrain.delacruz@ujat.mx](mailto:efrain.delacruz@ujat.mx)

<sup>5</sup>Veracruzana University-Xalapa Campus, Gonzalo Aguirre Beltrán University Circuit s/n, university area, Xalapa, Veracruz, Mexico; [llaracapistran@gmail.com](mailto:llaracapistran@gmail.com)

---

### Abstract (Please elaborate, extend; 200-250 words)

Nanotechnology is a tool that allows increasing germination and fast crop establishment under field conditions. This study determined the effect of copper oxide nanoparticles (CuONPs) on germination and production of phytochemical compounds in seedlings. Wheat seeds were treated with CuONPs suspensions: 0, 0.5, 1, 2, 4, and 6 mg mL<sup>-1</sup>. The use of 0.5 mg mL<sup>-1</sup> improved germination, vigor, plumule and radicle length besides increasing phytochemical compound biosynthesis in wheat sprouts. High concentration of CuONPs causes inhibiting effects due to Cu accumulation and phytotoxicity in plant tissue. However, their use in seed priming has a bearing in seedling germination and development, promoting bioactive compound biosynthesis.

---

**Keywords:** antioxidants; biosynthesized; CuONPs; *Triticum aestivum* L.

---

### Introduction

Nanotechnology is revolutionizing agriculture by producing agriproducts, such as nanopesticides, nanoherbicides and nanofertilizers that reduce environmental impact and increase crop yield (Kumar and Pandey, 2017; Younes et al., 2020). Among the main nanoproducts, metallic nanoparticles (NPs) have been the most used in producing nanofertilizers (Kolesnikov et al., 2019).

Metallic NPs increase seed germination in diverse plant species (Joshi et al., 2020) since they induce a greater synthesis of antioxidant compounds, such as enzymatic superoxide dismutase (SOD), catalase (CAT), ascorbate peroxidase (APX) and glutathione peroxidase (GPx) and non-enzymatic phenols, carotenoids,

glutathione, and vitamins. These compounds increase greater bioactivity (germination and content of phytochemical compounds) in seeds of *Sinapis alba* (Rajput et al., 2018), *Lactuca sativa L* (Pelegrino et al., 2020), *Medicago sativa* (Hong et al., 2015), *Eleusine coracana Gaertn* (Sathiyabama and Manikandan, 2018), *Ipomea batatas* (Bonilla-Bird et al., 2020), among others.

Antioxidant compounds are responsible for neutralizing and transforming toxic free radicals in non-harmful substances to maintain cellular homeostasis (Al-Hakkani, 2020). However, high doses of NPs of Ag, Ni, Zn, Ce, Ti, among other, have cytotoxic and genotoxic effects, decreasing plant germination and growth (Antisari et al., 2018).

Among the metallic NPs, those that stand out are copper (CuONPs) because they show antimicrobial and antiviral effects (Bridley et al., 2020). Their application in agriculture has demonstrated to have positive effects in plants because of their stability (Mali et al., 2019) and low concentrations used (Hernández-Hernández et al., 2019), besides showing they improve content of bioactive compounds (Lopez-Lima et al., 2020).

Moreover, wheat (*Triticum aestivum L.*) is one of the most important cereals at world level since it occupies a predominant role in total cereal production and global food security (Faraji et al., 2018). To assure an adequate productivity of this crop, high quality seed is required for an adequate germination since this stage is one of the most sensitive to adverse conditions and has an influence directly on crop yield (Victorava and Feoktistova, 2018). Therefore, the objective of this study is to assess the effect of CuONPs on wheat germination, growth, and phytochemical compounds.

## Materials and Methods

### *Study site and experimental conditions*

This study was performed in a biotechnology laboratory located in Institute Technologic, Torreon, Mexico at 24°30'27" N latitude, 102°10'40" W longitude at 1120 m.a.s.l.

### *Synthesis of copper oxide nanoparticles (CuONPs)*

The CuONPs synthesis was performed following the procedure described by Markova et al., (2010) and Taghavi et al. (2018), using copper salt  $\text{CuSO}_4 \bullet 5\text{H}_2\text{O}$  at 100 mM (>99.8%, Fermont<sup>®</sup>), aqueous solution and orange (*Citrus × sinensis*) peel as a reducing agent in a ratio of 4:1 (v/w). The mixture was maintained stirring at 250 rpm at 25°C for 24 h in a 4 L glass reactor on a heating blanket (EVELSA<sup>®</sup>). The activation of the chemical reaction was by colour change from emerald-green to dark green. Subsequently, the extract was vacuum filtered with Whatman #1 and #40, and heating was applied at 100 °C until solid CuONPs were obtained. Once the CuONPs synthesis was performed, the product was characterized by dynamic light scattering (DLS), X-Rays (XRD), Fourier-transform infrared (FTIR) spectroscopy by attenuated total reflection (ATR) and ultraviolet-visible (UV-vis) spectroscopy (synthesis and characterization were replicated five times).

### *Characterizing copper oxide nanoparticles by dynamic light scattering (DLS)*

The sample was prepared dispersing 1 mg CuONPs in 5 mL of deionized  $\text{H}_2\text{O}$  (Items S.A de C.V., MX) adding one drop of Tween<sup>®</sup> 80 (Merck, S.A de C.V., MX) at 1%. This technique was used to characterize nanoparticle size and distribution inferior to 1 nm with the particle size analyzer (Microtrac<sup>®</sup>, Nanotrac Wave II Q).

### *Characterizing copper oxide nanoparticles by fourier-transform infrared by ATR*

This study measured spectra in the region 4000-500  $\text{cm}^{-1}$  on an infrared spectrometer 550 FTIR (Nicolet<sup>®</sup>) using KBr powder. A spectrum was taken as the average of 160 scans to increase the signal of the

relationship noise and 4 cm<sup>-1</sup> spectral resolution and band frequency precision at 0.01 cm<sup>-1</sup>. Peak absorption intensities in this study were determined using the reference method (Vijilvani et al., 2019). The varieties in frequencies and band areas were determined with precision from the corrected spectrum from the original base line up to the corresponding control and treated samples. To detect absorption band intensities at the corresponding functional group concentration (Izumi et al., 2015), all spectral manipulation was performed with the software 8.0 for Windows (OMNIC Specta).

#### *Characterizing copper oxide nanoparticles by x-ray diffraction (XRD)*

The x-ray diffraction analysis was performed in a diffractometer (SIEMENS® D-500) using CuK $\alpha$  (25 mA, 35 kV) radiation, operating at a 40 kV and 40 mA at room temperature in a range of 2 $\theta$  from 10° to 80° and a speed of 0.02 °/s. This technique was used to identify the crystalline stage and approximate size of the biosynthesized CuONPs. The calculus to obtain average particle size was performed with the software Maud following Halder-Wagner's (Santos Filho et al., 2019) method.

#### *Characterizing copper oxide nanoparticles by ultraviolet-visible spectroscopy*

Absorbance of CuONPs solution was recorded using an UV-vis spectrophotometer (Shimadzu®, UV-2401PC) in a wavelength range of 190 nm at 500 nm. The sample was prepared by dispersing 1 mg CuONPs in 5 mL of deionized H<sub>2</sub>O (Items S.A de C.V., MX), adding one drop of Tween® 80 (Merck®) at 1%. This technique was used to identify transition metal ions in the solution.

#### *Treatments*

Copper oxide nanoparticles were dispersed in 50 mL sterile deionized H<sub>2</sub>O suspension; 0.5 mL of Agrex® (Agroenzimas S.A de C.V., MX) and Tween® 80 (Merck®) at 1% were added to the CuONPs solution with a concentration of 100 mg and stirred for 30 min in a shaker vortex (Fisher Scientific®, Vortex Genie 2). Subsequently, 950 mL of sterile deionized H<sub>2</sub>O were added, complementing the solution with 0.5 mL Agrex® (Agroenzimas S.A de C.V., MX) and 0.5 mL of Tween at 1% to a 1 L volume and stopped shaking at 12 h to avoid adding the dispersed particles (Chung et al., 2019). The CuONPs resulting suspension was diluted in distilled water and adjusted to concentrations 0, 0.5, 1, 2, 4, and 6 mg mL<sup>-1</sup> (Li et al., 2019).

#### *Plant material*

Wheat (*Triticum aestivum* L.) seeds -donated by the Cereal Program of Universidad Autónoma Agraria Antonio Narro were used with 96% germination, disinfected with ethanol solution at 75% for 5 min and washed four times with sterile distilled water (Li et al., 2019).

#### *Germination and growth parameter assay*

Seeds were placed in six batches with 10 seeds each (n = 60). The treatments were applied only once to the seed by inhibition using 50 mL of each CuONPs concentration. Seeds were placed within an artificial growth chamber (Labnet International Inc.®, 211DS) at a temperature of 25 °C ± 2 °C, respectively, with 60% relative humidity for 8 h in darkness (Salas-Pérez et al., 2016).

Once the inhibition period was completed, 10 seeds were deposited on Whatman #1 (Productos químicos del sur S.A. de C.V., MX) filter paper wet with 5 mL deionized water inside a Petri dish; then seeds were placed carefully with the embryo located downward. Subsequently, the Petri plates were sealed and placed in an artificial growth incubator (Labnet International Inc.®, 211DS) with a day/night 12-h cycle at 25 ± 2 °C, respectively, with 60% relative humidity measured with (BOSCH Digital Multi-Scanner GMS120G). The treatment distribution consisted of a completely randomized design with six replicates per treatment, considering a Petri plate as an experimental unit. Seed germination was quantified daily according to the International Seed Testing Association (ISTA, 1999), and growth parameters (germination percentage (G), germination Index (GI), seed vigor (V), plumule (PL), radicle (RL), fresh weight and dry weight were recorded

for seven days. Seed germination was considered when sprout length reached half the seed length (2 mm long) (Faraji et al., 2018).

#### *Parameters tested during bioassay development*

The following parameters were calculated: Germination percentage (G). For determining germination percentage, total count of normal seedlings was considered, and the result was expressed in percentage at seven days after germination.

$$G = \frac{\text{Nº germinated seeds}}{\text{Total nº of seeds}} \times 100 \quad (1)$$

Germination Index (GI). GI represents the product of relative seed germination by the radicle relative growth. It constitutes an interaction indicator of the factors that promote or inhibit germination, as well as the respective factors that favour or hinder radicle growth. This index expresses both germinated seed and growth percentages reached by the radicle during the bioassay. The germination index was calculated with seed relative germination (SRG = # of germinated seeds in the extract × 100) and relative radicle growth (RRG = radicle elongation in the extract × 100).

$$GI (\%) = (SRG \times RRG) / 100 \quad (2)$$

Seed vigor (V). The biological potential that favors rapid and uniform establishment under conditions is seed vigour, including unfavorable ones in field (Taghavi et al., 2018), that determine the activity and performance in growth since it allows identifying the differences between germination and emergence in the field, mainly when field conditions can cause stress. At day 4 after sowing, the first count, at 5 days the second count, at 6 days the third count and at seven days the fourth was performed for seedling (that have well root and plumule) data with a total development of 2.0 cm measured individually, with a digital vernier (Truper, CALDI-6MP; 14388), to determine seed vigour expressed in percentage.

$$V (\%) = \frac{\text{Normal seedlings}}{\text{Total nº of seeds}} \times 100 \quad (3)$$

Plumule (PL) and radicle (RL) length. Plumule length was measured from the root-hypocotyl junction up to the cotyledon base, whereas radicle length was measured from the hypocotyl base to the radicle at its apex. The values were expressed in centimetres.

Fresh weight. Fresh weight was recorded in sprouting mg<sup>-1</sup>. The sprouts of each experimental unit were placed in a watch glass recording sample weight in an analytical balance (Mettler Toledo, ADN® HR-200).

Dry weight. Dry weight was expressed in sprouting mg<sup>-1</sup>. Sprouts were placed in perforated and labelled brown paper bags. After that, they were placed in a drying stove (Novatech S.A de C.V, Ohasus® 547A) at a temperature of 72 °C for 24 h. At the end of each period, sample weight was recorded in an analytical balance (Mettler Toledo, ADN® HR-200).

#### *Extract preparation for phytochemical compounds*

To obtain the extracts, 2 g of fresh sprouts were mixed in 10 mL ethanol at 80% with constant centrifugation at 0.31 RCF (relative centrifugal force) for 24 h. Subsequently, extracts were centrifuged at 13.61 RCF for 5 min, and the supernatant was extracted for its subsequent analysis.

#### *Phytochemical compounds*

Total phenolic content was determined using a modification of Folin-Ciocalteau method (Singleton et al., 1999); 50 µL of ethanolic extract were taken, diluted in 3 mL of Milli-Q (MQ, Damstadt, DE) water; 250

$\mu\text{L}$  of Folin-Ciocalteau reagent (1N) was added, stirred, and left in reaction for 3 min. Subsequently, 750  $\mu\text{L}$  of  $\text{Na}_2\text{CO}_3$  (20%) and 950  $\mu\text{L}$  of MQ water were added. The solution was allowed to stand for 2 h, and the samples were quantified in an UV-Vis spectrophotometer at 760 nm. The standard was prepared with gallic acid, and the results were expressed in mg GAE 100  $\text{g}^{-1}$  fresh weight.

Total flavonoids were determined by colorimetry (Hidalgo et al., 2019); 250  $\mu\text{L}$  of ethanolic extract were taken, mixed with 1.25 mL of MQ water and 75  $\mu\text{L}$  of  $\text{NaNO}_2$  (5%). After a 5-min rest, 150  $\mu\text{L}$   $\text{AlCl}_3$  (aluminum chloride-1-Ethyl-3-methylimidazolium chloride, Sigma-Aldrich, St. Louis, MO, U.S.A.) were added. Subsequently, 500  $\mu\text{L}$  of  $\text{NaOH}$  (1M) and 275  $\mu\text{L}$  of MQ water were added and vigorously stirred; samples were quantified in an UV-Vis spectrophotometer at 510 nm. The standard was prepared with quercetin dissolved in absolute ethanol ( $y = 0.0122x - 0.0067$ ;  $r^2 = 0.965$ ), and the results were expressed in mg QE 100  $\text{g}^{-1}$  fresh weight.

Total antioxidant capacity was measured by in-vitro 2,2-diphenyl-1-picryl-hydrazyl-hydrate (DPPH<sup>+</sup>) method (Li et al., 2019). A DPPH<sup>+</sup> solution (Sigma-Aldrich, St. Louis) was prepared in ethanol at 0.025 mg  $\text{mL}^{-1}$  concentration; 50  $\mu\text{L}$  of ethanolic extract were mixed with 1950  $\mu\text{L}$  of DPPH<sup>+</sup> solution; after 30 min the samples were quantified in an UV-Vis spectrophotometer at 517 nm. The results were expressed in  $\mu\text{M}$  equivalent in Trolox 100  $\text{g}^{-1}$  fresh weight.

### Copper

Cu content in wheat sprouts was determined by atomic absorption spectrophotometry (Cartaya et al., 2017). Sprouts were dried in a stove at 70 °C in brown paper and subsequently macerated in a mortar. Cu was quantified in a hydride generation atomic absorption (AA) spectrophotometer (Lab Wrench Varian Spectr AA®, model 220Fast). The results were determined in  $\mu\text{g kg}^{-1}$  dry weight.


### Statistical analyses

The results obtained were analysed by analysis of variance (ANOVA) and comparison of means with Tukey's test ( $p \leq 0.05$ ) using the statistical package SAS (Statistical Analysis System Institute) version 9.4. Normality of data expressed as a percentage (germination percentage (G), germination Index (GI), seed vigor (V), was verified with Kolmogorov-Smirnov Test and were transformed applying arcsine and square root transformation before analysis of variance.


## Results and Discussion

### Green synthesis of copper oxide nanoparticles (CuONPs)

Nanoparticles were obtained in a size of >95 nm by DLS (Figure 1). The size of the particles depends on the biological entity and biochemical processing, as the effect of environmental conditions such as temperature, affect the size, shape, and morphology of the biosynthesized nanoparticles (Fardood et al., 2019). The nanoparticles obtained showed a cone shape with a length from 80-120 nm and thickness from 15-20 nm, specific characteristics of CuONPs (Ghidan et al., 2016). The nanoparticles measured by XRD indicated the specific formation of CuONPs with diffraction angles of 43° and 56°, assigned to the plane orientation 111 and 200 of CuO, respectively (Figure 2), indicating its crystalline nature (Wei et al., 2016). Therefore, the powder is characterized as CuONPs according to the international data base of the Joint Committee on Diffraction Standards for Dust (JCDPS 9-43) and the Joint Committee on Standards for Diffraction of Dust (No. 089-2838) (Trang et al., 2018; Van den Bruinhorst et al., 2016).



**Figure 1.** Micrographics obtained by histogram of the size distribution of the dynamic light scattering (DLS)  
Please replace, using a better image – quality, resolution etc.




**Figure 2.** XRD sample patterns of CuONPs powder obtained by green synthesized with orange peel extract  
and  $\text{CuSO}_4 \cdot 5 \text{ H}_2\text{O}$  to 100 mM for 24 h and treated subsequently at 100 °C

The functional groups of the responsible components of the ion bio-reduction and Cu ion stabilization were identified by FTIR spectrometer (Figure 3a) (Tadjarodi et al., 2015). The absorption peaks that appear at 1626 and 3297  $\text{cm}^{-1}$  could be water molecule flexing and stretching and the superficial hydroxyl groups in the CuONPs adsorbent surface (Nazar et al., 2018; Siddiqi et al., 2020). Absorption of infrared bands close to 589-660  $\text{cm}^{-1}$  show CuONPs characteristic vibration, confirming the formation of the expected CuO starting from green synthesis (Bhagat et al., 2020). Different authors have performed CuO synthesis studies and obtained characteristic links by FT-IR found from 400-700 nm (Siddiqi et al., 2020; Chaudhary et al., 2019; Qamar et al., 2020), which agree with CuO monoclinic phase (Taghavi et al., 2018). Possibly the carbonyl group of amino acid waste has a strong linkage capacity with metal, which suggests the formation of an organic cover on CuONPs to avoid agglomeration and provide medium stability (Chaudhary et al., 2019).

Figure 3b indicate the absorption spectra of UV-vis/ with the presence of the purified and dispersed CuONPs in deionized water, observing an absorption band close to 192 nm. This result confirms that reported

(Le Van et al., 2016) by the maximum appearance of absorbance at 200 nm for CuONPs, assigned to transition between central electron bands of Cu metals and CuO nanocrystals (Oli et al., 2018).



**Figure 3.** Spectra of (a) Fourier-Transform Infrared (FTIR) and (b) ultraviolet-visible (UV-vis) of CuONPs obtained by green synthesized with orange peel extract and  $\text{CuSO}_4 \cdot 5 \text{ H}_2\text{O}$  to 100 mM for 24 h and treated subsequently at 100 °C

#### *Effect of copper oxide nanoparticles on germination of wheat seed*


Germination and fresh weight are some of the main properties implied in seed quality (Prieto-Méndez et al., 2019). The highest dose of CuONPs affected the variables related with germination negatively (Table 1), due to cyto- and genotoxicity of CuONPs (Bonciu et al., 2018; Quiterio-Gutiérrez et al., 2019) and an alteration of DNA repairing kinetics (Escobedo-Paredes et al., 2020).

**Table 1.** Effect of copper oxide nanoparticle (CuONPs) doses on germination percentage (G), germination index (GI), vigor (V), plumule length (PL), radicle length (RL), fresh weight (FW) and dry weight (DW) in wheat sprouts

CuONPs mg mL⁻¹	G (%)	GI (%)	V (%)	PL (cm)	RL (cm)	FW (mg)	DW (mg)
0.0	90.0 a ± 1.02*	42.0 ab ± 0.25	63.9 ab ± 11.02	7.1 b ± 0.27	6.5 c ± 0.31	4891 a ± 0.04	325 a ± 0.04
0.5	90.0 a ± 0.98	43.2 a ± 0.07	81.4 a ± 9.25	9.1 a ± 0.057	7.2 a ± 1.17	4974 a ± 0.07	354 a ± 0.07
1.0	66.6 b ± 1.21	39.9 abc ± 0.09	64.1 ab ± 21.32	7.7 ab ± 0.047	6.8 b ± 0.51	4924 a ± 0.09	349 a ± 0.09
2.0	54.9 c ± 2.41	27.8 bc ± 0.04	61.3 ab ± 11.25	7.1 b ± 0.78	5.5 c ± 0.75	3564 b ± 0.06	302 b ± 0.06
4.0	45.1 d ± 1.04	26.7 c ± 0.05	55.1 b ± 13.44	6.0 b ± 0.065	5.2 d ± 0.85	3216 bc ± 0.07	295 bc ± 0.07
6.0	27.5 e ± 2.12	16.4 d ± 0.08	36.4 c ± 10.74	3.7 c ± 0.074	4.3 e ± 0.074	3051 c ± 0.05	115 c ± 0.05

\*Data are show as the mean ± standard deviation (SD, n = 50). Values with same letters in each column are equal according to Tukey's test ( $p \leq 0.05$ ).

The abiotic stress caused by NPs reduced germination growth (Heikal *et al.*, 2020). In contrast, with low doses of CuONPs, the variables related with germination improved (Chaudhary *et al.*, 2019; Shabbir *et al.*, 2020) when they showed a stimulating effect in shoot and radicle growth (Figure 4) as indicated by (García-Gómez and Fernández, 2019).



**Figure 4.** Germination and emergence of wheat seeds treated with copper oxide nanoparticles (CuONPs): (a) 0, (b) 0.5, (c) 1, (d) 2, (e) 4 and (f) 6 mg mL<sup>-1</sup>

Metal NPs improve water and micronutrient imbibition, accelerate mobilization of reserves during the first stages of the germination process (Costa Da and Sharma, 2016) and increase germination vigour (Quiterio-Gutiérrez *et al.*, 2019)[51] because of a greater activation of copper enzymes (Javed *et al.*, 2017) and a fast adenosine triphosphate (ATP) production by the copper dependent enzyme cytochrome C oxidase (Ramzan *et al.*, 2021). This process takes place when molecular oxygen ( $O_2$ ) reduction is catalyzed to water ( $H_2O$ ), generating an electric gradient used by the mitochondria (Kumar *et al.*, 2020; Mehdizadeh *et al.*, 2020). Where Cu enters the cell through a high affinity Cu (I) uptake system that includes CTR/COPT family proteins (Pham *et al.*, 2022). After entry, cytosolic chaperone proteins (of which ATX1 is a prototype) are responsible for subsequent Cu transfer to key metabolic Cu proteins (Shin *et al.*, 2012). More recently, GSH is implicated as a Cu ion carrier between the importers and metallochaperones (Miras *et al.*, 2008). Then, it activates the development of different efficient cellular mechanisms to cope with Cu excess (Paredes *et al.*, 2020; Rajput *et al.*, 2020), immobilizing it in the cytosol to maintain the metal at a low concentration (Xiong *et al.*, 2017) with vacuole and Cu chelation sequestration with metallothionein (MT) and phytochelatin (PC) (Singh *et al.*, 2018). CuONPs in the lowest concentration increased germination, a very important parameter because this stage is the most sensitive one to adverse conditions and influences directly in crop yield (Rastogi *et al.*, 2017). The germination parameter improvement by metallic NPs agrees with other works (Kolesnikov *et al.*, 2019; Mohan *et al.*, 2016), but a response pattern to the effects caused by CuONPs in seeds and sprouts is difficult to establish since it depends on species, concentration, type of NPs (Bhuvaneshwari *et al.*, 2018; Shabbir *et al.*, 2020), and application method (Xu *et al.*, 2020).

#### *Phytochemical compounds*

The high CuONPs dose decreased total phenols and flavonoids and the antioxidant capacity of germinated wheat seeds (Figure 5 a-c). CuONPs in high doses cause oxidative stress increasing biosynthesis of these phytochemicals (Ruttkay-Nedecky *et al.*, 2017) due to the high reactive oxygen species (ROS) production through the redox cycle (Mortezaee *et al.*, 2019). This process generates damages in the antioxidant defense system causing a rupture of the normal cellular function besides physiological and morphological damage in

different macromolecules, which help promote irreversible damage of lipids, nucleic acids, and cellular proteins (Rao et al., 2018).

Because free radicals tend to cause chain reactions (one radical generates another one, and so on), an unbalance emerges due to excess ROS production and a reduction of antioxidant defense mechanisms (Carvajal et al., 2017). Whereas the use of  $0.05 \text{ mg mL}^{-1}$  of CuONPs improves secondary metabolite biosynthesis in sprouts because of the stimulation exerted by antioxidant cellular biochemistry (Tocmo et al., 2020; Xu et al., 2020). This stimulation can be explained because copper can accept and donate stable electrons within the atom in its orbit, which avoids decoupling to other molecule, thus the importance of this mechanism for oxidation-reduction reactions. Then, the interaction of protector actions against ROS is possible and performed by regulating several enzymes (as superoxide dismutase [SOD], catalase [CAT], ascorbate-glutathione [AG] and glutathione peroxidase [GPx]) and non-enzymatic compounds (phenolic, flavonoids, vitamin C, and glutathione) as first line of defense against oxidative stress (Rai et al., 2018; Singh et al., 2018). Likely, CuONPs function bioactive molecules and dynamic systems that may provide an optimum environment to transport nutrients connectively guiding growth and cellular proliferation, as well as secondary metabolite production due to highly reactive and efficient NPs in their actions (López-Vargas et al., 2018). By participating as copper enzyme of the several enzymatic systems involved in formation and conversion of amino acids as in detoxification of superoxide radicals, they protect the cellular and subcellular system from the cytotoxic ROS effects (León-Silva et al., 2018). Thus, the parameters related with the antioxidant system of sprouts improve, which agrees with that reported by other authors (Fidelis et al., 2019; Prieto-Méndez et al., 2019).




Figure 5. Effect of CuONPs concentration in content of (a) total phenolic and (b) total flavonoid compounds; (c) antioxidant capacity; and (d) Cu content in wheat sprouts  
Data are show as the mean  $\pm$  standard deviation (SD,  $n = 50$ ).

Average values in the columns with different letters differ statistically among them (Tukey's test  $p \leq 0.05$ ).

#### *Concentration of copper in wheat sprouts*

Copper content in wheat sprouts increased as applied doses increased (Figure 5d). CuONPs distribution in plant species are contrasting because they depend on vegetal species, nanoparticle synthesis (concentration, size, shape) (Kolesnikov *et al.*, 2019) and application routes (seeds, leaves, substrate, among others) (Hernández-Hernández *et al.*, 2019; Kumar *et al.*, 2020).

Cu absorption and translocation within sprouts is determined by activating copper enzymes that oversee unfolding the proteins contained in storage places (Kranjc and Drobne, 2019), which in turn are digested in more simple substances to translocate to growth points by the cellular division of the embryo axis (Sun-Kou *et al.*, 2014). The greatest absorption of this element could have been due to copper, which is captured better by the passive apoplast membrane of the so-called transport proteins (Aguirre-Bolaños *et al.*, 2017; Ruddaraju *et al.*, 2020), since they are incorporated to metabolic pathways, such as plant copper enzymes altering the mitogen-activated protein kinase (El Shafey, 2020). Thus, these proteins participate in signaling the transduction induced by heavy metals (Hu and Gao, 2010) and protein phosphorylation, which are important in the enzymatic activity center (Parsai and Kumar, 2019) and a greater effect in antioxidant activity (Rajput *et al.*, 2020). Similar results have been reported, finding increase in Cu content because of adding CuONPs. Cu accumulation in sprouts could complement daily intake, which fluctuates from 440 to 900 µg (Lugo, 2016).

## Conclusions

This study showed that orange peel can be used efficiently for  $\text{CuSO}_4 \cdot \text{H}_2\text{O}$  reduction in CuONPs and the characterization by XRD which confirms copper present in the form of oxide with a particle size of 95 nm. Likewise, CuONPs in low concentration ( $0.5 \text{ mg mL}^{-1}$ ) stimulates germination, vigour, and plumule and root development, as well as non-enzymatic antioxidant compound synthesis. Moreover, high doses of CuONPs ( $6 \text{ mg mL}^{-1}$ ) caused an inhibitory effect due to phytotoxicity by high Cu accumulation in plant tissues. The use of CuO NPs for green synthesis is a viable alternative to obtain beneficial effects in germination and seedling development, as well as greater secondary metabolite production. Finally, future research should investigate the effects of the CuONPs in other varieties of wheat and confirmed their influence on the growth and productivity of the plants in field.

## Authors' Contributions

Conceptualization: JMGD, PPR; Methodology: JMGD, HOO; Validation: LLC, LGHM; Formal analysis: MFH, EdeCL; Investigation: JMGD, LLC; Data curation: PPR, LGHM; Funding acquisition: PPR, HOO; Project administration: PPR. Writing: JMGD, PPR; Review and editing HOO, PPR. All authors read and approved the final manuscript.

## Ethical approval (for researches involving animals or humans)

Not applicable.

## Acknowledgements

Jazmin Montserrat Gaucin Delgado appreciates the financial support provided by the National Council for Science and Technology of Mexico (CONACYT) for doctoral studies and the TECNM for the support

and sponsorship for the project. To Dr. Hortensia Ortega-Ortiz from CIQA for donating the nanoparticles and Bertha Alicia Puente Urbina for her technical assistance in obtaining the X-ray diffractograms.

### Conflict of Interests

The authors declare that there are no conflicts of interest related to this article.

### References

- Aguirre-Bolaños M, Benítez-Flores JC, González-Valle MR, Hernández-Portilla LB, Quintanar-Zúñiga RE, Flores-Ortiz CM (2017). Efecto del almacenamiento prolongado sobre la viabilidad y perfil de ácidos grasos en semillas de *Encyclia adenocarpa* (Lex.) Schltr. Revista Fitotecnia Mexicana 40(2):151-60. <https://doi.org/10.35196/rfm.2017.2.151-160>
- Al-Hakkani MF (2020). Biogenic copper nanoparticles and their applications: A review. Applied Sciences 2(3):1-20. <https://doi.org/10.1007/s42452-020-2279-1>
- Antisari LV, Carbone S, Bosi S, Gatti A, Dinelli G (2018). Engineered nanoparticles effects in soil-plant system: Basil (*Ocimum basilicum* L.) study case. Applied Soil Ecology 123:551-60. <https://doi.org/10.1016/j.apsoil.2018.01.007>
- Bhagat D, Suryawanshi I, Gurnule W, Sawant S, Chavan P (2021). Greener synthesis of CuO nanoparticles for enhanced development of latent fingerprints. Materials Today: Proceedings 36:747-750, <https://doi.org/10.1016/j.matpr.2020.05.357>
- Bhuvaneshwari V, Vaidehi D, Logpriya S (2018). Green synthesis of copper oxide nanoparticles for biological applications. Microbiol Curr Res. 2(1):9-10. <https://doi.org/10.4066/2591-8036.e105>
- Bonciu E, Firbas P, Fontanetti CS, Wusheng J, Karaismailoglu MC, Liu D (2018). An evaluation for the standardization of the *Allium cepa* test as cytotoxicity and genotoxicity assay. Caryologia 71(6):191-209. <https://doi.org/10.1080/00087114.2018.1503496>
- Bonilla-Bird N, Ye Y, Akter T, Valdes-Bracamontes C, Darrouzet-Nardi A, Saupe G (2020). Effect of copper oxide nanoparticles on two varieties of sweet potato plants. Plant Physiology Biochemistry 154:277-286. <https://doi.org/10.1016/j.plaphy.2020.06.009>
- Bridley A, Daffin Jr LW, Cuttler C, Campoli J, Cummings JA. (2020). Defining psychopathology. Abnormal Psychology. Retrieved 2021 August 20 from: <https://openpress.usask.ca/abnormalpsychology/chapter/part-3/>
- Cartaya O, Guridi F, Cabrera A, Moreno AM, Hernández Y (2017). Efecto de la aplicación foliar de oligogalacturónidos a plántulas de tomate (*Solanum Lycopersicum* L.) en la fitoextracción de cobre de suelo contaminado. Cultivos Tropicales 38:142-147.
- Carvajal C, Márquez M, Gutiérrez B, González-Vera A, Arellano J, Ávila M (2017). Aspectos de fisiología, deterioro y calidad en semilla de soya. Alcance 1:76-92.
- Chaudhary RG, Bhusari GS, Tiple AD, Rai AR, Somkuvar SR, Potbhare AK (2019). Metal/metal oxide nanoparticles: toxicity, applications, and future prospects. Current Pharmaceutical Design 25:4013-29. <https://doi.org/10.2174/138161282566191111091326>
- Chaudhary S, Rohilla D, Umar A, Kaur N, Shanavas A (2019). Synthesis and characterizations of luminescent copper oxide nanoparticles: toxicological profiling and sensing applications. Ceramics International 45:15025-15035. <https://doi.org/10.1016/j.ceramint.2019.04.239>
- Chung I-M, Rajakumar G, Subramanian U, Venkidasamy B, Thiruvengadam M (2019). Impact of copper oxide nanoparticles on enhancement of bioactive compounds using cell suspension cultures of *Gymnema sylvestre* (Retz.) R. Br. Applied Sciences 9:2165. <https://doi.org/10.3390/app9102165>
- Costa Da M, Sharma P (2016). Effect of copper oxide nanoparticles on growth, morphology, photosynthesis, and antioxidant response in *Oryza sativa*. Photosynthetica 54: 110-119. <https://doi.org/10.1007/s11099-015-0167-5>
- El Shafey AM (2020). Green synthesis of metal and metal oxide nanoparticles from plant leaf extracts and their applications: A review. Green Processing Synthesis 9:304-339. <https://doi.org/10.1515/gps-2020-0031>

- Escobedo Paredes CR, Rodríguez Soto JC, Contreras Quiñones M, Aspajo Villalaz C, Calderón Peña A, León Alcántara E (2020). Citotoxicidad y genotoxicidad de nanopartículas de cobre sobre *Allium cepa* L. (*Amaryllidaceae*). Arnaldoa 27:181-90. <http://dx.doi.org/10.22497/arnaldoa.27.27110>
- Faraji J, Sepehri A, Salcedo-Reyes JC (2018). Titanium dioxide nanoparticles and sodium nitroprusside alleviate the adverse effects of cadmium stress on germination and seedling growth of wheat (*Triticum aestivum* L.). Universitas Scientiarum 23:61-87. <https://doi.org/10.11144/javeriana.sc23-1.tdna>
- Fardood ST, Ramazani A, Moradi S (2017). Green synthesis of Ni–Cu–Mg ferrite nanoparticles using tragacanth gum and their use as an efficient catalyst for the synthesis of polyhydroquinoline derivatives. Journal of Sol-Gel Science Technology 82:432-439. <https://doi.org/10.1007/s10971-017-4310-6>
- Feoktistova Victorava L, Clark Feoktistova Y (2018). Metabolism of copper. Its consequences for human health. Medisur. 16(4):579-587.
- Fidelis M, de Moura C, Kabbas Junior T, Pap N, Mattila P, Mäkinen (2019). Fruit seeds as sources of bioactive compounds: sustainable production of high value-added ingredients from by-products within circular economy. Molecules 24:3854. <https://doi.org/10.3390/molecules24213854>
- García-Gómez C, Fernández MD. (2019). Impacts of metal oxide nanoparticles on seed germination, plant growth and development. Analysis, Fate, Toxicity of Engineered Nanomaterials in Plants 84:75-124. <https://doi.org/10.1016/bs.coac.2019.04.007>
- Ghidan AY, Al-Antary TM, Awwad AM (2016). Green synthesis of copper oxide nanoparticles using *Punica granatum* peels extract: Effect on green peach aphid. Environmental Nanotechnology Monitoring Management 6:95-98. <https://doi.org/10.1016/j.cnm.2016.08.002>
- Heikal YM, Şutan NA, Rizwan M, Elsayed A (2020). Green synthesized silver nanoparticles induced cytogenotoxic and genotoxic changes in *Allium cepa* L. varies with nanoparticles doses and duration of exposure. Chemosphere 243:125430. <https://doi.org/10.1016/j.chemosphere.2019.125430>
- Hernández-Hernández H, Quiterio-Gutiérrez T, Cadenas-Pliego G, Ortega-Ortiz H, Hernández-Fuentes AD, Cabrera de la Fuente M (2019). Impact of selenium and copper nanoparticles on yield, antioxidant system, and fruit quality of tomato plants. Plants 8:355. <https://doi.org/10.3390/plants8100355>
- Hidalgo A, Šaponjac VT, Ćetković G, Šeregelj V, Čadanović-Brunet J, Chiosa D, Brandolini AJ (2019). Antioxidant properties and heat damage of water biscuits enriched with sprouted wheat and barley. LWT 114:108423. <https://doi.org/10.1016/j.lwt.2019.108423>.
- Hong J, Rico CM, Zhao L, Adeleye AS, Keller AA, Peralta-Videa JR (2015). Toxic effects of copper-based nanoparticles or compounds to lettuce (*Lactuca sativa*) and alfalfa (*Medicago sativa*). Environmental Science: Processes and Applications 17:177-185. <https://doi.org/10.1039/C4EM00551A>
- Hu Y-L, Gao J-Q (2010). Potential neurotoxicity of nanoparticles. International Journal of Pharmaceutics 394:115-21. <https://doi.org/10.1016/j.ijpharm.2010.04.026>
- International Seed Testing Association (ISTA). (1999). International rules for seed testing. Rules 1:333.
- Izumi F, Ikeda T (2015). Implementation of the Williamson-Hall and Halder-Wagner methods into RIETAN-FP. Nagoya Institute of Technology Repository System 3:33-38.
- Javed R, Mohamed A, Yücesan B, Gürel E, Kausar R, Zia M (2017). CuO nanoparticles significantly influence *in vitro* culture, steviol glycosides, and antioxidant activities of *Stevia rebaudiana* Bertoni. Plant Cell, Tissue Organ Culture 131:611-620. <https://doi.org/10.1007/s11240-017-1312-6>
- Joshi H, Kumar R, Pandey D, Jariwala C (2020). Effect of plasma and nanotechnology on nutrient use efficiency of wheat (*Triticum aestivum* L.). Journal of Pharmacognosy Phytochemistry 9:137-141.
- Khan I, Raza MA, Awan SA, Shah GA, Rizwan M, Ali B (2020). Amelioration of salt induced toxicity in pearl millet by seed priming with silver nanoparticles (AgNPs): the oxidative damage, antioxidant enzymes and ions uptake are major determinants of salt tolerant capacity. Plant Physiology Biochemistry 156: 221-232. <https://doi.org/10.1016/j.jplphys.2020.09.018>
- Kolesnikov S, Timoshenko A, Kazeev KS, Akimenko YV, Myasnikova M (2019). Ecotoxicity of copper, nickel, and zinc nanoparticles assessment on the basis of biological indicators of chernozems. Eurasian Soil Science 52:982-987. <https://doi.org/10.18470/1992-1098-2020-1-130-136>
- Kranjc E, Drobne D (2019). Nanomaterials in plants: a review of hazard and applications in the agri-food sector. Nanomaterials 9:1094. <https://doi.org/10.3390/nano9081094>
- Kumar A, Pandey G (2017). A review on the factors affecting the photocatalytic degradation of hazardous materials. Material Science & Engineering International 1:1-10. <https://doi.org/10.15406/mseij.2017.01.00018>

- Kumar VK, Muthukrishnan S, Rajalakshmi RJ (2020). Phytostimulatory effect of phytochemical fabricated nanosilver (AgNPs) on *Psophocarpus tetragonolobus* (L.) DC. seed germination: An insight from antioxidative enzyme activities and genetic similarity studies. Current Plant Biology 23:100158. <https://doi.org/10.1016/j.cpb.2020.100158>
- Le Van N, Ma C, Shang J, Rui Y, Liu S, Xing BJ (2016). Effects of CuO nanoparticles on insecticidal activity and phytotoxicity in conventional and transgenic cotton. Chemosphere 144:661-670. <https://doi.org/10.1016/j.chemosphere.2015.09.028>
- León-Silva S, Fernández-Luqueño F, López-Valdez F (2018). Engineered nanoparticles: are they an inestimable achievement or a health and environmental concern? In: Packages E (Ed). Agricultural Nanobiotechnology. Springer 2:183-212. [https://doi.org/10.1007/978-3-319-96719-6\\_10](https://doi.org/10.1007/978-3-319-96719-6_10)
- Li R, He J, Xie H, Wang W, Bose SK, Sun Y (2019). Effects of chitosan nanoparticles on seed germination and seedling growth of wheat (*Triticum aestivum* L.). International Journal of Biological Macromolecules 126:91-100. <https://doi.org/10.1016/j.ijbiomac.2018.12.118>
- Lopez-Lima D, Mtz-Enriquez AI, Carrión G, Basurto-Cereceda S, Pariona N (2020). The bifunctional role of copper nanoparticles in tomato: Effective treatment for Fusarium wilt and plant growth promoter. Scientia Horticulturae 277:109810. <https://doi.org/10.1016/j.scienta.2020.109810>
- López-Vargas ER, Ortega-Ortíz H, Cadenas-Pliego G, de Alba Romenus K, Cabrera de la Fuente M, Benavides-Mendoza A (2018). Foliar application of copper nanoparticles increases the fruit quality and the content of bioactive compounds in tomatoes. Journal of Applied Sciences 8:1020. <https://doi.org/10.3390/app8071020>
- Lugo NT (2016). Papel del ácido fólico, zinc y cobre en la prevención primaria de los defectos congénitos. Revista Cubana de Medicina General Integral 32(4):1-10.
- Mali SC, Raj S, Trivedi RJ (2019). Biosynthesis of copper oxide nanoparticles using *Ericostemma axillare* (Lam.) leaf extract. Biochemistry Biophysics Reports 20:100699. <https://doi.org/10.1016/j.bbrep.2019.100699>
- Markova-Deneva I (2010). Infrared spectroscopy investigation of metallic nanoparticles based on copper, cobalt, and nickel synthesized through borohydride reduction method. Journal of the University of Chemical Technology Metallurgy 45:351-378.
- Mehdizadeh T, Zamani A, Froushan SMA (2020). Preparation of Cu nanoparticles fixed on cellulosic walnut shell material and investigation of its antibacterial, antioxidant and anticancer effects. Heliyon 6:e03528. <https://doi.org/10.1016/j.heliyon.2020.e03528>
- Minkina T, Rajput V, Fedorenko G, Fedorenko A, Mandzhieva S, Sushkova S (2020). Anatomical and ultrastructural responses of *Hordeum sativum* to the soil spiked by copper. Environmental Geochemistry Health 42:45-58. <https://doi.org/10.1007/s10653-019-00269-8>
- Miras R, Morin I, Jacquin O, Cuillel M, Guillain F, Mintz E (2008). Interacción entre glutatión, Atx1 y cobre. 1. Dimerización de Atx1 inducida por glutatión de cobre (I). Revista de Química Biológica Inorgánica: JBIC: Una publicación de la Sociedad de Química Biológica Inorgánica 13:195-205. <https://doi.org/10.1007/s00775-007-0310-2>
- Mortezaee K, Najafi M, Samadian H, Barabadi H, Azarnezhad A, Ahmadi AJ (2019). Redox interactions and genotoxicity of metal-based nanoparticles: A comprehensive review. Chemico-Biological Interactions 312:108814. <https://doi.org/10.1016/j.cbi.2019.108814>
- Mosa KA, El-Naggar M, Ramamoorthy K, Alawadhi H, Elnaggar A, Wartanian S (2018). Copper nanoparticles induced genotoxicity, oxidative stress, and changes in Superoxide Dismutase (SOD) gene expression in cucumber (*Cucumis sativus*) plants. Frontiers in Plant Science 9:872-882. <https://doi.org/10.3389/fpls.2018.00872>
- Nazar N, Bibi I, Kamal S, Iqbal M, Nouren S, Jilani K (2018). Cu nanoparticles synthesis using biological molecule of *P. granatum* seeds extract as reducing and capping agent: Growth mechanism and photo-catalytic activity. International Journal of Biological Macromolecules 106:1203-10. <https://doi.org/10.1016/j.ijbiomac.2017.08.126>
- Oli HB, Sharma N, Ekaraj K, Subedee A, Timilsina R (2018). Green synthesis of copper nanoparticles using *Zingiber officinale* extract and characterization. Journal of Nepal Chemical Society 39:10-17. <https://doi.org/10.3126/jncts.v39i0.27008>
- Paredes CRE, Soto JCR, Quiñones MC, Villalaz CA, Peña AC, Alcántara EL (2020). Citotoxicidad y genotoxicidad de nanopartículas de cobre sobre *Allium cepa* L. (Amaryllidaceae). Arnaldoa 27:e108-e112. <http://dx.doi.org/10.22497/arnaldoa.271.27110>

- Parsai T, Kumar A (2019). Understanding effect of solution chemistry on heteroaggregation of zinc oxide and copper oxide nanoparticles. Chemosphere 235:457-469. <https://doi.org/10.1016/j.chemosphere.2019.06.171>
- Pelegrino MT, Kohatsu MY, Seabra AB, Monteiro LR, Gomes DG, Oliveira HC, ... Lange CN (2020). Effects of copper oxide nanoparticles on growth of lettuce (*Lactuca sativa* L.) seedlings and possible implications of nitric oxide in their antioxidative defense. Environmental Monitoring and Assessment 192:1-14. <https://doi.org/10.1007/s10661-020-8188-3>
- Pham KL, Schmollinger S, Merchant SS, Strenkert D (2022). Chlamydomonas ATX1 is essential for Cu distribution to multiple cupro-enzymes and maintenance of biomass in conditions demanding cupro-enzyme-dependent metabolic pathways. Plant Direct 6:e383. <https://doi.org/10.1002/pld3.383>
- Prieto-Méndez J, Aquino-Torres E, Pérez-Ríos SR, Trejo-González N, Prieto-García F (2019). Seed germination of beans (*Phaseolus vulgaris*) with nano-particles of iron. Bulgarian Journal of Agricultural Science 25:283-287.
- Qamar H, Rehman S, Chauhan DK, Tiwari AK, Upmanyu V (2020). Green synthesis, characterization and antimicrobial activity of the copper oxide nanomaterial derived from *Momordica charantia*. International Journal of Nanomedicine 15:2541-2553. <https://doi.org/10.2147/IJN.S240232>
- Quiterio-Gutiérrez T, Ortega-Ortiz H, Cadenas-Pliego G, Hernández-Fuentes AD, Sandoval-Rangel A, Benavides-Mendoza A (2019). The application of selenium and copper nanoparticles modifies the biochemical responses of tomato plants under stress by *Alternaria solani*. International Journal of Molecular Sciences 20:1950. <https://doi.org/10.3390/ijms20081950>
- Rafique M, Shaikh AJ, Rasheed R, Tahir MB, Bakhat HF, Rafique MS (2017). A review on synthesis, characterization and applications of copper nanoparticles using green method. Nano Energy 12:1750043. <https://doi.org/10.1142/S1793292017500436>
- Rai M, Ingle AP, Pandit R, Paralikar P, Shende S, Gupta I (2018). Copper and copper nanoparticles: role in management of insect-pests and pathogenic microbes. Nanotechnology Reviews 7:303-315. <https://doi.org/10.1515/ntrev-2018-0031>
- Rajput V, Minkina T, Sushkova S, Behal A, Maksimov A, Blicharska E (2020). ZnO and CuO nanoparticles: A threat to soil organisms, plants, and human health. Environmental Geochemistry Health 42:147-158. <https://doi.org/10.1007/s10653-019-00317-3>
- Rajput V, Minkina T, Suskova S, Mandzhieva S, Tsitsuashvili V, Chapligin V (2018). Effects of copper nanoparticles (CuO NPs) on crop plants: a mini review. BioNanoScience 8:36-42. <https://doi.org/10.1007/s12668-017-0466-3>
- Ramzan R, Kadenbach B, Vogt S (2021). Multiple mechanisms regulate eukaryotic cytochrome C oxidase. Cells 10:514. <https://doi.org/10.3390/cells10030514>
- Rao MP, Sathishkumar P, Mangalaraja RV, Asiri AM, Sivashanmugam P, Anandan S (2018). Simple and low-cost synthesis of CuO nanosheets for visible-light-driven photocatalytic degradation of textile dyes. Journal of Environmental Chemical Engineering 6:2003-2010. <https://doi.org/10.1016/j.jece.2018.03.008>
- Rastogi A, Zivcak M, Sytar O, Kalaji HM, He X, Mbarki S (2017). Impact of metal and metal oxide nanoparticles on plant: a critical review. Frontiers in Chemistry 5:78. <https://doi.org/10.3389/fchem.2017.00078>
- Ruddaraju LK, Pammi SVN, sankar Guntuku G, Padavala VS, Kolapalli VRM (2020). A review on anti-bacterials to combat resistance: From ancient era of plants and metals to present and future perspectives of green nano technological combinations. Asian Journal of Pharmaceutical Sciences 15:42-59. <https://doi.org/10.1016/j.japs.2019.03.002>
- Ruttkay-Nedecky B, Krystofova O, Nejdl L, Adam V (2017). Nanoparticles based on essential metals and their phytotoxicity. Journal of Nanobiotechnology 15:33-39. <https://doi.org/10.1186/s12951-017-0268-3>
- Salas-Pérez L, Gaucín-Delgado JM, Preciado-Rangel P, Fortis-Hernández M, Valenzuela-García JR, Ayala-Garay AV (2016). Efecto del ácido benzoico en la capacidad antioxidante de germinados de trigo. Revista Mexicana de Ciencias Agrícolas 17:3397-3404.
- Santos Filho Rd, Vicari T, Santos SA, Felisbino K, Mattoso N, Sant'Anna-Santos BF (2019). Genotoxicity of titanium dioxide nanoparticles and triggering of defense mechanisms in *Allium cepa*. Genetics Molecular Biology 42:425-435. <https://doi.org/10.1590/1678-4685-GMB-2018-0205>
- Sathiyabama M, Manikandan A (2018). Application of copper-chitosan nanoparticles stimulate growth and induce resistance in finger millet (*Eleusine coracana* Gaertn.) plants against blast disease. Journal of Agricultural Food Chemistry 66:1784-1790. <https://doi.org/10.1021/acs.jafc.7b05921>

- Shabbir Z, Sardar A, Shabbir A, Abbas G, Shamshad S, Khalid S (2020). Copper uptake, essentiality, toxicity, detoxification and risk assessment in soil-plant environment. *Chemosphere* 259:127436. <https://doi.org/10.1016/j.chemosphere.2020.127436>
- Shin, LJ, Lo, JC y Yeh, KC (2012). Copper chaperone antioxidant protein1 is essential for copper homeostasis. *Plant Physiology* 159:1099-1110. <https://doi.org/10.1104/pp.112.195974>
- Siddiqi KS, Husen A (2020). Current status of the manufacture of copper / copper oxide nanoparticles based on plant metabolites and their applications: a review. *Biomaterials Research* 24:11-16. <https://doi.org/10.1186/s40824-020-00188-1>
- Singh J, Dutta T, Kim K-H, Rawat M, Samddar P, Kumar PJJon (2018). 'Green'synthesis of metals and their oxide nanoparticles: Applications for environmental remediation. *Tropical Plant Research* 16:84-75. <https://doi.org/10.1186/s12951-018-0408-4>
- Singleton VL, Orthofer R, Lamuela-Raventós RM (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. Elsevier 1:152-178. [https://doi.org/10.1016/S0076-6879\(99\)99017-1](https://doi.org/10.1016/S0076-6879(99)99017-1)
- Sun-Kou MdR, Obregón-Valencia D, Pinedo-Flores Á, Paredes-Doig AL, Aylas-Orejón J (2014). Adsorción de metales pesados empleando carbones activados preparados a partir de semillas de aguaje. *Revista de la Sociedad Química del Perú* 80:225-236.
- Tadjarodi A, Akhavan O, Bijanzad K (2015). Photocatalytic activity of CuO nanoparticles incorporated in mesoporous structure prepared from bis (2-aminonicotinate) copper (II) microflakes. *Transactions of Nonferrous Metals Society of China* 25:3634-3642. [https://doi.org/10.1016/S1003-6326\(15\)64004-3](https://doi.org/10.1016/S1003-6326(15)64004-3)
- Taghavi FS, Ramazani A, Asiabi P, Joo S (2018). A novel green synthesis of copper oxide nanoparticles using a Henna extract powder. *Journal of Structural Chemistry* 59:1796-1801. <https://doi.org/10.1134/S0022476618070302>
- Trang NTH, Dung PD, Van PTH, Thang PQ (2018). Green synthesis of copper nanoparticles using mandarin (*Citrus reticulata*) peel extract and antifungal study. *Asian Journal of Biotechnology and Bioresource Technology* 3:1-9. <https://doi.org/10.9734/AJB2T/2018/39126>
- Van den Bruinhorst A, Kouris P, Timmer J, de Croon M, Kroon M (2016). Exploring orange peel treatment with deep eutectic solvents and diluted organic acids. *Natural Products Chemistry Research* 4:1-5. <https://doi.org/10.4172/2329-6836.1000242>
- Vijilvani C, Bindhu MR, Frincy FC, AlSalhi MS, Sabitha S, Saravanakumar K, ... Atif M (2020). Antimicrobial and catalytic activities of biosynthesized gold, silver and palladium nanoparticles from *Solanum nigrum* leaves. *Journal of Photochemistry and Photobiology B: Biology* 202:111713. <https://doi.org/10.1016/j.jphotobiol.2019.111713>
- Wei Y, Fang Z, Zheng L, Tan L, Tsang EP (2016). Green synthesis of Fe nanoparticles using *Citrus maxima* peels aqueous extracts. *Materials Letters* 185:384-386. <https://doi.org/10.1016/j.matlet.2016.09.029>
- Wojtkowiak K, Stępień A (2015). Nutritive value of spelt (*Triticum aestivum* spp. *spelta* L.) as influenced by the foliar application of copper, zinc and manganese. *Zemdirbyste-Agriculture* 102:389-396. <https://doi.org/10.13080/z-2015.102.049>
- Xiong T, Dumat C, Dappe V, Vezin H, Schreck E, Shahid M (2017). Copper oxide nanoparticle foliar uptake, phytotoxicity, and consequences for sustainable urban agriculture. *Environmental Science* 51:5242-5251. <https://doi.org/10.1021/acs.est.6b05546>
- Xu W, Xiang P, Liu X, Ma LQJC (2020). Closely-related species of hyperaccumulating plants and their ability in accumulation of As, Cd, Cu, Mn, Ni, Pb and Zn. *Quimiosfera* 34:126334. <https://doi.org/10.1016/j.chemosphere.2020.126334>
- Younes N, Hassan HS, Elkady MF, Hamed A, Dawood MF (2020). Impact of synthesized metal oxide nanomaterials on seedlings production of three Solanaceae crops. *Heliyon* 6:e03188. <https://doi.org/10.1016/j.heliyon.2020.e03188>



The journal offers free, immediate, and unrestricted access to peer-reviewed research and scholarly work. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.



**License** - Articles published in *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* are Open-Access, distributed under the terms and conditions of the Creative Commons Attribution (CC BY 4.0) License.  
© Articles by the authors; UASVM, Cluj-Napoca, Romania. The journal allows the author(s) to hold the copyright/to retain publishing rights without restriction.