

Evaluating Ecohydrological Theories of Woody Root Distribution in the Kalahari

Abinash Bhattachan¹, Mokganedi Tatlhego², Kebonye Dintwe^{3,4}, Frances O'Donnell⁵, Kelly K. Caylor⁵, Gregory S. Okin³, Danielle O. Perrot⁶, Susan Ringrose², Paolo D'Odorico^{1,7*}

1 Department of Environmental Sciences, University of Virginia, Charlottesville, Virginia, United States of America, **2** Okavango Research Institute, University of Botswana, Maun, Botswana, **3** Department of Geography, University of California Los Angeles, Los Angeles, California, United States of America, **4** Department of Forestry and Range Resources, Gaborone, Botswana, **5** Department of Civil and Environmental Engineering, Princeton University, Princeton, New Jersey, United States of America, **6** Department of Geography, University of Colorado, Boulder, Colorado, United States of America, **7** School of Architecture, Civil and Environmental Engineering, École Polytechnique Fédérale De Lausanne, Lausanne, Switzerland

Abstract

The contribution of savannas to global carbon storage is poorly understood, in part due to lack of knowledge of the amount of belowground biomass. In these ecosystems, the coexistence of woody and herbaceous life forms is often explained on the basis of belowground interactions among roots. However, the distribution of root biomass in savannas has seldom been investigated, and the dependence of root biomass on rainfall regime remains unclear, particularly for woody plants. Here we investigate patterns of belowground woody biomass along a rainfall gradient in the Kalahari of southern Africa, a region with consistent sandy soils. We test the hypotheses that (1) the root depth increases with mean annual precipitation (root optimality and plant hydrotropism hypothesis), and (2) the root-to-shoot ratio increases with decreasing mean annual rainfall (functional equilibrium hypothesis). Both hypotheses have been previously assessed for herbaceous vegetation using global root data sets. Our data do not support these hypotheses for the case of woody plants in savannas. We find that in the Kalahari, the root profiles of woody plants do not become deeper with increasing mean annual precipitation, whereas the root-to-shoot ratios decrease along a gradient of increasing aridity.

Citation: Bhattachan A, Tatlhego M, Dintwe K, O'Donnell F, Caylor KK, et al. (2012) Evaluating Ecohydrological Theories of Woody Root Distribution in the Kalahari. PLoS ONE 7(3): e33996. doi:10.1371/journal.pone.0033996

Editor: Han Y. H. Chen, Lakehead University, Canada

Received: July 5, 2011; **Accepted:** February 23, 2012; **Published:** March 28, 2012

Copyright: © 2012 Bhattachan et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Funding for this study came from the National Science Foundation DEB #0717360. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: paolo@virginia.edu

Introduction

Savannas are mixed plant communities with tree and grass species coexisting in the same landscape. They cover about 20% of the global land surface, including approximately one-half of Africa and Australia, 45% of South America and 10% of India and Southeast Asia [1,2,3]. Savannas are home to a large portion of the human population, and provide important ecosystem services such as rangelands for livestock grazing [2] and carbon storage [4,5]. Investigating root distribution in savanna vegetation and associated soil carbon pools is an important step towards the assessment of the global carbon budget [6].

Research in savanna ecology has long-recognized four major determinants of ecosystem structure, namely: fire, herbivory, water, and nutrient availability [1,3,7,8,9,10]. The structure of savanna vegetation is strongly governed by the spatiotemporal distribution of these four factors. Vegetation structure may, in turn, play a major role in the distribution of abiotic resources such as energy, water and nutrients [2,11,12]. For example, tree canopies redistribute rain water [13] and shade the ground, reducing the rates of soil evaporation and maintaining higher moisture levels in the subcanopy soils [2,11]. Similarly, the presence of tree roots and higher soil organic matter content enhance soil infiltration capacity beneath trees [14,15]. As a result,

the dynamics of soil moisture are strongly affected by the distribution of tree canopies and their accompanying root systems [11,16,17].

Processes governing vegetation composition and structure in savannas have been, for the most part, inferred from the study of patterns of aboveground plant biomass. Canopy cover, tree spacing, and the amount of woody biomass in relation to herbaceous biomass have been often considered the descriptors of vegetation structure in these ecosystems (e.g., [18,19,20]). However, the dynamics of savanna vegetation are strongly dependent on belowground processes such as competition for water and nutrients [21,22,23]. Root distribution affects plants' ability to compete for soil resources [2,16,24], ecosystem carbon storage [13,14,25] and water redistribution within the soil profile [26,27,28]. Some classic theories of tree-grass coexistence and plant geographic distribution are based on specific assumptions on root structure and function (e.g., [29,30]). For example, it has long been assumed (e.g., [29]) that tree-grass coexistence is due to niche separation between the woody and herbaceous life forms that exploit different soil layers with minimal overlap between their root zones (e.g., [31]). However, the assumption that woody plants grow most of their roots in deeper soil layers while herbaceous vegetation has roots only in the shallow soils has been repeatedly challenged by a number of studies [21,24,32,33,34]. Therefore




Figure 1. Geographic location of the study region and of the research sites.

doi:10.1371/journal.pone.0033996.g001

Table 1. Grain size analysis of the soil samples was conducted using a particle size analyzer (LS 13 320, Beckman Coulter®).

Site	Depth (m)	% clay	% silt	% sand
Shakawe	0.1	0.6	1	98.4
	0.3	0.7	1.7	97.6
	0.7	0.6	0.7	98.7
	1.2	0.5	0.7	98.8
Ghanzi	0.1	3	1	96.0
Tshane	0.1	0.9	4.1	95.0
	0.3	1	5.9	93.1
	0.7	1	6.8	92.2
	1.2	1.1	8.2	90.7
Bokspits	0.1	0.8	3.1	96.1
	0.3	0.7	2.1	97.2
	0.7	0.7	2.6	96.7
	1.2	0.6	2.7	96.7

Soil samples were collected from depths 0–0.1 m, 0.1–0.3 m, 0.3–0.7 m, and 0.7–1.2 m at all sites except for the Kuke/Ghanzi area, where we have used only soils from the top 10 cm.

doi:10.1371/journal.pone.0033996.t001

new advances in savanna ecology require a better understanding of patterns of root distribution. Studies on rooting depth and root profiles are also essential for regional and global-scale assessments of belowground carbon storage and climate modeling, as they define the thickness of the soil layer exploited by plants for water and nutrient uptake (e.g., [35,36]).

Two major issues need to be addressed to improve our current understanding of root structure and patterns of belowground biomass in savannas. First, the niche separation hypothesis is based on the assumption that savanna trees are deeply rooted. However, the rooting depth of woody plants in savannas remains poorly investigated. In fact, it is not clear how the rooting depth varies with different conditions of water availability. Theoretical studies have shown that optimal root profiles (i.e., the profiles that maximize plant transpiration while minimizing water stress) become deeper in wetter climates [37,38,39,40,41,42]. However, field observations along aridity gradients have shown a good agreement with this theory only in the case of herbaceous vegetation, while no significant relationship has been found between the rooting depth of woody vegetation and precipitation [21]. A possible explanation could be that, while in water-limited ecosystems roots are expected to become deeper with increasing mean annual precipitation, in more mesic environments root profiles are determined also by nutrient limitations. Schenk [43] listed a number of other reasons why there is an advantage for roots to be shallow and noted that roots “tend to be as shallow as

Table 2. Coordinates, mean annual precipitation (MAP), and plant community composition of the study sites.

Site	Coordinates	MAP (mm)	Main Woody Species	Main Grass Species
Shakawe	18°21'51"S, 21°50'31"E	539±170 (209.4)	<i>Terminalia sericea</i> , <i>Ochna pulchra</i> , <i>Pterocarpus angolensis</i> , <i>Burkea africana</i>	<i>Aristida meridionales</i> , <i>Eragrostis</i> spp, <i>Schmidtia pappophoroides</i>
Kuke	20°58'36"S, 22°28'48"E	439±157 (162.2)	<i>Terminalia sericea</i> , <i>Lonchocarpus nelssi</i> , <i>Acacia erubescence</i> , <i>Acacia fleckii</i>	<i>Eragrostis lehmanniana</i> , <i>Schmidtia</i> <i>pappophoroides</i> , <i>Anthphora pubescens</i>
Tshane	24°01'01"S, 21°52'08"E	358±133 (156.5)	<i>Lycium</i> species, <i>Acacia mellifera</i> , <i>Acacia luderitzii</i>	<i>Aristida congesta</i> , <i>Eragrostis lehmanniana</i> , <i>Eragrostis pallensis</i>
Bokspits	26°53'39"S 20°41'54"E	177±107 (55.2)	<i>Rhigozum trichotomum</i> <i>Acacia mellifera</i> , <i>Acacia erioloba</i> , <i>Boscia albitrunca</i>	<i>Stipagrostis amabilis</i> , <i>Stipagrostis</i> <i>uniplumis</i> , <i>Schmidtia kalahariensis</i>

Information on precipitation includes mean annual precipitation (MAP) ± the standard deviation of annual precipitation and (in parentheses) the minimum annual precipitation recorded in 1971–2006. Because rainfall data from Kuke do not exist, the values reported are from Ghanzi.
doi:10.1371/journal.pone.0033996.t002

possible and as deep as needed to fulfill evapotranspirational demands".

Second, the assessment of carbon storage in savannas requires a better understanding of the relation between above and below-ground biomass and its dependence on the rainfall regime. Root systems are often referred to as "the hidden half" (e.g., [44]) and carbon budget studies sometime assume live belowground biomass to be similar in magnitude to the above ground biomass [45]. However, the relation between root and shoot biomass appears to be more complex [46]. It has been argued that plants have a way of regulating the growth of their above- and belowground biomass [47] maintaining a "functional equilibrium" [47,48,49]. According to this equilibrium theory, limitations in the availability of aboveground resources (for example, light) induce an increase in aboveground plant biomass (e.g., leaves), whereas root growth is stimulated by limitations in belowground resources. The "functional equilibrium theory" would suggest that the root-to-shoot ratio (R:S, the ratio of belowground to aboveground biomass) should increase from mesic to xeric environments, due to the decrease in soil moisture levels and decrease in vegetation density and competition for light. Patterns of belowground primary production are consistent with the "functional equilibrium hypothesis" in the case of herbaceous vegetation in tropical savannas [50]. However, it remains to be seen whether this hypothesis can explain patterns of above- and belowground woody biomass in savannas.

Although global scale reviews (e.g., [4,21,51,52]) are in agreement with ecological theories on the relationship between root structure and water availability (e.g., [30,37,47]), they tend to mix studies made on different soil textures and presumably using

different sampling methods. To avoid these issues, we investigate root depths and R:S ratios at sites in the Kalahari, where a relatively homogeneous sandy soil exists along a latitudinal rainfall gradient [53,54]. This unique observational setting allows us to examine the "optimal rooting depth" (e.g., [21,37,38,42,52]) and the "functional equilibrium" [47] theories of root structure across a large gradient of climate independent of major changes in soil type. More specifically, we will use data from a major root sampling effort to study how the rooting depth of woody plants and the ratio between belowground and aboveground biomass vary along the Kalahari's rainfall gradient.

Methods

Study sites

The Kalahari sand sheet is one of the largest continuous sand deposits on Earth, stretching for several thousand kilometers across a rainfall gradient, with no substantial variability in the soil physical properties [54]. Known as the Kalahari Transect (KT), this region is an ideal natural laboratory for ecohydrologic global change studies [55,56] in that the effects of changing hydrologic conditions can be investigated without the confounding effects of different soil types.

This study involves four field sites located within the Botswana portion of the KT (Fig. 1), along a south-north rainfall gradient ranging between 180 and about 550 mm/yr with an average storm depth of 10 mm d⁻¹ [57]. More details on the seasonality of precipitation are provided in Figure 1. All sites exhibit savanna vegetation found on a very thick (>100 m) homogeneous sandy substrate with consistent physical and geochemical properties. The grain size analysis of soil collected at depths 0–10 cm, 10–30 cm, 30–70 cm, and 70–120 cm shows that soils are consistently sandy (>90% sand) throughout the top 1.2 m profile and across the entire transect (Table 1). Given the geomorphic context (i.e., the Kalahari sand sheet) there is no reason to believe that soil texture differences would play a role in root distribution. Water table depth ranges from about 25 m at the wettest site (Shakawe) to about 100 m deep at the driest (Bokspits). Vegetation composition and structure vary across the study region from an open fine-leaf shrub savanna in the south, through a bush savanna, to a mixture of bush and broad leaf woodland savanna in the northwestern part of Botswana [58].

Individual study site characteristics (i.e. location, rainfall regime and dominant woody and herbaceous vegetation) are summarized in Table 2. Mean annual precipitation (MAP) values used in this study were calculated using meteorological records (Botswana

Table 3. Results of the analysis of covariance performed on the wet to dry biomass ratio, *k*, using tissue type (i.e., above ground or below ground biomass) as a factor and size of the wet sample as a covariate.

Variable	Estimate	Std. Error	t - value	p - value
<i>k</i> (aboveground)	1.64	0.0264	61.9	1.33E-138
<i>k</i> (roots)	1.87	0.0372	50.2	2.80E-120
<i>k</i> (aboveground): wet mass	−0.000387	0.000342	−1.13	0.259
<i>k</i> (roots): wet mass	0.00128	0.000570	2.25	0.0253

doi:10.1371/journal.pone.0033996.t003




Figure 2. Dry biomass density of roots in the four sites across the Kalahari aridity gradient. The error bars indicate the minimum and maximum data values, unless outliers are present (shown as circles). The black line indicates the median, while the box boundaries are the lower and upper quartiles. Based on a set of 60 soil profiles sampled at each site. doi:10.1371/journal.pone.0033996.g002

Bureau of Meteorology, 1971–2000) from Shakawe, Ghanzi, Tshane, and Bokspits. All necessary permits were obtained (from the Ministry of Environment, Wildlife and Tourism of the Republic of Botswana) to perform the described field studies at these sites.

Field scale surveys of above- and below-ground biomass

The survey of aboveground and belowground biomass took place during the dry season over the course of three field campaigns (June–August) in 2008 (in Shakawe), 2009 (in Bokspits

Table 4. Parameters, a and b , and associated statistics from the fitting of an exponential distribution $r(z) = a e^{-bz}$ to the average root profile in the vertical (z) direction.

Site	a (g m^{-3})	b (m^{-1})	R^2	RMSE (g m^{-3})
	a_L (cm m^{-3})	b_L (m^{-1})	R^2	RMSE (cm m^{-3})
	[95% confidence bounds]	[95% confidence bounds]		
Shakawe	5250 [4833.0, 5668.0]	4.874 [4.425, 5.323]	0.9995	18.8
	7595 [4911, 10280]	1.367 [0.3384, 2.396]	0.9834	229.68
Kuke	540.4 [404.0, 676.7]	0.2413 [−0.334, 0.817]	0.9928	33.5
	5555 [5121, 5988]	0.6726 [0.4773, 0.868]	0.9993	48.52
Tshane	330.0 [271.8, 388.1]	1.636 [1.093, 2.180]	0.9959	8.3
	3946 [3223, 4670]	2.37 [1.71, 3.03]	0.9957	42.54
Bokspits	257.7 [204.8, 310.5]	2.775 [1.974, 3.576]	0.9948	5.0
	7299 [6709, 7889]	2.298 [2.011, 2.584]	0.9991	35.62

The parameters a_L , b_L , and associated statistics from the fitting of an exponential distribution $r_L(z) = a_L e^{-b_L z}$ to the average root density (figure 3) are shown in italic fonts. The distribution is calculated as an average of three plots (20 soil pits per plot) at five depths. doi:10.1371/journal.pone.0033996.t004




Figure 3. Linear root density (average root lengths per volume) across the Kalahari rainfall gradient in Botswana. Kuke site exhibits the highest length/volume readings compared to the other sites. The error bars indicate the minimum and maximum data values, unless outliers are present (shown as circles). The black line indicates the median, while the box boundaries are the lower and upper quartiles. Based on a set of 60 soil profiles sampled at each site.

doi:10.1371/journal.pone.0033996.g003

and Tshane), and 2010 (in Kuke and Tshane). At each of the four sites, three 20 m×20 m plots were established in randomly selected locations within 1000 m of each other. In each plot, the spatial location of each tree and shrub was mapped and plant height, tree stem coordinates, basal diameter, and canopy size were measured. The aboveground tree and shrub biomass was then harvested and weighed. An estimate of aboveground wet biomass in g m^{-2} was calculated by dividing the total aboveground biomass by the plot area (400 m^2).

Root sampling

A variety of methods are commonly used to investigate root structure and function, including microsatellite markers [59], tracer uptake [22,60], ground penetrating radar [61], minirhizotron cameras [62], coring, and excavations using shovels or air pressure systems (e.g., [44,63]). While non-destructive methods (e.g., tracer-based methods) are used to determine the zones of influence [22,60] of plant roots and the function of fine roots [64],

they are not effective at quantifying woody root structure and biomass in that only functional (i.e., active) roots contribute to tracer uptake. Root excavation is the most direct approach to determine root biomass because it samples both active and inactive roots, and allows for direct root mass measurement. Therefore, excavation methods were utilized to quantify below-ground biomass. Each plot was partitioned into a square grid of 20×20 square subplots of 1 m×1 m area. Twenty of these 1 m^2 subplots were randomly selected in each plot (i.e., a total of 60 plots per site). Each of these 1 m^2 subplots were dug to a depth of 110 cm, with a 30 cm-thick surface layer and four subsequent 20 cm-thick layers.

During the excavation of these 1 m^2 soil pits, all woody roots in each soil layer were harvested, cleaned of soil, and weighed; length and diameter were measured with a caliper. Roots finer than 2 mm in diameter were grouped and then weighed together for each soil layer. The same procedure was repeated in all three plots at each of the four study sites for a total of 240 soil pits. In 2009, a subset of both root and branch pieces of various lengths and widths




Figure 4. Root diameter distribution along the soil profile across the Kalahari's rainfall gradient. The error bars indicate the minimum and maximum data values, unless outliers are present (shown as circles). The black line indicates the median, while the box boundaries are the lower and upper quartiles. Based on a set of 60 soil profiles sampled at each site.

doi:10.1371/journal.pone.0033996.g004

were collected at the Tshane site. These samples were oven-dried and reweighed to determine the ratio between dry and wet biomass as explained in section 2.4.

Data Analyses

A relation between dry and wet biomass ($k = \text{wet mass}/\text{dry mass}$) was determined from a subset of roots ($n = 50$), stems ($n = 75$) and branch ($n = 102$) samples collected in the field and weighted both before and after oven drying for 24 hours at 60°C . This subset of samples was used to perform an analysis of covariance (ANCOVA) on the ratio of fresh mass to dry mass of each sample using tissue type (i.e., above ground or below ground biomass) and species as factors and size of the wet sample as a covariate. It was found that, while for above ground biomass the k ratio does not significant vary across the three species here considered (i.e., $k = \text{Boschia albutrunca}$, $\text{Terminalia sericea}$, and Acacia mellifera), significant differences in the k ratios of below ground biomass existed among the same species. However, because the sampling protocol used for belowground biomass consisted of excavating a number of soil pits, and collecting and measuring the root samples in each pit, we were unable to determine to what species each sample belonged to. For this reason, we used the same k ratio for

all above ground biomass samples and another one for all the belowground samples, without accounting for differences among species. The analysis of covariance was then repeated using tissue type (i.e., above ground or below ground biomass) as a factor and size of the wet sample as a covariate. The results (Table 3) show that k is significantly different for aboveground ($k = 1.64$) and belowground ($k = 1.87$) biomass. For aboveground biomass there was a significant interaction ($p = 0.259$) between k and the size of the sample (expressed in terms of wet biomass); however, the slope of the interaction between branch and wet mass is close to zero (see Table 3) and the effect of this interaction on the estimate of aboveground biomass is overall negligible. Conversely for belowground biomass the interaction was not significant ($p = 0.0253$). The above analysis was then used to convert wet biomass to dry biomass with two different values of k for roots and stems or branches.

Plot-level averages of aboveground biomass, belowground biomass, and root-to-shoot ratio were tested for normality using the Jarque-Bera test [65] at the 5% significance level. To determine how rainfall influences allocation, we calculated the correlation coefficient of MAP and each of these variables with the plot as the unit of replication.

Root biomass data were expressed in terms of root density, $r(z)$, (i.e., dry root mass per unit volume of soil) and analyzed to assess changes with depth and across sites. For each of the 20×20 plots we fit an exponential distribution, $r(z) = a \exp[-bz]$, to the data where a and b are two parameters obtained by fitting the cumulative distribution, $R(z) = \int r(u) du$, to the data. Following Schenk and Jackson [52], we expressed root depth in terms of the portions, Z_{50} and Z_{95} , of the soil profile, which comprise 50% and 95% of the total root biomass, respectively. The 50% and 95% percentiles were derived from the fitted equations, while the total belowground biomass for the whole soil profile (in g m^{-2}) was calculated by extrapolation as $R(\infty) = a/b$. The same analysis was then repeated for the vertical profiles of average linear root density, $r_L(z)$, defined as the average root length per unit volume of soil. Based on the linear root density, the depths, $Z_{L,50}$ and $Z_{L,95}$ - containing 50% and 95% of the total root length, respectively - were calculated.

Results

Root biomass decreases with depth at all sites except Kuke, which exhibits a relatively uniform distribution of root biomass in the top 1.10 m (Fig. 2). At all sites, an exponential distribution provides a good fit of the field data ($R^2 = 0.99$ for each of these four sites, see Table 4). The 95% confidence bounds reported in Table 4 show that at Kuke the b parameter is not significantly different from zero, suggesting that at this site the root profile is uniform. Moreover, these bounds show that at all sites the within-site variability was smaller than the variability of a and b among sites. An indicator of root distribution, such as the linear root density (Fig. 3), exhibits a well-defined exponential decrease with depth at all sites ($R^2 > 0.98$ at all sites, see Table 4), while root diameters do not exhibit a well-defined dependence on depth (Fig. 4).

Overall, tree roots in the Kalahari are relatively shallow with Z_{50} depths (i.e., 50% of root biomass is shallower than Z_{50}) ranging between 0.14 and 0.43 m (Table 5), while the Z_{95} depths (i.e., 95% of the roots are shallower than Z_{95}) range between 0.61 and 1.88 m. The only exception to this is the Kuke site, where the root profile in the top 1.10 m was relatively uniform. Therefore, at this site the estimation of root depth by fitting an exponential function provided values of Z_{50} and Z_{95} that were much larger than the 1.10 m depth explored in our excavations. Thus, these values of Z_{50} and Z_{95} are not reported in Table 5. Linear root density profiles (e.g., $Z_{L,50}$) exhibit relatively shallow distributions at all sites. The rooting depths reported in Table 5, however, do not show any clear relationship with MAP since the deepest root systems were found in areas in the middle of the transect (Kuke and Tshane) rather than at either end. Similarly, neither the interannual variability nor the minimum values of precipitation (see Table 2) could explain root depth variability among sites.

The Jarque-Bera test indicated that above- and belowground biomass and R:S ratio were not normally distributed, but the log transforms of these variables are. Belowground biomass decreases with decreasing MAP (Table 5) with relatively large variations among sample plots, particularly at the wetter sites (Fig. 5). The correlation coefficient of MAP and log-transformed belowground biomass was 0.90 ($p < 0.01$) for a set of 3 plot replicates at each site. The same relationship can be also found in the extrapolated values of total belowground biomass, $R(\infty)$. Nonetheless, the plots used in this study did not show any clear dependence of aboveground woody biomass on MAP (Fig. 5). There was no significant correlation between log-transformed aboveground biomass and MAP ($p = 0.66$) for a set of 3 plot replicates at each site. However,

Table 5. Z_{50} and Z_{95} root depths calculated assuming an exponential profile of root biomass.

Site	Z_{50} (m)	Z_{95} (m)	Root Biomass in Top 1.10 m (g m^{-2})	Extrapolated Total Root Biomass, $R(\infty)$ (g m^{-2})	Above-ground Biomass (g m^{-2})	Basal Area ($\text{m}^2 \text{ ha}^{-1}$)	R:S (top 1.1 m)	$Z_{L,50}$ (m)	$Z_{L,95}$ (m)
Shakawe	0.14	0.61	1070.76 ± 676.9	1078.03	438.65 ± 272.46	3.6	3.10 ± 2.91	0.26	1.16
Kuke	n/a	n/a	536.16 ± 100.34	n/a	842.08 ± 236.53	6.4	0.64 ± 0.08	0.90	3.89
Tshane	0.43	1.88	163.75 ± 40.44	201.71	513.37 ± 68.20	2.1	0.30 ± 0.03	0.31	1.32
Bokspits	0.25	1.08	91.89 ± 28.75	92.86	556.50 ± 116.76	1.9	0.17 ± 0.03	0.28	1.25

Measured dry belowground biomass (top 1.1 m), extrapolated total root biomass (i.e., $R(\infty)$, see Methods section), dry above ground biomass, basal area, root-to-shoot ratio (R:S, based on dry root biomass in the top 1.10 m), and depths, $Z_{L,50}$ and $Z_{L,95}$ of soil column containing 50% and 95% of the total root length, respectively (assuming exponential profile of linear root density). Biomass variability among plots expressed as \pm standard deviation calculated for a set of 3 plot replicates at each site.
doi:10.1371/journal.pone.0033996.t005




Figure 5. Measured average woody plant biomass (above and belowground) per unit area (top 1.1 m) across the Kalahari transect aridity gradient. Inset: ratio of below to above ground biomass. The error bars represent \pm standard deviation calculated for a set of 3 plot replicates at each site.

doi:10.1371/journal.pone.0033996.g005

in fire-prone areas above ground biomass is difficult to estimate because of its dependence on the time since last disturbance. The ratio of below- to aboveground biomass has a clear positive relationship with rainfall, with a correlation between MAP and log-transformed R:S ratio of 0.89 ($p < 0.001$). Shakawe exhibited the highest R:S ratio and Bokspits the lowest (Fig. 5-inset), indicating that woody plants at the more mesic (northern) sites allocate more of their biomass to roots relative to woody plants in the dryer (southern) sites.

Discussion

We found that there is a positive relationship between belowground biomass and mean annual rainfall, which is consistent with previous studies in other systems [5,52]. However, because there is no clear relationship between above ground biomass and annual rainfall at our sites, the R:S ratio increases with rainfall (Fig. 5), which is inconsistent with the results by Schulze et al. [5].

According to the functional equilibrium hypothesis of plant growth [48] plants should allocate more carbon to belowground biomass in areas with stronger limitations in belowground resources (e.g., soil moisture), while they should allocate more carbon to above ground biomass in areas that are more limited by above ground resources (e.g., light). Thus, this hypothesis predicts decreasing R:S with increasing MAP. For studies on biomass allocation between below- and aboveground biomass in grasses and in potted plants some authors [48,50,66,67] have shown that the R:S ratios increase with aridity (i.e., decrease with increasing MAP). Our data for woody species in the Kalahari directly contradict the predictions of the functional equilibrium hypothesis. The inconsistency between our results and those from studies of potted plants and herbaceous vegetation suggests that different vegetation growth forms respond differently to climatic and environmental drivers (e.g. [5,21,33]).

In recent years, theories of root distribution [37,38,39,40,41,68] have provided a framework to determine “optimal root profiles”, i.e., the vertical root distributions that optimize the use of soil moisture by water limited vegetation for given climate and soil properties. According to these theories, optimal root profiles in semiarid environments have the following characteristics: 1) the root density profile decreases exponentially with depth (e.g., [37]), and 2) root depth (e.g., Z_{50} and Z_{95} or $Z_{L,50}$ and $Z_{L,95}$) increases with MAP due to the fact that rainwater infiltrates deeper into the soil column in mesic environments [11,37,38,39] and plants grow more roots where there are more resources to take up, a phenomenon known as “hydrotropism” (e.g., [47,69]).

Our data show that, in the Kalahari, the linear root density of woody species – an indicator of roots’ ability to take up water and other soil resources (e.g., [35]) – decreases exponentially with depth at all sites (Fig. 3), but values of $Z_{L,50}$ and $Z_{L,95}$ calculated from these fits do not increase with increasing MAP. Indeed the lowest values of $Z_{L,50}$ and $Z_{L,95}$ occur at Shakawe, the wettest site. The same is observed with Z_{50} and Z_{95} calculated from average root mass profiles, though in this case there is one site (Kuke) with a root mass profile that does not exponentially decrease with depth (Fig. 2c) and for which Z_{50} and Z_{95} values cannot be calculated. Thus, our data do not support the predictions of optimality theories of root distribution.

Using data from different biomes and across a wide precipitation range, Schenk and Jackson [21] found that for grasses and forbs, the rooting depth exhibits a strong positive relationship with MAP. However, this relationship with MAP was less clear for trees and shrubs [21]. This study shows that woody vegetation does not exhibit an increase in rooting depth with MAP.

Overall, our data show that woody vegetation in the Kalahari has relatively shallow roots, despite the sandy texture of Kalahari soils, which would be expected to favor plants with deep roots because of the deeper and faster infiltration typical of soils with




Figure 6. Left: Fire frequency from 2000 to 2011 calculated from the MODIS Burned Area Product (MCD45) [72] in yr^{-1} . White areas experienced no fires during this period. **Right: Average fire frequency (in yr^{-1}) calculated along a longitudinal transect (21.3°) using a moving box of approximately 100×100 km.**
doi:10.1371/journal.pone.0033996.g006

coarse texture [30]. The vegetation we sampled has 50% of its root biomass in the top 0.14 to 0.43 m of the soil column (and 95% in the top 0.61 or 1.88 m), except for the Kuke site, which exhibits deeper roots (Table 5).

Given the failure of existing theories to explain this new and unique dataset, alternative explanations of root distribution must be explored. We hypothesize that our data could emerge from differing strategies of water and carbon utilization and storage along the Kalahari Transect. In the Kalahari, as in savannas worldwide, woody vegetation must contend with ongoing threats of fire and drought. In the northern portion of the transect, rainfall is higher, and consequently, the incidence of fire during the dry season is also higher (see Fig. 6 and e.g., [70,71]). In the southern portion of the transect precipitation is lower, so there is insufficient biomass to sustain significant fires (Fig. 6). Thus, different strategies are appropriate at different points along the rainfall gradient. We hypothesize that in the more mesic portion of the transect a strategy that stores carbon is better suited to the climatic and fire conditions. In this strategy, plants invest in significant belowground root systems to provide ample storage for water and carbon. Although the large root system brings with it significant metabolic costs, the ability for storage that it entails means that loss of the aboveground

structure in a fire can be compensated for by quick growth from the belowground stores. Due to the additional cost of respiration required by this large belowground storage system, the plant may be in carbon deficit in some years (especially during droughts), but will still have excess metabolic carbon in most years. In contrast, the lack of fire and the more frequent drought occurrence in the dry portion of the transect favors a second strategy. In this strategy, the root system is not used for carbon and water storage, but is almost exclusively used for foraging of soil resources, mainly water. Because an extensive root system cannot be sustained in these water-limited environments, the plants have a lower R:S ratio. Large belowground biomass is not required under this strategy because no storage to be used for re-growth of aboveground biomass is required where fire frequency is low. In this strategy, a small root system can harvest ample water to support the organism, as long as it is also shallow and spatially extensive, and does not provide a large metabolic burden, especially in areas where the risk of drought, and therefore carbon deficit, is high.

This hypothesis of life-history strategies explains several important features of our data, including why the R:S ratio increases with increasing MAP (due to the need for additional storage as the risk of fire increases), and why rooting depth is

shallow in the dry portion of the transect (so roots can easily access soil moisture from small precipitation events), while not requiring that rooting depth be deeper in wetter areas.

Acknowledgments

We acknowledge our field assistants, O. Mathata, U. Mathata, David Rachal, Mowkaledi Mafa, Molly O'Connor, Ming Lu, Matthew Patterson, Christine Bonthius, Jeremy Chen, Rodrigo Munoz-Rogers and teacher-researchers, Heather Riffel, Richard Wellbeloved-Stone, and Ashley

Pollard. A special thanks to Mr. Johannes Martin for allowing us to use his land for this research and the Botswana Department of Forestry and Range Resources for their invaluable help.

Author Contributions

Conceived and designed the experiments: PD GSO KKC SR. Performed the experiments: MT FCO DOP. Analyzed the data: MT AB KD. Wrote the paper: MT PD AB.

References

- Scholes RJ, Walker BH (1993) An African savanna, Synthesis of the Nylsvley Study. New York: Cambridge University Press.
- Scholes RJ, Archer SR (1997) Tree-grass interactions in Savannas. *Annual Review Ecology and Systematics* 28: 517–544.
- Sankaran M, Hanan PN, Scholes RJ, Ratnam J, Augustine DJ, et al. (2005) Determinants of woody vegetation in African Savannas. *Nature* 438: 846–849.
- Canadell J, Jackson RB, Ehleringer JR, Mooney HA, Sala OE, et al. (1996) Maximum rooting depth for vegetation types at the global scale. *Oecologia* 108: 583–595.
- Schulze ED, Mooney HA, Sala OE, Jobbagy E, Buchmann N, et al. (1996) Rooting depth, water availability, and vegetation cover along an aridity gradient in Patagonia. *Oecologia* 108: 503–511.
- Gill RA, Jackson RB (2000) Global patterns of root turnover for terrestrial ecosystems. *New Phytologist* 147: 13–31.
- Skarpe C (1992) Dynamics of savanna ecosystems. *Journal of Vegetation Science* 3: 293–300.
- House JI, Archer S, Breshears DD, Scholes RJ (2003) Conundrums in mixed woody–herbaceous systems. *Journal of Biogeography* 30: 1763–1777.
- D'Odorico P, Laio F, Ridolfi L (2006a) A probabilistic analysis of fire-induced tree-grass coexistence in savannas. *American Naturalist* 167: E79–E87.
- D'Odorico P, Porporato A (2006b) *Dryland ecohydrology*. Dordrecht: Springer.
- D'Odorico P, Caylor KK, Okin GS, Scanlon TM (2007) On soil moisture–vegetation feedbacks and their possible effects on the dynamics of dryland ecosystems. *Journal of Geophysical Research* 112: G04010. doi: 10.1029/2006JG000379.
- Wang L, D'Odorico P, Okin GS, Macko SA (2009) Isotope composition and anion chemistry of soil profiles along the Kalahari Transect. *Journal of Arid Environments* 73: 480–486.
- Kimmins JP (2004) *Forest Ecology, a foundation for sustainable forest management and environmental ethics in forestry* (3rd ed.). London: Pearson Education Inc..
- Jackson RB, Mooney HA, Schulze ED (1997) A global budget for fine root biomass, surface area, and nutrient contents. *Proceedings of the National Academy of Sciences* 94: 7362–7366.
- Greene RSB (1992) Soil physical-properties of three geomorphic zones in a semiarid Mulga woodland. *Australian Journal of Soil Research* 30: 55–69.
- Sankaran M, Ratnam J, Hanan NP (2004) Tree–grass coexistence in savannas revisited – insights from an examination of assumptions and mechanisms invoked in existing models. *Ecology Letters* 7: 480–490.
- Caylor KK, D'Odorico P, Rodriguez-Iturbe I (2006) On the ecohydrology of structurally heterogeneous semi arid landscapes. *Water Resource Research* 42: W07424. doi: 10.1029/2005WR004683.
- Breshears DD, Barnes EJ (1999) Interrelationships between plant functional types and soil moisture heterogeneity for semiarid landscapes within the grassland/forest continuum, a conceptual model. *Landscape Ecology* 14: 465–478.
- Caylor KK, Shugart HH, Dowty PR, Smith TM (2003) Tree Spacing along the Kalahari Transect in southern Africa. *Journal of Arid Environments* 54: 281–296.
- Barbier N, Couteron P, Lefever R, Deblauwe V, Lejeune O (2008) Spatial decoupling of facilitation and competition at the origin of gapped vegetation patterns. *Ecology* 89: 1521–1531.
- Schenk JH, Jackson RB (2002b) Rooting depths, lateral root spreads and below-ground/above-ground allometries of plants in water-limited ecosystems. *Journal of Ecology* 90: 480–494.
- Hartle RT, Fernandez GCJ, Nowak RS (2006) Horizontal and vertical zones of influence for root systems of four Mojave Desert shrubs. *Journal of Arid Environments* 64: 586–603.
- Caylor KK, Shugart HH, Rodriguez-Iturbe I (2005) Tree canopy effects on simulated water stress in southern African savannas. *Ecosystems* 8: 17–32.
- Hipondoka MHT, Versfeld WD (2006) Root system of *Terminalia sericea* shrubs across rainfall gradient in a semi-arid environment of Etosha National Park, Namibia. *Ecological Indicators* 6: 516–524.
- Norby RJ, Jackson RB (2000) Root dynamics and global change, seeking an ecosystem perspective. *New Phytologist* 147: 3–12.
- Burgess SSO, Adams MA, Turner NC, Ong CK (1998) The redistribution of soil water by tree root systems. *Oecologia* 115: 306–311.
- Horton JL, Hart SC (1998) Hydraulic lift, A potentially important ecosystem process. *Tree* 13: 232–235.
- Ludwig F, Dawson TE, Kroon H, Berendse F, Prins HHT (2002) Hydraulic lift in *Acacia tortilis* trees on and East African savanna. *Oecologia* 134: 293–300.
- Walter H (1971) *Ecology of tropical and subtropical vegetation*. Edinburgh: Oliver & Boyd.
- Sala OE, Lauenroth WK, Golluscio RA (1997) Plant functional types in temperate semi arid regions Plant Functional Types. Smith TM, Shugart HH, Woodward FI, eds. 217–233, Cambridge: Cambridge University Press.
- Kulmatiski A, Beard KH, Verweij RJT, February EC (2010) A depth-controlled tracer technique measures vertical, horizontal and temporal patterns of water use by trees and grasses in a subtropical savanna. *New Phytologist* 188: 199–209. doi: 10.1111/j.1469-8137.2010.03338.x.
- Le Roux X, Bariac T, Mariotti A (1995) Spatial partitioning of the soil water resource between grass and shrub components in a West African humid savanna. *Oecologia* 104: 147–155.
- Hipondoka MHT, Arinabar AR, Chirara C, Lihavha M, Macko MA (2003) Vertical distribution of tree and grass roots in Southern African ecosystems, niche differentiation or competition? *Journal of Arid Environments* 54: 319–325.
- Mommer L, van Ruijven J, de Caluwe H, Smit-Tiekstra AE, Wagemaker CAM, et al. (2010) Unveiling below-ground species abundance in a biodiversity experiment: a test of vertical niche differentiation among grassland species. *Journal of Ecology* 98: 1117–1127.
- Feddes RA, Holger H, Bruen M, Dawson T, de Rosnay P, et al. (2001) Modeling root-water uptake in hydrological and climate models. *Bulletin of American Meteorology Society* 82: 2797–2809.
- Bonan G (2002) *Ecological Climatology*. Cambridge: Cambridge University Press.
- Laio F, D'Odorico P, Ridolfi L (2006) An analytical model to relate the vertical root distribution to climate and soil properties. *Geophysical Research Letters* 33: L18401. doi: 10.1029/2006GL027331.
- Collins DBG, Bras RL (2007) Plant rooting strategies in water-limited ecosystems. *Water Resources Research* 43: W06407. doi: 10.1029/2006WR005541.
- Guswa AJ (2008) The influence of climate on root depth, A carbon cost-benefit analysis. *Water Resources Research* 44: W02427. doi:10.1029/2007WR006384.
- Guswa AJ (2010) Effect of plant uptake strategy on the water–optimal root depth. *Water Resources Research* 46: W09601. doi:10.1029/2010WR009122.
- Preti F, Dani A, Laio F (2010) Root profile assessment by means of hydrological, pedological and above-ground vegetation information for bio-engineering purposes. *Ecological Engineering* 36: 305–316.
- Van Wijk MT (2011) Understanding plant rooting patterns in semi-arid systems, an integrated model analysis of climate, soil type and plant biomass. *Global Ecology and Biogeography* 20(2): 331–342. doi: 10.1111/j.1466-8238.2010.00601.x.
- Schenk HJ (2008) The shallowest possible water extraction profile, A null model for global root distributions. *Vadose Zone Journal* 7: 1119–1124.
- Waisel Y, Eshel A, Kafkafi U (1991) *Plant roots, the hidden half*. New York: Marcel Dekker Inc.
- Grace J, San Jose J, Meir P, Miranda HS, Montes RA (2006) Productivity and carbon fluxes of tropical savannas. *Journal of Biogeography* 33: 387–400.
- Schutz AEN, Bond WJ, Cramer MD (2009) Juggling carbon, allocation patterns of a dominant tree in a fire-prone savanna. *Oecologia* 160: 235–246.
- Pooter H, Nagel O (2000) The role of biomass allocation in the growth response of plants to different levels of light, CO₂, nutrients and water, a quantitative review. *Australian Journal of Plant Physiology* 27: 595–607.
- Brouwer R (1962) Nutritive influences on the distribution of dry matter in the plant. *Netherlands Journal of Agricultural Sciences* 10: 361–376.
- Klepper B (1991) Root-shoot relationships. In: *Plant roots, the hidden half* Waisel Y, Eshel A, Kafkafi U, eds. 265–286, New York: Marcel Dekker Inc.
- Hui DF, Jackson RB (2006) Geographical and interannual variability in biomass partitioning in grassland ecosystems, a synthesis of field data. *New Phytologist* 169: 85–93.
- Jackson RB, Canadell J, Ehleringer JR, Mooney HA, Sala OE, et al. (1996) A global analysis of root distributions for terrestrial biomes. *Oecologia* 108: 389–411.
- Schenk JH, Jackson RB (2002a) The global biogeography of roots. *Ecological Monographs* 72: 311–328.

53. Ringrose S, Matheson W, Wolski P, Huntsman-Mapila P (2003) Vegetation cover trends along the Botswana Kalahari transect. *Journal of Arid Environments* 54: 297–317.
54. Wang L, D'Odorico P, Ringrose S, Coetzee S, Macko SA (2007) Biogeochemistry of Kalahari Sands. *Journal of Arid Environments* 71: 259–27.
55. Koch GW, Vitousek PM, Steffen WL, Walker BH (1995) Terrestrial transects for global change research. *Plant Ecology* 121: 53–65.
56. Shugart HH, Macko SA, Lesolle P, Szuba TA, Mukelebai MM, et al. (2004) The Safari 2000- Kalahari Transect wet season campaign of year 2000. *Global Change Biology* 10: 273–280.
57. Porporato A, Laio F, Ridolfi L, Caylor KK, Rodriguez-Iturbe I (2003) Soil moisture and plant stress dynamics along the Kalahari precipitation gradient. *Journal of Geophysical Research* 108: 4127. doi: 10.1029/2002JD002448.
58. Scholes RJ, Dowty PR, Caylor KK, Parsons DB, Frost PGH, et al. (2002) Trends in savanna structure and composition along an aridity gradient in the Kalahari. *Journal of Vegetation Science* 13: 419–428.
59. Saari SK, Campbell CD, Russell J, Alexander AJ, Anderson IC (2005) Pine microsatellite markers allow roots and ectomycorrhizas to be linked to individual trees. *New Phytologist* 165: 295–304.
60. Casper BB, Schenk HJ, Jackson JB (2003) Defining a plant's belowground zone of influence. *Ecology* 84: 2313–2321.
61. Butnor JR, Doolittle JA, Kress L, Cohen S, Johnsen KH (2001) Use of ground-penetrating radar to study tree roots in the southeastern United States. *Tree Physiology* 21: 1269–1278.
62. Cheng W, Coleman DC, Box JE, Jr. (1990) Root dynamics, production and distribution in agroecosystems on the Georgia Piedmont using minirhizotrons. *Journal of Applied Ecology* 27: 592–604.
63. Nadezhkina N, Cermak J (2003) Instrumental methods for studies of structure and function of root systems of large trees. *Journal of Experimental Botany* 54: 1511–1521.
64. Göttlicher SG, Taylor AFS, Grip H, Betson NR, Valinger E, Höglberg MN, et al. (2007) The lateral spread of tree root systems in boreal forests, Estimates based on ¹⁵N uptake and distribution of sporocarps of ectomycorrhizal fungi. *Forest Ecology and Management* 255: 75–81.
65. Jarque CM, Bera AK (1987) A test for normality of observations and regression residuals. *International Statistical Review* 55: 163–172.
66. Fan JW, Wang K, Harris H, Zhong HP, Hu ZM, et al. (2009) Allocation of vegetation biomass across a climate related gradient on the grasslands of Inner Mongolia. *Journal of Arid Environments* 73: 521–528.
67. Boutraa T (2010) Growth performance and biomass partitioning of the desert shrub *Calotropis procera* under water stress conditions. *Research Journal of Agriculture and Biological Sciences* 6: 20–26.
68. Van Wijk MT, Bouten W (2001) Towards understanding tree root profiles, simulating hydrologically optimal strategies for root distribution. *Hydrology and Earth System Science* 5: 629–644.
69. Schenk JH, Jackson RB (2005) Mapping the global distribution of deep roots in relation to climate and soil characteristics. *Geoderma* 126: 129–140.
70. Du Toit JT, Rogers KH, Biggs HC (2003) The Kruger experience, ecology and management of savanna heterogeneity. Washington DC: Island Press.
71. Hely C, Caylor KK, Dowty P, Alleaume S, Swap RJ, et al. (2007) A temporally explicit production efficiency model for fuel load allocation in Southern Africa. *Ecosystems* 10: 1116–1132.
72. Roy DP, Boschetti L, Justice CO, Ju J (2008) The Collection 5 MODIS Burned Area Product - Global Evaluation by Comparison with the MODIS Active Fire Product. *Remote Sensing of Environment* 112: 3690–370.