

Too few, too many, or just right? Optimizing sample sizes for population-level inferences in animal tracking projects

Inês Silva^{1,*}, Christen H. Fleming^{2,3}, Michael J. Noonan^{4,5,6}, William F. Fagan⁷, Justin M. Calabrese^{1,7,8}

¹Center for Advanced Systems Understanding (CASUS), Helmholtz-Zentrum Dresden-Rossendorf (HZDR), Görlitz, Germany;

²Department of Biology, University of Central Florida, Orlando, Florida, USA;

³Smithsonian National Zoo and Conservation Biology Institute, Virginia, USA;

⁴Department of Biology, University of British Columbia Okanagan, Kelowna, Canada;

⁵Okanagan Institute for Biodiversity, Resilience, and Ecosystem Services, University of British Columbia Okanagan, Kelowna, Canada;

⁶Department of Computer Science, Math, Physics, and Statistics, University of British Columbia Okanagan, Kelowna, Canada;


⁷Department of Biology, University of Maryland, Maryland, USA;

⁸Department of Ecological Modelling, Helmholtz Centre for Environmental Research – UFZ, Leipzig, Germany


***Corresponding author:** Inês Silva, **E-mail:** i.simoaes-silva@hzdr.de, imss.silva@gmail.com

S3. Single combination plots

18 Home range estimation



19
20 **Figure S3.1. Relative error (%) in home range estimates (AKDE) as a function of population sample size (m), ranging from 2 to 50 simulated African**
21 **buffalos (*Syncerus caffer*).** Each vertical facet corresponds to one of five sampling durations: 2 months, 4 months, 1 year, 3 years, and 9 years. The
22 plot shows mean estimates based on a **single** combination of individuals per m (no resampling); estimates in **blue** are within the error threshold and
23 in **red** if estimates fall outside of it. Horizontal dotted lines mark the $\pm 5\%$ error threshold.



24
25 **Figure S3.2. Relative error (%) in home range estimates (AKDE) as a function of population sample size (m), ranging from 2 to 50 simulated Mongolian**
26 **gazelles (*Procapra gutturosa*).** Each vertical facet corresponds to one of five sampling durations: 2 months, 4 months, 1 year, 3 years, and 9 years. The
27 plot shows mean estimates based on a **single** combination of individuals per m (no resampling); estimates in **blue** are within the error threshold and
28 in **red** if estimates fall outside of it. Horizontal dotted lines mark the $\pm 5\%$ error threshold.

Speed & distance estimation




Figure S3.3. Relative error (%) in speed estimates (CTSD) as a function of *population* sample size (m), ranging from 2 to 50 simulated African buffalos (*Syncerus caffer*). Each vertical facet corresponds to one of five sampling intervals: 4 hours, 3 hours, 2 hours, 1 hour, and 30 minutes. The plot shows mean estimates based on a single combination of individuals per m (no resampling); estimates in blue are within the error threshold and in red if estimates fall outside of it. Horizontal dotted lines mark the $\pm 5\%$ error threshold.




Figure S3.4. Relative error (%) in speed estimates (CTSD) as a function of *population* sample size (m), ranging from 2 to 50 simulated Mongolian Gazelles (*Procapra gutturosa*). Each vertical facet corresponds to one of five sampling intervals: 4 hours, 3 hours, 2 hours, 1 hour, and 30 minutes. The plot shows mean estimates based on a **single** combination of individuals per m (no resampling); estimates in blue are within the error threshold and in red if estimates fall outside of it. Horizontal dotted lines mark the $\pm 5\%$ error threshold.