

Supplementary Material

Sensor-Family Composition of the In-Situ Archive Changes the Apparent Bias and RMSD of Satellite Soil-Moisture Products

Chang-Hwan Park
 Department of Civil Systems Engineering, Ajou University
 Suwon, Republic of Korea
 Department of Civil Systems Engineering, Ajou University in Tashkent
 Tashkent, Uzbekistan

July 8, 2026

S1 Archive composition and per-reference residuals

Table S1 gives the full manufacturer-family composition of the retained 2020–2023 ISMN archive and the mean ISMN–reference residual for each family against all three reference products, with 95 % bootstrap confidence intervals. The HydraProbe-family ERA5-Land CI does not overlap the Decagon/METER TDR CI, and the family swing ($0.17 \text{ m}^3 \text{ m}^{-3}$) is an order of magnitude larger than the within-family disagreement among references.

Table S1: ISMN 2020–2023 composition and mean ISMN–reference residual ($\text{m}^3 \text{ m}^{-3}$) by manufacturer family, 95 % bootstrap CI (5,000 resamples). Residual means use the valid-coverage matchup set; the composition count and the ERA5 residual count differ because not every station-sensor has valid coverage in every product (e.g., 1,231 of 1,274 HydraProbe-family sensors have valid ERA5-Land coverage).

Sensor family	n	Share (%)	ERA5-Land	SMAP L3.E	SMAP-SMI
HydraProbe family	1,274	78.4	$-0.097 [-0.103, -0.091]$	$-0.025 [-0.035, -0.016]$	$-0.037 [-0.043, -0.032]$
Decagon/METER TDR	95	5.8	$+0.068 [+0.028, +0.109]$	$+0.006 [-0.041, +0.057]$	$-0.074 [-0.103, -0.042]$
Campbell CS6xx	65	4.0	$-0.065 [-0.088, -0.042]$	$+0.019 [-0.001, +0.039]$	$-0.024 [-0.050, +0.005]$
ThetaProbe	59	3.6	$-0.039 [-0.061, -0.017]$	$+0.001 [-0.025, +0.027]$	$-0.001 [-0.028, +0.025]$
TRIME	2	0.1	—	—	—
other	130	8.0	$-0.056 [-0.079, -0.031]$	$+0.003 [-0.026, +0.033]$	$-0.022 [-0.046, +0.002]$
Total	1,625	100.0			

S2 Per-archive dielectric-free CRNS attribution

The pooled cross-check reported in the main text (92 CRNS↔ISMN pairs, Welch’s $t = +4.07$, $p = 2.3 \times 10^{-4}$) decomposes by archive as follows. *COSMOS-USA* (reprocessed CRNS–ERA5-Land bias table of Zheng et al. [1], matched within 0.2° , $n = 56$): HydraProbe $+0.142 \text{ m}^3 \text{ m}^{-3} [+0.114, +0.170]$ vs non-HydraProbe $-0.067 [-0.174, +0.037]$; Welch’s $t = +3.61$, $p = 1.4 \times 10^{-3}$. *COSMOS-Europe* [2] (matched within 0.1° , $n = 31$): HydraProbe $+0.058 [+0.038, +0.078]$ vs non-HydraProbe $+0.010 [-0.016, +0.040]$; $t = +2.60$, $p = 0.019$. *COSMOS-UK* [3] on-site CRNS–TDT (same-footprint

dielectric probe, $n = 49$): $+0.035$ [$+0.010, +0.061$], excluded from the pooled within-archive family contrast because its design has no non-HydraProbe channel. *CosmOz Australia* [4] ($n = 5$): CRNS–HydraProbe $+0.162$ ($n = 2$), CRNS–Campbell CS6xx -0.043 ($n = 3$)—directionally consistent but too small for an independent test. Leave-one-archive deletion: dropping USA gives $t = +3.91$, $p = 6.3 \times 10^{-4}$; dropping Europe $t = +4.05$, $p = 3.7 \times 10^{-4}$; dropping Australia $t = +3.68$, $p = 7.8 \times 10^{-4}$. Figure S1 summarizes the archive-level contrasts.

The COSMOS-USA row is a synthesized residual: the Zheng et al. CRNS–ERA5-Land bias is differenced against the station-matched ISMN–ERA5-Land residual from this work. The two ERA5-Land overlays are not redistributed as common-cell grids in this compact review package, so the USA row should be interpreted as a supporting rather than sole-identifying contrast. The USA-excluded leave-one-archive result above is therefore the conservative check used in the main text.




Figure S1: Per-archive dielectric-free CRNS attribution. Each row is the mean (CRNS – dielectric probe) residual for one archive-family comparison; error bars are 95 % bootstrap CI. The neutron reference is systematically wetter than the HydraProbe family and closer to non-HydraProbe families across three continents; Australia is shown for direction only.

S3 Near-surface constraint for the paired-sensor test

The main-text paired test uses $n = 21$ stations that host both a HydraProbe-family and a non-HydraProbe sensor in the same ISMN grid cell; both members are in the retained near-surface layer (depth-from ≤ 0.15 m). Table S2 reports the paired family contrast used in the main text. The inference is therefore restricted to co-located near-surface pairs.

Table S2: Paired-sensor offset (non-HydraProbe – HydraProbe) for co-located near-surface pairs.

Pair subset	pairs	mean diff (m ³ m ⁻³)	t	p
All co-located pairs	21	$+0.075$ [$+0.031, +0.119$]	$+3.19$	4.7×10^{-3}

S4 Real 2021 SMAP temporal validation check

The main letter focuses on mean-state bias and RMSD. To connect the family-composition result to standard satellite calibration/validation language, we also computed a real overpass-matched daily matchup for 2021 using raw ISMN `.stm` hourly files and SMAP L3 SM_P descending/AM granules. The filters were ISMN quality flag "G", depth-from ≤ 0.15 m, recommended-quality SMAP retrievals, and at least 20 matched days per station-sensor. The resulting table contains 785 station-sensors and 78,600 matched station-days.

Table S3: Real 2021 overpass-matched temporal metrics by manufacturer family against SMAP L3 SM_P (AM). Bias is ISMN minus SMAP; values are station-sensor medians; brackets give 95 % network-cluster bootstrap CI. ThetaProbe ($n = 3$) is shown for completeness only.

Family	Stations	Bias	ubRMSE	R	Anomaly R
HydraProbe	637	−0.032 [−0.099, −0.017]	0.062 [0.051, 0.074]	0.317 [0.150, 0.443]	0.329 [0.088, 0.465]
Campbell CS6xx	58	−0.107 [−0.151, +0.099]	0.058 [0.032, 0.064]	0.166 [0.087, 0.622]	0.140 [−0.036, 0.657]
Decagon/METER TDR	57	+0.058 [+0.031, +0.139]	0.048 [0.028, 0.068]	0.622 [0.441, 0.647]	0.423 [0.264, 0.635]
other	30	−0.071 [−0.085, −0.033]	0.078 [0.056, 0.080]	0.055 [0.001, 0.436]	0.059 [−0.016, 0.283]
ThetaProbe	3	+0.195	0.030	0.639	0.608




Figure S2: Real 2021 overpass-matched SMAP L3 SM_P (AM) temporal validation by sensor family. Bars are family medians and error bars are network-cluster 95 % CI.

The HydraProbe-versus-non-HydraProbe contrast is -0.032 versus $-0.036 \text{ m}^3 \text{ m}^{-3}$ for bias ($p = 9.8 \times 10^{-3}$), 0.062 versus $0.057 \text{ m}^3 \text{ m}^{-3}$ for ubRMSE ($p = 8.2 \times 10^{-4}$), 0.317 versus 0.273 for R ($p = 0.70$), and 0.329 versus 0.179 for anomaly R ($p = 0.055$). Because this check uses the 36-km SMAP SM_P product, one year, AM overpasses only, and point-to-grid matchups, it is interpreted as a standard-metric diagnostic rather than as a replacement for the mean-state SPL3SMP_E result in the main text. The pooled non-HydraProbe bias masks family heterogeneity: Decagon/METER TDR is wetter than HydraProbe ($+0.058$ versus $-0.032 \text{ m}^3 \text{ m}^{-3}$), whereas Campbell CS6xx and other records are drier (-0.107 and $-0.071 \text{ m}^3 \text{ m}^{-3}$).

S5 Temporal-window invariance

Because SMAP L3_E was extracted for 2023 only while ERA5-Land and SMAP-SMI span 2020–2023, we recomputed the ERA5-Land swvl1 (0–7 cm) grand mean for (i) the full 2020–2023 window and (ii) 2023 only, using an identical 0.1° nearest-neighbour protocol. The HydraProbe-family mean residual changes by $0.004 \text{ m}^3 \text{ m}^{-3}$ (3.9 %) between the two windows; all five families change by $\leq 0.012 \text{ m}^3 \text{ m}^{-3}$; and the driest→wettest family ranking (HydraProbe → Decagon/METER → other

→ ThetaProbe → Campbell CS6xx) is identical in both windows. The reported family ordering is therefore not an artifact of the ERA5 2020–2023 versus SMAP L3_E 2023-only window choice.

S6 Soil-texture control

Soil organic carbon (SOC; g kg^{-1}) was log-transformed before regression. A single ordinary-least-squares regression of the pooled three-reference mean residual on $\log_{10}(\text{SOC})$ gives a slope of $-0.078 \text{ m}^3 \text{ m}^{-3}$ per decade ($p = 1.7 \times 10^{-8}$, $n = 786$). At face value this reads as an organic-matter effect, but it is not robust to texture control. In a linear mixed-effects model with a random intercept for ISMN network and fixed-effect covariates for SoilGrids 2.0 [5] clay fraction and bulk density, sensor depth-from, and $|\text{latitude}|$, implemented with Statsmodels [6], the $\log_{10}(\text{SOC})$ coefficient collapses in all three references (ERA5-Land -0.029 , $p = 0.11$; SMAP L3_E -0.009 , $p = 0.77$; SMAP-SMI $+0.030$, $p = 0.08$), while the clay coefficient is highly significant in every reference ($p < 10^{-9}$). A leave-one-network jackknife of the OLS slope stays negative across all networks (-0.074 ± 0.011 ; range -0.096 to -0.034). The apparent organic-matter dependence is thus confounded with clay fraction and with HydraProbe deployment geography, consistent with a sensor-family/texture confounder rather than a direct organic-matter effect; the sensor-family ordering itself is unchanged by these controls.

References

- [1] Y. Zheng *et al.*, "Evaluation of reanalysis soil moisture products using cosmic ray neutron sensor observations across the globe," *Hydrol. Earth Syst. Sci.*, vol. 28, pp. 1999–2022, 2024, doi:10.5194/hess-28-1999-2024.
- [2] H. R. Bogaen *et al.*, "COSMOS-Europe: a European network of cosmic-ray neutron soil moisture sensors," *Earth Syst. Sci. Data*, vol. 14, pp. 1125–1151, 2022, doi:10.5194/essd-14-1125-2022.
- [3] H. M. Cooper *et al.*, "Soil moisture estimates from the UK COSMOS-UK network," *Earth Syst. Sci. Data*, vol. 13, pp. 1737–1757, 2021, doi:10.5194/essd-13-1737-2021.
- [4] A. Hawdon, D. McJannet, and J. Wallace, "Calibration and correction procedures for cosmic-ray neutron soil moisture probes across Australia," *Water Resour. Res.*, vol. 50, pp. 5029–5043, 2014, doi:10.1002/2013WR015138.
- [5] L. Poggio *et al.*, "SoilGrids 2.0: producing soil information for the globe with quantified spatial uncertainty," *SOIL*, vol. 7, pp. 217–240, 2021, doi:10.5194/soil-7-217-2021.
- [6] S. Seabold and J. Perktold, "Statsmodels: econometric and statistical modeling with Python," in *Proc. 9th Python in Science Conf.*, 2010.