

1 Supporting Standard Results in Real Analysis

{fact:realanalysis}

Proposition 1. *Let $f : \mathbb{R}_{++} \rightarrow \mathbb{R}_+$ be a continuous function. Consider sequences x^n in \mathbb{R}_{++} and $f^n(x^n)$ in \mathbb{R}_+ . If $f^n(x^n) \rightarrow f(x)$ as $n \rightarrow \infty$, then $x^n \rightarrow x$ as $n \rightarrow \infty$.*

Proof. Given that f is continuous at x (with $x \in \mathbb{R}_{++}$), for every $\epsilon > 0$, there exists a $\delta > 0$ such that for all y in \mathbb{R}_{++} with $|y - x| < \delta$, we have $|f(y) - f(x)| < \epsilon$.

Given $f^n(x^n) \rightarrow f(x)$, for the above ϵ , there exists an N such that for all $n > N$, $|f^n(x^n) - f(x)| < \epsilon$.

Assume for the sake of contradiction that x^n doesn't converge to x . This implies that there exists a $\delta > 0$ such that for infinitely many terms of the sequence x^n , $|x^n - x| \geq \delta$.

By the continuity of f at x , if $|x^n - x| \geq \delta$ for infinitely many n , then $|f^n(x^n) - f(x)| \geq \epsilon$ for those n , contradicting our assumption that $f^n(x^n) \rightarrow f(x)$.

Therefore, our assumption for contradiction is false, and it follows that $x^n \rightarrow x$ as $n \rightarrow \infty$. \square

Fact 1. *Let $g : X \rightarrow \mathbb{R}_+$ be a continuous function, where $X \subseteq \mathbb{R}^n$ is an open convex set. Define the weighted supremum norm $\|\cdot\|_g$ of a real-valued function $f : X \rightarrow \mathbb{R}$ by*

$$\|f\|_g := \sup_{x \in X} \frac{|f(x)|}{g(x)}. \quad (1)$$

If $\lim_{n \rightarrow \infty} \|f_n - f^\|_g = 0$, f_n converges to f^* uniformly on compact sets.*

Proof. Let \tilde{X} be an arbitrary compact subset of X . Since \tilde{X} is compact, there exists a positive lower bound for g on this subset, denoted as

$$\bar{g} = \min_{x \in \tilde{X}} g(x) > 0. \quad (2)$$

Hence, on \tilde{X} , if $\lim_{n \rightarrow \infty} \|f_n - f^*\|_g = 0$, then $\lim_{n \rightarrow \infty} \|f_n - f^*\|_\infty = 0$ on \tilde{X} , where $\|\cdot\|_\infty$ denotes the supremum norm.

Now, let K be a compact subset of X . Given the continuity of g , there exists a positive maximum value for g on K , denoted as M_K . Then, we have

$$\sup_{x \in K} |f_n(x) - f(x)| \leq M_K \sup_{x \in K} \frac{|f_n(x) - f(x)|}{g(x)} \leq M_K \sup_{x \in X} \frac{|f_n(x) - f(x)|}{g(x)}. \quad (3)$$

Thus, $\lim_{n \rightarrow \infty} \|f_n - f\|_\infty = 0$ implies that f_n converges uniformly to f on the compact set K . It's also worth noting that the convexity and openness of X aren't strictly necessary for this argument. \square

{fact:compactnt}

Fact 2. *Let $\{f_n\}$ be a sequence of continuous functions defined on a subset of the real line and converging uniformly to a function f on compact sets. If $\{x_n\}$ is a convergent sequence of real numbers with limit x , then $f_n(x_n)$ converges to $f(x)$.*

Proof. Let \tilde{X} be an arbitrary compact subset of X . Since \tilde{X} is compact, there exists a positive lower bound for g on this subset, denoted as

$$\bar{g} = \min_{x \in \tilde{X}} g(x) > 0. \quad (4)$$

Hence, on \tilde{X} , if $\lim_{n \rightarrow \infty} \|f_n - f^*\|_g = 0$, then $\lim_{n \rightarrow \infty} \|f_n - f^*\|_\infty = 0$ on \tilde{X} , where $\|\cdot\|_\infty$ denotes the supremum norm.

Now, let K be a compact subset of X . Given the continuity of g , there exists a positive maximum value for g on K , denoted as M_K . Then, we have

$$\sup_{x \in K} |f_n(x) - f(x)| \leq M_K \sup_{x \in K} \frac{|f_n(x) - f(x)|}{g(x)} \leq M_K \sup_{x \in X} \frac{|f_n(x) - f(x)|}{g(x)}. \quad (5)$$

Thus, $\lim_{n \rightarrow \infty} \|f_n - f\| = 0$ implies that f_n converges uniformly to f on the compact set K . It's also worth noting that the convexity and openness of X aren't strictly necessary for this argument. \square

Fact 3. *Let $\{f_n\}$ be a sequence of continuous functions defined on a subset of the real line and converging uniformly to a function f on compact sets. If $\{x_n\}$ is a convergent sequence of real numbers with limit x , then $f_n(x_n)$ converges to $f(x)$.*

{fact:compactnts}

Proof. Since x_n converges to x , the sequence $\{x_n\}$ is bounded. Therefore, there exists a compact set K (specifically, a closed interval in the real line) that contains all the x_n as well as x .

Given the uniform convergence of f_n to f on K , for every $\epsilon > 0$, there exists an N such that for all $n \geq N$ and for all $y \in K$, we have

$$|f_n(y) - f(y)| < \epsilon.$$

In particular, for $y = x_n$, we have

$$|f_n(x_n) - f(x_n)| < \epsilon$$

for all $n \geq N$.

Now, each f_n being continuous and x_n converging to x implies that $f(x_n)$ converges to $f(x)$. Thus, for sufficiently large n , $f(x_n)$ can be made arbitrarily close to $f(x)$.

Combining the two inequalities and taking n large enough, we deduce that $|f_n(x_n) - f(x)|$ can be made smaller than any given ϵ . Hence, $f_n(x_n)$ converges to $f(x)$. \square