

Penguin trait variation in the Palmer Archipelago

Cindy Gao

2025-09-29

Introduction

Species' trait variation can give crucial insights into both the responses of species' to environmental gradients and the strength of filtering by those same gradients (Fitzgerald *et al.* 2025). For example, strong environmental filtering may cause a specific set of trait values to be best suited to a system, decreasing overall trait variation with knock-on effects for overall community diversity (Craven *et al.* 2018). Constrained trait values may also indicate low plasticity, developmental limits, or certain trait tradeoffs, which may also have negative implications for overall biodiversity (Arnold 1992). Consequently, identifying areas where species' trait variation is low or high may be of relevance to determining where species may be most threatened.

Methods

The study used the Palmer Penguins dataset, which has physical measurements for adult penguins observed on Biscoe, Dream, and Torgersen Island (Horst *et al.* 2020). The study sought to answer two questions related to spatial variation in penguin traits: 1) Do mean penguin trait values vary across Biscoe, Dream, and Torgersen Island, and 2) Do relationships between trait differ across Biscoe, Dream, and Torgersen Island?

The traits used in the study included flipper length (mm), bill length (mm), and body mass (g). Question 1 was answered using ANOVAs that tested for differences in mean trait values across islands. Pairwise comparisons were performed using a post-hoc Tukey test.

Question 2 was answered using a linear regression model that included an interaction term of the island. Pairwise comparisons were performed using a post-hoc Tukey test.

Figures




Figure 1: Boxplots of flipper length (mm) across islands. Biscoe Island had the widest distribution and was significantly different from Dream and Torgersen Islands.




Figure 2: Comparison of the relationship between flipper and bill length across Biscoe, Dream, and Torgersen Island. Bill and flipper measurements were grouped by island and each group was fitted with a separate linear regression line. The relationship between traits on Biscoe Island is significantly different from Dream and Torgersen Island, whereas the relationship between traits is not significantly different between Dream and Torgersen Island.

Tables

Table 1: Mean trait measurements across sites.

island	count	Mean Bill Length (mm)	Mean Flipper Length (mm)	Mean Body Mass (g)
Biscoe	163	45.24847	209.5583	4719.172
Dream	123	44.22195	193.1870	3718.902
Torgersen	47	39.03830	191.5319	3708.511

References

- Arnold, S.J. (1992). Constraints on phenotypic evolution. *The American Naturalist*, 140, S85–S107.
- Craven, D., Hall, J.S., Berlyn, G.P., Ashton, M.S. & Breugel, M. van. (2018). Environmental filtering limits functional diversity during succession in a seasonally wet tropical secondary forest. *Journal of Vegetation Science*, 29, 511–520.
- Fitzgerald, J.L., Ogilvie, J.E. & CaraDonna, P.J. (2025). Intraspecific body size variation across distributional moments reveals trait filtering processes. *Journal of Animal Ecology*, 94, 394–409.

Horst, A.M., Hill, A.P. & Gorman, K.B. (2020). *Allisonhorst/palmerpenguins: v0.1.0*. Zenodo.