Emerald ash borer

Simon Frost

Table of contents

1 Overview 1

2 The model 2

3 Spread year by year 2

4 The arrangement of ash 4

5 How many founders 6

6 Deviations from the source 7

1 Overview

The emerald ash borer, *Agrilus planipennis*, a beetle from Asia found near Detroit in 2002, has killed hundreds of millions of ash trees in North America. Its larvae feed in the phloem under the bark, girdling the tree. Mercader et al. (2011) asked how far an infestation spreads from a single point, and how that depends on where the ash is and on how the females choose where to lay. Their model is a coupled map lattice: cells of 50 m hold amounts of ash phloem and numbers of beetles, which spread, breed and eat phloem each year.

This implementation is a reproduction of the model's two behaviours in its published code, and an inferred version of a third; it reproduces the source's radii of infestation year by year.

```
ENV["GKSwstype"] = "100"
using EcologicalCellularAutomata
using CellularAutomataCore
using Plots
using Random
using Statistics
```

2 The model

Each year:

- adults emerge from larvae that took one year, and from those that took two;
- they disperse from their cell by a negative exponential kernel (`b = 0.037` per metre), normalised over the cells within 50 cells; with avoidance, cells without phloem are treated as 1.8 km further away;
- each female lays `r0 = 10.35` female offspring, as far as the phloem allows at `C = 0.0113` m^2 per larva, which the larvae eat;
- a share `-0.194 ln(x) + 0.986` of the larvae, `x` larvae per m^2 of phloem, take two years to develop.

The model is deterministic. The landscape is surrounded by a border of 2 cells where dispersing beetles are lost, as in the source's code.

```
prob = problem(:emerald_ash_borer)
size(prob.u0.phloem), sum(prob.u0.phloem[3:(end - 2), 3:(end - 2)])
```

```
((209, 209), 8.405e6)
```

The landscape is 205 × 205 cells (about 10 × 10 km) holding 8,405,000 m^2 of phloem, here spread evenly, 200 m^2 a cell. One hundred adults start at its centre.

3 Spread year by year


```
function spread(; years=15, kwargs...)
    integ = init(problem(:emerald_ash_borer; years, kwargs...);
save_everystep=false)
    out = [eab_spread(integ.u)]
    while integ.t < years
        step!(integ)
        push!(out, eab_spread(integ.u))
    end
    return out, integ.u
end
avoid, final = spread()
standard, _ = spread(; behaviour=:standard)
Int.(getfield.(avoid, :radius))
```

```
16-element Vector{Int64}:
    0
  150
  150
  250
  300
```

```
 400
 400
 500
 600
 700
 800
 900
1050
1150
1250
1350
```


These are the radii of infestation, from year 0 to year 15, with avoidance: the farthest cell from the origin holding at least one larva. They are those of the source's Fig. 9, year by year.

```
p1 = plot(0:15, [getfield.(standard, :radius) getfield.(avoid, :radius)];
marker=:circle,
          ms=3, lw=2, label=["standard" "avoidance"], xlabel="year", ylabel="radius
(m)",
          legend=:topleft)
plot!(p1, 0:15, [getfield.(standard, :median) getfield.(avoid, :median)]; ls=:dash,
      color=[1 2], label=["median, standard" "median, avoidance"])
p2 = plot(0:15, [getfield.(standard, :beetles) getfield.(avoid, :beetles)] ./ 1e6;
          lw=2, marker=:circle, ms=3, label=["standard" "avoidance"], xlabel="year",
          ylabel="beetles (millions)", legend=:topleft)
plot(p1, p2; layout=(1, 2), size=(850, 320), left_margin=4Plots.mm,
     bottom_margin=4Plots.mm)
```



Without avoidance, larvae fill the cells near the origin, eat them out, and the beetles emerging there keep laying in empty cells nearby. With avoidance they fly over the eaten-out core to fresh ash, so the infestation spreads further and holds more beetles: after 15 years 1350 m and 5.5 million here, against 1200 m and 3.9 million without (the source: 1350 m and about 5.3 million, 1200 m and about 3.8 million).


```
heatmap(final.phloem[3:(end - 2), 3:(end - 2)][63:143, 63:143]; aspect_ratio=1,
c=:YlGn,
        title="phloem left after 15 years, central 4 × 4 km", titlefontsize=10,
        xticks=false, yticks=false, size=(420, 380))
```



4 The arrangement of ash

The source compares three landscapes with the same total phloem: even, clumped (blocks of 5 × 5 cells separated by gaps of 5 empty cells) and random (cells holding 0 to 400 m² at random).

```
plot([heatmap(eab_environment(k; rng=Random.Xoshiro(1))[73:133, 73:133];
aspect_ratio=1,
              c=:YlGn, title=string(k), titlefontsize=10, xticks=false,
yticks=false,
              colorbar=false)
      for k in (:homogeneous, :clumped, :random)]...; layout=(1, 3), size=(800,
270))
```



The central 3 × 3 km of each. Fifteen years in each landscape, with each behaviour, and the source's values read from its Figs 4 and 5 (radius in m, beetles in millions):

```
source = Dict((:homogeneous, :standard) ⇒ (1200, 3.8),
              (:homogeneous, :avoidance) ⇒ (1350, 5.3),
              (:homogeneous, :resource_bias) ⇒ (1350, 5.3),
              (:clumped, :standard) ⇒ (750, 1.0), (:clumped, :avoidance) ⇒ (1650,
9.0),
              (:clumped, :resource_bias) ⇒ (2600, 10.5),
              (:random, :standard) ⇒ (1200, 3.8), (:random, :avoidance) ⇒ (1500,
6.2),
              (:random, :resource_bias) ⇒ (2050, 11.0))
cases = [(e, b) for e in (:homogeneous, :clumped, :random)
         for b in (:standard, :avoidance, :resource_bias)]
finals = fetch.([Threads.@spawn eab_spread(solve(problem(:emerald_ash_borer;
environment=e,
                                                 behaviour=b);
saveat=15).u[end])
                 for (e, b) in cases])
[(environment=e, behaviour=b, radius=r.radius, beetles=round(r.beetles / 1e6;
digits=2),
  source=source[(e, b)]) for ((e, b), r) in zip(cases, finals)]
```

```
9-element Vector{@NamedTuple{environment::Symbol, behaviour::Symbol,
radius::Float64, beetles::Float64, source::Tuple{Int64, Float64}}}:
 (environment = :homogeneous, behaviour = :standard, radius = 1200.0, beetles =
 3.88, source = (1200, 3.8))
 (environment = :homogeneous, behaviour = :avoidance, radius = 1350.0, beetles =
 5.46, source = (1350, 5.3))
 (environment = :homogeneous, behaviour = :resource_bias, radius = 1350.0, beetles =
 5.46, source = (1350, 5.3))
 (environment = :clumped, behaviour = :standard, radius = 750.0, beetles = 1.02,
 source = (750, 1.0))
 (environment = :clumped, behaviour = :avoidance, radius = 1650.0, beetles = 8.39,
 source = (1650, 9.0))
 (environment = :clumped, behaviour = :resource_bias, radius = 2050.0, beetles =
 9.65, source = (2600, 10.5))
 (environment = :random, behaviour = :standard, radius = 1200.0, beetles = 3.45,
 source = (1200, 3.8))
```

```
 (environment = :random, behaviour = :avoidance, radius = 1450.0, beetles = 6.27,
 source = (1500, 6.2))
 (environment = :random, behaviour = :resource_bias, radius = 1850.0, beetles =
 9.04, source = (2050, 11.0))
```

Standard and avoidance reproduce the source in every landscape. Clumping hinders a beetle that does not avoid empty ground: most of its offspring land in the gaps, so in the clumped landscape it spreads only 750 m. A beetle that avoids empty ground spreads furthest there, as each clump is eaten out and the beetles fly to the next. The source's third behaviour, a bias towards cells rich in phloem, is inferred here from its description; it reproduces the homogeneous result, and spreads further than avoidance in the other two landscapes, as in the source, but less far.

5 How many founders

The source compares 100 and 1,000 founders, and reports the difference in the final number of beetles as 67.5, 21.8 and 10.3% in the clumped, homogeneous and random landscapes:

```
envs = (:clumped, :homogeneous, :random)
founders = fetch.([Threads.@spawn eab_spread(solve(problem(:emerald_ash_borer;
environment=e,
                                                    founders=f);
saveat=15).u[end])
                   for e in envs for f in (100, 1000)])
[(environment=e, beetles_100=round(founders[2k - 1].beetles / 1e6; digits=2),
  beetles_1000=round(founders[2k].beetles / 1e6; digits=2),
  difference=round(100 * (founders[2k].beetles - founders[2k - 1].beetles) /
                   founders[2k].beetles; digits=1),
  radius_1000=founders[2k].radius)
 for (k, e) in enumerate(envs)]
```

```
3-element Vector{@NamedTuple{environment::Symbol, beetles_100::Float64,
beetles_1000::Float64, difference::Float64, radius_1000::Float64}}:
 (environment = :clumped, beetles_100 = 8.39, beetles_1000 = 4.24, difference =
 -98.0, radius_1000 = 1650.0)
 (environment = :homogeneous, beetles_100 = 5.46, beetles_1000 = 6.84, difference =
 20.2, radius_1000 = 1600.0)
 (environment = :random, beetles_100 = 6.27, beetles_1000 = 6.84, difference = 8.3,
 radius_1000 = 1650.0)
```

In the homogeneous and random landscapes, ten times as many founders give an infestation about 20% and 8% larger after 15 years (21.8% and 10.3% in the source), far from ten times larger, as the phloem near the origin limits it. In the clumped landscape the number of beetles swings from year to year, as the clumps are eaten out and the beetles fly to the next, and at year 15 the run with 1,000 founders happens to be in a trough; the source, whose clumped runs also oscillate strongly, finds 67.5% more.

6 Deviations from the source

```
foreach(d -> println("- ", d), model_info(:emerald_ash_borer).deviations)
```

```
- The source's Eq. 1 is misprinted; dispersal is normalised source by source over
the 101 x 101 window clipped at the edge of the lattice, as the source's code does,
with the distance of a cell from itself 0.25 cells.
- Avoidance adds 1.8 km (36 cells of 50 m), as the text says, to the distance of
cells without phloem; the code's example uses 18 cells. The results barely change.
- Of the source's five foraging behaviours, Standard and Avoidance (the two in its
code) are implemented as described. 'Resource Bias All' is inferred from the text:
cells whose initial phloem is at or above the landscape's upper quartile are 33%
nearer and those at or below the lower quartile 33% further, with no bias when the
quartiles are equal. It reproduces the homogeneous result (1350 m) but spreads less
than the source on the clumped (2050 against 2600 m) and random (1850 against 2050
m) landscapes. 'Stress bias' and 'Resource Bias Within 200 m', not in the code and
ambiguously described, are not implemented.
- The clumped landscape is 5 x 5 blocks of cells separated by 5 empty cells, one
centred, rescaled to the source's total of 8,405,000 m² (762.4 m² a cell, against
the 784.4 m² the source states, which does not divide its total among whole blocks).
The random landscape puts the 17 levels from 0 to 400 m² on equal numbers of cells
as far as they divide, sets the centre to 200 m² if it would be empty, and rescales
to the total; the source does not say how its levels were matched to the total.
- The radius of infestation is the largest distance from the origin of a cell with
at least one larva, floored to whole cells and times 50 m, as the source's code
computes it; with this rule the source's year-by-year radii are reproduced exactly.
- The cluster-gradient, habitat-quality and Saginaw County experiments are not
included; the Saginaw phloem map is not published.
```

Mercader, Rodrigo J., Nathan W. Siegert, Andrew M. Liebhold, and Deborah G. McCullough. 2011. “Influence of Foraging Behavior and Host Spatial Distribution on the Localized Spread of the Emerald Ash Borer, *Agrilus Planipennis*.” *Population Ecology* 53 (2): 271–85. https://doi.org/10.1007/s10144-010-0233-6.