Coastal dunes

Simon Frost

Table of contents

1 Overview 1

2 The model 2
2.1 Water levels . . . 2
2.2 Profiles from pieces . . . 3

3 The same potential transport 4

4 Fifty years of dune growth 5

5 Deviations from the source 9

1 Overview

Coastal foredunes grow where wind blows sand off the beach and dune plants trap it. Teixeira et al. (2023) study this with DuBeVeg, a cellular automaton in the family of Werner's sand-slab model: the surface is a stack of slabs on each cell, slabs are picked up and hop downwind until they land, avalanches keep slopes below the angle of repose, the sea resets the beach below the water level, and two kinds of plant grow or die back with the sand they gain. In the original model a slab hops one cell; the authors make it hop a saltation length of 5 cells, with slabs five times thinner so that the potential transport is the same, and let vegetation trap slabs part-way through a hop. On a wide beach this changes how the dune grows.

This implementation is an adaptation: the authors' code is not published, the sea is forced by synthetic water levels, the domain is narrower, and several constants come from the base model's documentation.


```
ENV["GKSwstype"] = "100"
using EcologicalCellularAutomata
using CellularAutomataCore
using Plots
using Random
using Statistics
```

2 The model

Each aeolian iteration (50 a year), every bare, dry cell outside the shadow of higher ground gives up a slab with probability `(1 - ρ) 0.5`, where ρ is its plant cover. A slab moves landward and, at the end of each hop of `L` cells, lands with probability `ρ (1 - 0.1) + 0.1` (always in the shadow); with `L = 5`, it can also land part-way through a hop on cells whose plant cover is at least 0.3. Every second iteration the sea, with run-up from waves, resets the cells it reaches towards the reference beach; once a year the plants grow or die back with the sand each cell gained.

The beach rises from the shoreline to the foot of a foredune 15 m high and 100 m wide; the source compares a narrow beach of 100 m with a wide one of 500 m.


```
plot(; xlabel="cross-shore distance (m)", ylabel="elevation (m)", size=(700, 300),
     legend=:topleft)
for (beach, c) in ((100, 1), (500, 2))
    prof = dune_profile(; beach, h=0.02)
    plot!(prof.elevation .* 0.02; lw=2, color=c, label="$beach m beach")
end
hline!([0]; color=:gray, ls=:dash, label="mean sea level")
```



2.1 Water levels

The source forces the sea with the highest water level in each spring-neap cycle at Scheveningen, 1970 to 2020, which it plots (its Fig. 5d) but does not tabulate. `dune_water_levels` stands in with a synthetic series, 25 levels a year: 1.43 m plus a Gumbel-distributed surge of scale 0.24 m, fitted to the plotted series read from the figure (a median of about 1.5 m, a tenth of the cycles above 2 m, a highest level of about 3.2 m in 50 years; one draw is below). Wave run-up adds a few tenths of a metre more. The level the sea reaches most years keeps the beach below it bare, and so sets where new dunes can start.

```
levels = dune_water_levels(Random.Xoshiro(1))
plot((1:length(levels)) ./ 25, levels; lw=0.6, color=:black, legend=false,
     xlabel="year", ylabel="water level (m)", size=(700, 240),
left_margin=4Plots.mm,
     bottom_margin=4Plots.mm)
```




```
(median=round(median(levels); digits=2), q90=round(quantile(levels, 0.9); digits=2),
 highest=round(maximum(levels); digits=2))
```

```
(median = 1.52, q90 = 1.94, highest = 3.48)
```

2.2 Profiles from pieces

A profile is built by joining pieces from the sea to the land: a Dean shoreface, a straight beach, the dune ridge and the plateau behind it. A piece runs from one elevation to another, and ∗ joins two pieces only where the first ends at the elevation where the second starts. Joining is associative and an empty piece changes nothing, so profiles compose like paths between elevations (morphisms in a category whose objects are elevations), and a scenario is a composite:

```
narrow = shoreface_segment(60) ∗ slope_segment(100, 0, 2.5) ∗
         ridge_segment(100, 2.5, 15, 5) ∗ flat_segment(20, 5; kind=:hinterland)
(narrow, profile_heights(narrow, 280; h=0.02) == dune_profile(;
beach=100).elevation)
```


```
(ProfileSegment(shoreface ∗ beach ∗ dune ∗ hinterland: -1.8391426377445272 → 5.0 m
over 280 m), true)
```

The reference surface the sea works towards is the same profile with the dune and the land behind it levelled at the dune's foot. `reference_profile` does this piece by piece, so it respects joins: it maps composites to composites.

Teixeira et al.'s third scenario is a prograding beach, which builds out 4 m a year, from 100 m to 300 m wide over 50 years. In their Fig. 2b the profile below a berm at 2 m moves seaward and the berm lengthens. With pieces, that is one more piece: the beach is cut where it reaches

2 m and a level berm of `4t` m is joined in. Level pieces at one elevation add, a berm of `a` m joined to one of `b` m being a berm of `a + b` m, so building out is time acting on the profile by joining berms. `problem(:dune_formation; scenario=:prograding)` puts the narrow beach on a domain 200 m longer out to sea, anchored at the land, and replaces the reference surface with the year's at the end of every year; the sea, resetting cells towards it, builds the beach out.

```
plot(; xlabel="cross-shore distance (m)", ylabel="elevation (m)", size=(700, 300),
     legend=:topleft, ylims=(-3, 16))
for y in 0:10:50
    ref = reference_profile(dune_landform(; beach=100, offshore=260, berm=4y))
    plot!(profile_heights(ref, 480); lw=1.5, color=cgrad(:copper)[y / 50],
    label="year $y")
end
plot!(profile_heights(dune_landform(; beach=100, offshore=260), 480); color=:black,
lw=2,
      label="the dune at the start")
```



3 The same potential transport

The source's Eq. 3 gives the potential transport `Q = h L (pe0 / pd0) n`: with slabs of 0.1 m hopping 1 cell or of 0.02 m hopping 5, 25 m³ per metre of beach a year. On a bare, dry beach the slabs crossing a line do carry that:

```
function flat_flux(h, L; rows=40, iterations=20)
    p = merge(defaults(build(:dune_formation)),
              (; h, L, rho_star=Inf, hydro_every=10^6, iterations=10^6))
    total = 0
    for t in 1:iterations
        u = (topo=fill(Int32(round(2.9 / h)), rows, 400), pioneer=zeros(rows, 400),
             stabilizer=zeros(rows, 400), balance=zeros(Int32, rows, 400),
             zref=zeros(Int32, rows, 400), sea=zeros(rows, 400))
```

```
        u.topo[:, 200:end] .= round(Int32, 3.1 / h)
        total += dune_step!(u, p, t, Random.Xoshiro(t))
    end
    return round(total * h * 50 / (rows * iterations); digits=1)
end
(hop_1=flat_flux(0.1, 1), hop_5=flat_flux(0.02, 5))
```

```
(hop_1 = 24.6, hop_5 = 24.9)
```

4 Fifty years of dune growth


One run of 50 years for each beach and representation, on 20 rows of 1 m (the source runs 400 rows five times). We record the sand crossing the dune toe, the first cell above 3 m, each year, and the dune's volume and width every ten years.

```
function dune_run(scenario, representation; years=50, rows=20, seed=1)
    prob = problem(:dune_formation; scenario, representation, alongshore=rows,
years, seed)
    p, u, h = prob.p, deepcopy(prob.u0), prob.p.h
    flux = zeros(years)
    measures = [dune_measures(u; h)]
    profiles = [vec(mean(u.topo; dims=1)) .* h]
    for t in 1:(50years)
        crossed = dune_step!(u, p, t, Random.Xoshiro(hash((prob.seed, t, :dune))))
        flux[cld(t, 50)] += crossed * h / rows
        if t % 500 == 0
            push!(measures, dune_measures(u; h))
            push!(profiles, vec(mean(u.topo; dims=1)) .* h)
        end
    end
    return (; flux, measures, profiles, final=u.topo .* h)
end
cases = [(s, r) for s in (:narrow, :wide, :prograding) for r in (:default,
:saltation)]
runs = Dict(zip(cases, fetch.([Threads.@spawn dune_run(c...) for c in cases])))
nothing
```


The same runs drive the integrator: `dune_step!` with the generator seeded from the problem's seed and the iteration is what `solve` runs at each step.

```
plot([begin
          r = runs[(s, rep)]
          plot(r.profiles[1]; color=:black, lw=1.5, label="start",
               title="$s beach, hop $(rep === :saltation ? 5 : 1)",
titlefontsize=10,
               xlabel="cross-shore (m)", ylabel="elevation (m)", legend=:topleft,
               xlims=s === :wide ? (400, 680) : s === :prograding ? (100, 480) : (0,
280),
```

```
            ylims=(-1, 22))
        plot!(r.profiles[end]; color=2, lw=2, label="after 50 years")
    end
    for (s, rep) in cases]...; layout=(3, 2), size=(850, 800))
```




```
heatmap(hcat(permutedims(runs[(:wide, :default)].final[:, 400:end]),
             permutedims(runs[(:wide, :saltation)].final[:, 400:end])); c=:terrain,
        clims=(-1, 20), xticks=false, ylabel="cross-shore (m, from 400)", size=(600,
360),
        title="wide beach after 50 years: hop 1 (left), hop 5 (right)",
titlefontsize=10)
```





```
source_volume = Dict((:narrow, :default) => 1500, (:narrow, :saltation) => 1611,
                     (:wide, :default) => 1563, (:wide, :saltation) => 2234,
                     (:prograding, :default) => 1540, (:prograding, :saltation) =>
2130)
source_width = Dict((:narrow, :default) => 145, (:narrow, :saltation) => 150,
                    (:wide, :default) => 220, (:wide, :saltation) => 230,
                    (:prograding, :default) => 304, (:prograding, :saltation) =>
218)
[(beach=s, hop=rep === :saltation ? 5 : 1,
  toe_flux=round(mean(runs[(s, rep)].flux[2:end]); digits=1),
  volume_gain=round(Int,
                    runs[(s, rep)].measures[end].volume - runs[(s,
rep)].measures[1].volume),
  source_gain=source_volume[(s, rep)] - 1150,
  width=round(Int, runs[(s, rep)].measures[end].width),
source_width=source_width[(s, rep)])
 for (s, rep) in cases]
```

```
6-element Vector{@NamedTuple{beach::Symbol, hop::Int64, toe_flux::Float64,
volume_gain::Int64, source_gain::Int64, width::Int64, source_width::Int64}}:
 (beach = :narrow, hop = 1, toe_flux = 18.5, volume_gain = 384, source_gain = 350,
 width = 166, source_width = 145)
 (beach = :narrow, hop = 5, toe_flux = 18.9, volume_gain = 541, source_gain = 461,
 width = 165, source_width = 150)
 (beach = :wide, hop = 1, toe_flux = 13.4, volume_gain = 366, source_gain = 413,
 width = 280, source_width = 220)
 (beach = :wide, hop = 5, toe_flux = 23.0, volume_gain = 1040, source_gain = 1084,
 width = 252, source_width = 230)
```

```
(beach = :prograding, hop = 1, toe_flux = 13.0, volume_gain = 377, source_gain =
390, width = 241, source_width = 304)
(beach = :prograding, hop = 5, toe_flux = 22.7, volume_gain = 1028, source_gain =
980, width = 213, source_width = 218)
```

The sand crossing the dune toe, in m³ per metre a year, the gain in dune volume over 50 years and the dune's width after them, against the source's medians (from an initial volume of about 1150 m³/m; the medians of the narrow and wide beaches are in its text, those of the prograding beach and the wide beach's widths with hops of 1 cell read from its Fig. 4). As in the source, hops of 5 cells carry sand across a wide beach to the dune close to the potential transport, while single-cell hops build bedforms on the beach that hold sand back. So the dune gains much more with hops of 5 cells on the wide beach, here as in the source; on the narrow beach the two differ less.

```
plot(1:50, [runs[c].flux for c in cases]; lw=1.5,
     label=permutedims(["$(c[1]), hop $(c[2] === :saltation ? 5 : 1)" for c in
cases]),
     xlabel="year", ylabel="flux at the dune toe (m³/m/yr)", ylims=(0, 30),
     size=(650, 340), legend=:bottomright)
hline!([25]; color=:gray, ls=:dash, label="potential transport")
```



On the prograding beach the model follows the source closely. With hops of one cell the beach builds out faster than the dune can follow, and a row of small separate foredunes forms on it, near the shoreline, and the dune is wide (241 m against the source's 304 m, its widest). With hops of 5 cells the sand reaches the old dune, which grows a wide new ridge in front of it (213 m against 218 m), and the flux at the dune toe rises over the 50 years as the beach widens, as in the source's Fig. 5c. The gains in volume are 377 and 1028 m³/m, against about 390 and 980.

The dunes of the narrow and wide beaches grow somewhat wider than the source's: 166 and 165 m against 145 and 150 on the narrow beach, 280 and 252 m against 220 and 230 on the wide

one. New dunes start on the beach where the sea rarely reaches, and their seaward faces then advance until the storms that reach them balance the sand they trap. On the narrow beach that face stops at about 1.4 m above mean sea level here, where the source's stops nearer the dune's old foot at 2.5 m. The waves follow DuBeVeg's code: their initial strength is the total water level (in slabs) times an energy factor of 1, which keeps them strong far up the beach and, near the shoreline, overrules the protection of plants; the source's Eq. 4 leaves the factor out, and without it the narrow beach's face stopped at 1.1 m. An earlier synthetic series of water levels, 0.4 m lower than the source's, let them start much further down the beach (the wide beach's dune was then 340 to 375 m wide); fitting the series to the source's figure removed most of the difference. Raising the least water depth in the waves' dissipation from 0.01 m to 0.4 m (the base model's calibrated value) changes nothing here, as the waves reach the dune with their least strength either way. Vignette 28 shows that a wind steered by the topography also narrows the dunes, though the source's wind is not steered.

5 Deviations from the source

```
foreach(d -> println("- ", d), model_info(:dune_formation).deviations)
```

```
- The source forces the sea with observed water levels at Scheveningen (1970-2020),
the highest in each spring-neap cycle; here a synthetic series stands in: 1.43 m
plus a Gumbel-distributed surge of scale 0.24 m, 25 a year, fitted to the series the
source plots (its Fig. 5d, read from the figure: a median of about 1.5 m, a tenth of
the cycles above 2 m, a highest level of about 3.2 m). An earlier series 0.4 m lower
let new dunes start far down the beach and made the dunes much too wide.
- The domain is 20 to 100 rows of 1 m alongshore (400 in the source) and only the
cross-shore profile from 60 m offshore to 20 m behind the dune (1250 cells in the
source). The initial profile is a Dean-like shoreface, a linear beach and a smooth
dune ridge to 15 m read from the source's figure; the beach rises to 2.5 m at the
dune foot, as the figure shows, where the text says 3 m (with 3 m, small bumps on
the upper beach count as the dune).
- Slab transport is synchronous, as in DuBeVeg's code: all pick-ups are drawn from
the state at the start of an iteration, with the shadow and the probabilities fixed
for its duration.
- Part-way through a hop (Eq. 5), a slab lands only where vegetation is at least
rho_star, and the shadow does not force it to; after a failed landing at a hop's end
the next hop begins. Slabs reaching the land side of the domain are lost.
- The shadow test uses the continuous threshold i tan(15°) rather than DuBeVeg's
floored one.
```

- Constants not in the source come from DuBeVeg: the growth functions of the pioneer (burial -1.4, 0.2, 0.6, 2.0, 2.2 m/yr, peak 0.2) and stabilizer (-1.4, -0.65, 0, 0.2, 2.8, peak 0.05), lateral expansion 0.2 and establishment 0.05 a year (stabilizers establishing only where the sand balance is zero), the wave run-up relation, the dissipation by the inverse water depth summed from the sea, the least wave probability 0.1, and the waves' initial strength, the total water level in slabs times Fenergy = 1 (wave_energy), which multiplies WE - Wdiss D in DuBeVeg's code (the source's Eq. 4 leaves it out); near the shoreline it overrules the protection of vegetation, as in DuBeVeg. Plant cover is multiplied each year by the chance that the sea did not reset the cell. The initial split of species is not stated: pioneers seaward of the crest, stabilizers behind.
- The dune is wider than in the source after 50 years on the narrow and wide beaches (narrow 166 and 165 m with hops of 1 and 5 cells, against 145 and 150; wide 280 and 252 m against 220 and 230): the seaward faces of new dunes advance further down the beach before the storms that reach them balance the sand they trap (on the narrow beach to about 1.4 m above mean sea level, against about 2.5 m in the source). Raising the least water depth of the wave dissipation from 0.01 to 0.4 m (the base model's calibrated value) does not change this.
- The prograding beach builds out through the sea alone: the reference surface is replaced at the end of each year by the profile with a berm 4 m longer, the profile below the berm moved seaward within a domain extending 200 m further out (the source's profile reaches a depth of closure of 8 m, beyond the domain here), and the sea's resets move the beach towards it; the shoreline then follows the reference about a year behind.
- The sensitivity to rho_star is not included; one run per case on 20 rows replaces the source's five runs on 400.
- Beyond the source: three more landforms built from the same profile pieces (an embryo dune and slack seaward of the foredune on the wide beach, a seawall and promenade at the back of the narrow beach, and a breach through the foredune in a fifth of the rows), and a non-erodible layer (hard) for the seawall; the source runs none of them, so they test the model's behaviour, not its fidelity. A breach fills from its flanks rather than growing into a blowout.
- Beyond the source: an optional wind steered by the topography (speedup > 0; off by default), from the linear theory of turbulent flow over low hills (Kroy et al. 2002 along the wind, Schwammle and Herrmann 2005 across it), computed on the height above the reference surface with the shadow envelope in the lee. The erosion probability scales with the shear along the wind and flying slabs turn alongshore with the shear across it. With A = 4 it narrows the dunes of the narrow and wide beaches towards the source's widths, but a breach still fills.

Teixeira, Manuel, Erik M. Horstman, and Kathelijne M. Wijnberg. 2023. “Conceptualizing Aeolian Sediment Transport in a Cellular Automata Model to Simulate the Bio-Geomorphological Evolution of Beach–Dune Systems.” *Journal of Marine Science and Engineering* 11 (7): 1278. https://doi.org/10.3390/jmse11071278.