Dune landforms

Simon Frost

Table of contents

1 Overview 1

2 Landforms from pieces 2

3 Fifty years on each landform 3

4 An embryo dune starves the foredune 4

5 Sand piles against a seawall 5

6 A breach heals from its flanks 7

7 A landform of one's own 8

8 Summary 9

1 Overview

The coastal dune model of Teixeira et al. (2023) (vignette 20) starts from one shape: a beach rising to a single foredune with a plateau behind it. Real coasts have other shapes. An embryo dune may grow on the beach in front of the foredune, a seawall may stand where a dune would be, and a dune may be breached. In this package a profile is built by joining pieces, so each of these is a short composite of the same pieces.

The source runs none of these landforms. Running them tests the model's behaviour, not its fidelity: does it do what geomorphology leads one to expect on shapes it was not tuned for? One addition is needed: a seawall must not erode. So the model has a non-erodible layer, `hard`, below which neither wind, avalanches nor the sea remove sand, and on which plants grow only once sand covers it. It is empty by default, and the source's scenarios run exactly as before.

```
ENV["GKSwstype"] = "100"
using EcologicalCellularAutomata
using CellularAutomataCore
using Plots
using Random
using Statistics
```

2 Landforms from pieces

`dune_landform(; form)` builds four shapes, each `offshore + beach + 120` metres long:

- the source's foredune: shoreface, beach, a ridge rising to 15 m, and a plateau;
- a double ridge: an embryo dune 40 m wide and 7 m high on the wide beach, with a level dune slack 160 m long between it and the foredune;
- a seawall: the beach ends at a wall rising to 7.5 m, with a promenade behind it in place of the dune and plateau;
- a breach: the foredune lowered to a trough rising only to the plateau's 5 m.

Each is a composite of pieces. The kind of each piece says how the model treats it. The sea's reference surface levels pieces of kind `:dune` and everything landward of them, and pieces of kind `:seawall` and `:promenade` are non-erodible:

```
forms = (foredune=dune_landform(; beach=100), double_ridge=dune_landform(;
form=:double_ridge),
         seawall=dune_landform(; form=:seawall, beach=100),
         breach=dune_landform(; form=:breach, beach=100))
map(f -> join(unique(profile_kinds(f, length(f))), " * "), forms)
```

```
(foredune = "shoreface * beach * dune * hinterland", double_ridge = "shoreface *
beach * dune * slack * hinterland", seawall = "shoreface * beach * seawall *
promenade", breach = "shoreface * beach * dune * hinterland")
```

```
plot([begin
          x = name === :double_ridge ? (301:680) : (1:280)
          plot(x, f.heights[x]; lw=2, color=:sienna, label="profile",
          title=string(name),
               titlefontsize=10, ylims=(-2, 16), legend=name === :foredune ?
:topleft : false)
          plot!(x, reference_profile(f).heights[x]; ls=:dash, color=:steelblue,
                label="reference")
      end
      for (name, f) in pairs(forms)]...; layout=(2, 2), size=(800, 500),
     xlabel="cross-shore (m)", ylabel="elevation (m)")
```




Figure 1: The four landforms (land to the right): the source's foredune on the narrow beach, the double ridge on the wide beach (shown from 300 m), the seawall and promenade, and the breached foredune. Dashed lines are the reference surfaces the sea works towards.

`problem(:dune_formation; scenario)` sets each up: `:double_ridge` on the wide beach, `:seawall` on the narrow one, and `:blowout`, the narrow beach with the breach in the middle fifth of the rows and the source's foredune in the rest. A `landform` keyword accepts any composite, or one for each row.

3 Fifty years on each landform

Two runs of 50 years on each landform, with hops of 5 cells (the source's saltation representation), beside the source's narrow and wide beaches. The runs use 20 rows of 1 m, or 50 for the blowout so that the breach is 10 m wide. Each run records the surface every ten years.

```
function landform_run(scenario; rows=20, years=50, seed=1)
    prob = problem(:dune_formation; scenario, alongshore=rows, years, seed)
    p, u, h = prob.p, deepcopy(prob.u0), prob.p.h
    snaps = [u.topo .* h]
    intact = true
    for t in 1:(50years)
        dune_step!(u, p, t, Random.Xoshiro(hash((prob.seed, t, :dune))))
        t % 500 == 0 && push!(snaps, u.topo .* h)
        intact &= all(u.topo .>= u.hard)
```

```
    end
    return (; snaps, intact)
end
cases = [(s, seed) for s in (:narrow, :wide, :double_ridge, :seawall, :blowout) for
seed in 1:2]
runs = Dict(zip(cases,
                fetch.([Threads.@spawn landform_run(s; seed, rows=s === :blowout ?
                50 : 20)
                        for (s, seed) in cases])))
volume(z, cols) = sum(max.(z[:, cols] .- 3, 0)) / size(z, 1)    # m³/m above 3 m
gain(r, cols) = volume(r.snaps[end], cols) - volume(r.snaps[1], cols)
sand(r, cols) = sum(r.snaps[end][:, cols] .- r.snaps[1][:, cols]) / size(r.snaps[1],
1)
nothing
```

4 An embryo dune starves the foredune

The double ridge has the same domain as the wide beach, and its foredune occupies the same columns (561 to 680). The two can be compared column for column:

```
[(seed=seed, foredune_gain_wide=round(Int, gain(runs[(:wide, seed)], 561:680)),
  foredune_gain_double=round(Int, gain(runs[(:double_ridge, seed)], 561:680)),
  embryo_and_slack_gain=round(Int, gain(runs[(:double_ridge, seed)], 361:560)),
  embryo_crest=round(maximum(mean(runs[(:double_ridge, seed)].snaps[end][:,
361:400]; dims=1));
                     digits=1),
  slack_lowered=round(-sand(runs[(:double_ridge, seed)], 401:560) / 160; digits=2))
 for seed in 1:2]
```

```
2-element Vector{@NamedTuple{seed::Int64, foredune_gain_wide::Int64,
foredune_gain_double::Int64, embryo_and_slack_gain::Int64, embryo_crest::Float64,
slack_lowered::Float64}}:
 (seed = 1, foredune_gain_wide = 220, foredune_gain_double = 89,
 embryo_and_slack_gain = 179, embryo_crest = 12.6, slack_lowered = 0.46)
 (seed = 2, foredune_gain_wide = 235, foredune_gain_double = 89,
 embryo_and_slack_gain = 160, embryo_crest = 12.0, slack_lowered = 0.48)
```

The embryo dune intercepts the sand. It grows from 7 m to about 12 m, and the foredune behind it gains less than half what it gains on the open wide beach. The two ridges together gain a little more than the single foredune does. The open wide beach grows embryo dunes of its own (the dashed line in the figure below), as vignette 20 notes, but starting from bare sand they let more of it through to the foredune.

The bare slack between them loses sand. It is lowered by about half a metre: from 2.5 m to the groundwater surface, which the model puts at 0.8 times the reference surface, 2.0 m. Below that the sand is wet and the wind cannot lift it. Deflation down to the water table is how dune slacks form, and the model does it without being told.

```
x = 300:680
plot(x, vec(mean(runs[(:double_ridge, 1)].snaps[1]; dims=1))[x]; color=:black,
lw=1.5,
     label="double ridge, start", xlabel="cross-shore (m)", ylabel="elevation (m)",
     legend=:topleft, size=(750, 380), ylims=(-1, 22), left_margin=4Plots.mm,
     bottom_margin=4Plots.mm)
plot!(x, vec(mean(runs[(:double_ridge, 1)].snaps[end]; dims=1))[x]; color=2, lw=2,
      label="double ridge, 50 years")
plot!(x, vec(mean(runs[(:wide, 1)].snaps[end]; dims=1))[x]; color=1, lw=1.5,
ls=:dash,
      label="wide beach, 50 years")
hline!([2.0]; color=:gray, ls=:dot, label="groundwater surface")
```




Figure 2: Mean profiles after 50 years on the wide beach with and without an embryo dune (one run each). The embryo dune grows and the foredune behind it grows less; the slack between them deflates to the groundwater surface (2 m, dotted).

5 Sand piles against a seawall

On the narrow beach, a seawall stands where the dune foot would be:

```
[(seed=seed, wall_intact=runs[(:seawall, seed)].intact,
  upper_beach_sand=round(Int, sand(runs[(:seawall, seed)], 131:160)),
  promenade_sand=round(Int, sand(runs[(:seawall, seed)], 163:280)),
  promenade_depth_first_20_m=round(mean(runs[(:seawall, seed)].snaps[end][:,
163:182] .-
                                        runs[(:seawall, seed)].snaps[1][:,
163:182]);
```

```
                                    digits=1),
 dune_gain_without_wall=round(Int, gain(runs[(:narrow, seed)], 161:280)))
for seed in 1:2]
```

```
2-element Vector{@NamedTuple{seed::Int64, wall_intact::Bool,
upper_beach_sand::Int64, promenade_sand::Int64, promenade_depth_first_20_m::Float64,
dune_gain_without_wall::Int64}}:
 (seed = 1, wall_intact = 1, upper_beach_sand = 251, promenade_sand = 2,
 promenade_depth_first_20_m = 0.1, dune_gain_without_wall = 210)
 (seed = 2, wall_intact = 1, upper_beach_sand = 261, promenade_sand = 3,
 promenade_depth_first_20_m = 0.1, dune_gain_without_wall = 226)
```

The wall and promenade are never eroded. Sand blown off the beach piles against the wall, and plants colonise the pile. In 50 years a dune more than 11 m high grows on the upper beach in front of the wall, higher than the wall itself. Almost no sand gets past it: the promenade gains only a few m³ per metre, a decimetre near the wall, as the vegetated dune traps what the wind brings. A dune in front of a seawall shelters what lies behind it, here without any rule written for it. The balance is fine, though: with water levels 0.4 m lower (an earlier version of this model's synthetic series) the dune was lower and the promenade was buried by 2 to 3 m near the wall, the familiar nuisance of sand on seafront boulevards.

```
x = 100:280
z = runs[(:seawall, 1)].snaps
plot(x, dune_landform(; form=:seawall, beach=100).heights[x]; color=:gray, ls=:dash,
lw=2,
     label="wall and promenade", xlabel="cross-shore (m)", ylabel="elevation (m)",
     legend=:bottomright, size=(750, 380), left_margin=4Plots.mm,
bottom_margin=4Plots.mm)
for (k, y) in enumerate((0, 10, 30, 50))
    plot!(x, vec(mean(z[y ÷ 10 + 1]; dims=1))[x]; color=cgrad(:copper)[k / 4],
    lw=1.5,
          label="year $y")
end
plot!()
```




Figure 3: Mean profile at the seawall after 0, 10, 30 and 50 years (one run), against the non-erodible surface (dashed). A dune builds against the wall, higher than it, and little sand reaches the promenade.

6 A breach heals from its flanks

In the blowout scenario the foredune is breached in 10 of 50 rows, down to 5 m. The table gives the crest height in the breach every ten years, and the dune's gain in volume in the breached and intact rows:

```
gap, rest = 21:30, [1:20; 31:50]
crest(z, rows) = mean(maximum(z[rows, 161:260]; dims=2))
rows_of(r, rows) = (; snaps=[s[rows, :] for s in r.snaps])
[(seed=seed,
  breach_crest_by_decade=[round(crest(z, gap); digits=1) for z in runs[(:blowout,
seed)].snaps],
  intact_crest=round(crest(runs[(:blowout, seed)].snaps[end], rest); digits=1),
  gain_breach=round(Int, gain(rows_of(runs[(:blowout, seed)], gap), 161:280)),
  gain_intact=round(Int, gain(rows_of(runs[(:blowout, seed)], rest), 161:280)),
  gain_without_breach=round(Int, gain(runs[(:narrow, seed)], 161:280)))
 for seed in 1:2]
```

```
2-element Vector{@NamedTuple{seed::Int64, breach_crest_by_decade::Vector{Float64},
intact_crest::Float64, gain_breach::Int64, gain_intact::Int64,
gain_without_breach::Int64}}:
 (seed = 1, breach_crest_by_decade = [5.0, 13.4, 14.9, 15.1, 15.3, 15.6],
 intact_crest = 17.9, gain_breach = 542, gain_intact = 123, gain_without_breach =
 210)
```

```
(seed = 2, breach_crest_by_decade = [5.0, 13.1, 13.9, 14.7, 15.1, 15.2],
intact_crest = 17.4, gain_breach = 531, gain_intact = 116, gain_without_breach =
226)
```

The breach fills within ten years, to a crest of 13 to 14 m. After fifty its crest stands above the dune's original 15 m, though still a metre or two below the intact rows, which have grown too. Most of the sand comes from its flanks: the walls of the breach stand 10 m above its floor, far steeper than the angle of repose, and avalanche into it. The breached rows gain four to five times as much as the intact ones, and the intact rows gain less than the dune does on the same beach without a breach. Over all rows, the dune gains about what it gains without a breach.

```
plot([heatmap(120:280, 1:50, z[:, 120:280]; c=:terrain, clims=(-1, 20), title="year
$y",
              titlefontsize=10, xlabel="cross-shore (m)", ylabel=y == 0 ?
"alongshore (m)" : "")
      for (y, z) in zip((0, 10, 50), runs[(:blowout, 1)].snaps[[1, 2, 6]])]...;
     layout=(1, 3), size=(1000, 330), left_margin=4Plots.mm,
bottom_margin=5Plots.mm)
```




Figure 4: Elevation around the breached foredune at the start and after 10 and 50 years (one run; rows alongshore, cross-shore from 120 m).

Real breaches often grow into blowouts instead. They do so because the breach steers and speeds up the wind, which then scours it. By default the model has no such steering: slabs travel straight along their row whatever the topography. Vignette 28 adds a wind steered by the topography, from the linear theory of flow over low hills; the breach still fills, a little faster.

7 A landform of one's own

Any composite can be run. A common Dutch coastal defence, the dune-fronted dike, puts a small dune in front of a seawall. Here it is, with the seawall kept non-erodible behind a dune 40 m wide:

```
own = shoreface_segment(60) * slope_segment(100, 0, 2.5) * ridge_segment(40, 2.5, 8,
2.5) *
      slope_segment(2, 2.5, 7.5; kind=:seawall) * flat_segment(78, 7.5;
      kind=:promenade)
prob = problem(:dune_formation; landform=own, alongshore=4, years=1)
(cells=size(prob.u0.topo), hard_cells=count(>(typemin(Int32)), prob.u0.hard[1, :]),
 run_up_slope=prob.p.slope)
```

```
(cells = (4, 280), hard_cells = 80, run_up_slope = 0.025)
```

The run-up slope comes from the beach pieces, 2.5 m over 100 m. The source uses 3 m over the beach width, and the source's scenarios keep that.

8 Summary

landform	expected	in the model
embryo dune on the beach	intercepts sand, starving the foredune	foredune gains less than half; slack deflates to the water table
seawall	sand piles against it and blows onto the promenade	a dune higher than the wall in front of it, which shelters the promenade
breached foredune	may heal or grow into a blowout	mostly refilled within a decade by avalanching; no growth, even with the wind steered (vignette 28)

Teixeira, Manuel, Erik M. Horstman, and Kathelijne M. Wijnberg. 2023. "Conceptualizing Aeolian Sediment Transport in a Cellular Automata Model to Simulate the Bio-Geomorphological Evolution of Beach–Dune Systems." *Journal of Marine Science and Engineering* 11 (7): 1278. https://doi.org/10.3390/jmse11071278.