
Mathematics of Machine Learning


Integral Calculus



Riemann integration, integration by parts, substitution, change of variables


Outline

1. **Defining Integrals**
2. Doing Integrals
3. Changing Variables



Riemann Integral

- Integrals represent area beneath a curve



Linearity of Integration

- Integration is a linear operator

$$\int_a^b (r\,f(x) + s\,g(x))\,\mathrm{d}x = \lim_{\delta x_i \to 0} \sum_{i=0}^{n-1} (x_{i+1} - x_i)(r\,f(x_i) + s\,g(x_i))$$

$$= \lim_{\delta x_i \to 0} \left(\sum_{i=0}^{n-1} (x_{i+1} - x_i)\,r\,f(x_i) + \sum_{i=0}^{n-1} (x_{i+1} - x_i)\,s\,g(x_i) \right)$$

$$= \lim_{\delta x_i \to 0} \left(r \sum_{i=0}^{n-1} (x_{i+1} - x_i)\,f(x_i) + s \sum_{i=0}^{n-1} (x_{i+1} - x_i)\,g(x_i) \right)$$

$$= r \lim_{\delta x_i \to 0} \sum_{i=0}^{n-1} (x_{i+1} - x_i)\,f(x_i) + s \lim_{\delta x_i \to 0} \sum_{i=0}^{n-1} (x_{i+1} - x_i)\,g(x_i)$$


$$= r \int_a^b f(x)\,\mathrm{d}x + s \int_a^b f(x)\,\mathrm{d}x$$

Fundamental Theorem of Calculus

- Let

$$I(a,x) = \int_a^x f(z)\,\mathrm{d}z = \lim_{\delta z_i \to 0} \sum_{i=0}^{n-1} (z_{i+1} - z_i)\, f(z_i)$$

- Now for small δx



$$I(a,x+\delta x) = \int_a^{x+\delta x} f(z)\,\mathrm{d}z = \lim_{\delta z_i \to 0} \sum_{i=0}^{n-1} (z_{i+1} - z_i)\, f(z_i) + \delta x\, f(x)$$

- Thus

$$\frac{\mathrm{d}I(a,x)}{\mathrm{d}x} = \lim_{\delta x \to 0} \frac{I(x+\delta x) - I(x)}{\delta x} = \lim_{\delta x \to 0} \frac{\delta x\, f(x)}{\delta x} = f(x)$$

The Other Way Around

- Consider

$$\int_a^b \frac{\mathrm{d}f(x)}{\mathrm{d}x}\mathrm{d}x = \int_a^b \lim_{\delta x \to 0} \frac{f(x+\delta x) - f(x)}{\delta x}\mathrm{d}x$$

$$= \lim_{x_{i+1}-x_i \to 0} \sum_{i=0}^{n-1} (x_{i+1} - x_i) \frac{f(x_{i+1}) - f(x_i)}{x_{i+1} - x_i}$$

$$= \lim_{x_{i+1}-x_i \to 0} \sum_{i=0}^{n-1} (f(x_{i+1}) - f(x_i))$$

$$= (f(x_1) - f(x_0)) + (f(x_2) - f(x_1)) + (f(x_3) - f(x_2)) + \cdots$$
$$+ (f(x_{n-1}) - f(x_{n-2})) + (f(x_n) - f(x_{n-1}))$$

$$= f(x_n) - f(x_0) = f(b) - f(a)$$

- We can think of integration as an **anti-derivative**—it undoes differentiation

Indefinite Integrals

- So far we have considered **definite integrals** where we integrate between two points (a and b)
- However, when think about integration as an anti-derivative, it is useful to think of a function $F(x) = \int f(x)\mathrm{d}x$
- So that $F'(x) = f(x)$
- However the function $F(x)$, $F(x) + 1$, $F(x) + \pi$, etc. all have the same derivative so $F(x)$ is only defined up to an additive constant
- Note that the definite integral is given by


$$\int_a^b f(x)\mathrm{d}x = F(b) - F(a)$$

Multiple Integrals

- For functions involving many independent variables (e.g. $f(x,y)$, $f(x,y,z)$, $f(\boldsymbol{x})$) we can integrate over multiple dimensions

- For example

$$\iint_A f(x,y)\,\mathrm{d}x\,\mathrm{d}y$$




- It gets tedious writing multiple integral signs and I tend to write just one

$$\int \cdots \int f(x_1,x_2,\ldots,x_n)\,\mathrm{d}x_1\,\mathrm{d}x_2\cdots\mathrm{d}x_n = \int f(\boldsymbol{x})\,\mathrm{d}\boldsymbol{x}$$

Outline

1. Defining Integrals
2. **Doing Integrals**
3. Changing Variables



Performing Integration

- A key method for performing integrals is through knowledge of the anti-derivative
- If we know $F'(x) = f(x)$ then $F(x) + c = \int f(x)\mathrm{d}x$
- E.g. we know that $\mathrm{d}x^n/\mathrm{d}x = nx^{n-1}$ therefore

$$\int x^{n-1}\mathrm{d}x = \frac{1}{n}\int \frac{\mathrm{d}x^n}{\mathrm{d}x}\mathrm{d}x = \frac{x^n}{n} + c$$

and

$$\int_a^b x^{n-1}\mathrm{d}x = \frac{b^n}{n} - \frac{a^n}{n}$$

Is Integration Straightforward?

- We saw, due to the product and chain rules, that we can differentiate almost anything. Given integration is the anti-derivative can we integrate anything?

- Products and compositions

$$\int f(x)g(x)\mathrm{d}x = ? \qquad \int f(g(x))\mathrm{d}x = ?$$

- Unfortunately, unlike differentiation we don't have a small parameter we can expand in

- In general integration is hard

Integration by Parts

- Recall the product rule $\frac{\mathrm{d}f(x)g(x)}{\mathrm{d}x} = \frac{\mathrm{d}f(x)}{\mathrm{d}x}g(x) + f(x)\frac{\mathrm{d}g(x)}{\mathrm{d}x}$
- Integrating we get

$$\int_a^b \frac{\mathrm{d}f(x)g(x)}{\mathrm{d}x}\mathrm{d}x = \int_a^b \frac{\mathrm{d}f(x)}{\mathrm{d}x}g(x)\mathrm{d}x + \int_a^b f(x)\frac{\mathrm{d}g(x)}{\mathrm{d}x}\mathrm{d}x$$
$$= [f(x)g(x)]_a^b = f(b)g(b) - f(a)g(a)$$

- Unfortunately we get two integrals, but we can turn this around

$$\int_a^b f(x)\frac{\mathrm{d}g(x)}{\mathrm{d}x}\mathrm{d}x = [f(x)g(x)]_a^b - \int_a^b \frac{\mathrm{d}f(x)}{\mathrm{d}x}g(x)\mathrm{d}x$$

whether this is helpful depends on $f(x)$ and $g(x)$

Example of Integration by Parts

- Consider

$$\Pi(z) = \int_0^\infty x^z \mathrm{e}^{-x} \mathrm{d}x = \int_0^\infty x^z \frac{\mathrm{d}(-\mathrm{e}^{-x})}{\mathrm{d}x} \mathrm{d}x$$
$$= \left[x^z(-\mathrm{e}^{-x})\right]_0^\infty - \int_0^\infty \frac{\mathrm{d}x^z}{\mathrm{d}x}(-\mathrm{e}^{-x})\mathrm{d}x$$
$$= \int_0^\infty (zx^{z-1})\mathrm{e}^{-x}\mathrm{d}x = z\int_0^\infty x^{z-1}\mathrm{e}^{-x}\mathrm{d}x = z\Pi(z-1)$$

- Thus $\Pi(z) = z\Pi(z-1)$, but

$$\Pi(0) = \int_0^\infty \mathrm{e}^{-z}\mathrm{d}z = \left[-\mathrm{e}^{-x}\right]_0^\infty = -\mathrm{e}^{-\infty} - (-\mathrm{e}^0) = 1$$

- Now
$\Pi(n) = n\Pi(n-1) = n(n-1)\Pi(n-2) = n(n-1)(n-2)\ldots 1 = n!$

Differentiating Through the Integral

- A trick that sometimes works is differentiating through an integral, e.g. consider finding moments

$$M_n = \mathbb{E}[X^n] = \int_{-\infty}^{\infty} x^n f_X(x) \mathrm{d}x$$

- We can define a moment generating function

$$Z(\ell) = \int_{-\infty}^{\infty} \mathrm{e}^{\ell x} f_X(x) \mathrm{d}x$$

- Then $M_n = Z^{(n)}(0)$ (the nth derivative of $Z(\ell)$ at $\ell = 0$)

$$\left.\frac{\mathrm{d}^n Z(\ell)}{\mathrm{d}\ell^n}\right|_{\ell=0} = \int_{-\infty}^{\infty} \left.\frac{\mathrm{d}^n \mathrm{e}^{\ell x}}{\mathrm{d}\ell^n}\right|_{\ell=0} f_X(x) \mathrm{d}x = \int_{-\infty}^{\infty} x^n f_X(x) \mathrm{d}x = M_n$$

Cumulant Generating Function

- Note that $\mathrm{e}^{\ell x} = 1 + \ell x + \frac{1}{2}\ell^2 x^2 + \frac{1}{3!}\ell^3 x^3 + \cdots$
- So

$$Z(\ell) = \int_{-\infty}^{\infty} \mathrm{e}^{\ell x} f_X(x)\mathrm{d}x = 1 + \ell M_1 + \frac{1}{2}\ell^2 M_2 + \frac{1}{3!}\ell^3 M_3 + \cdots$$

- Now using $\log(1+\epsilon) = \epsilon - \frac{1}{2}\epsilon^2 + \frac{1}{3}\epsilon^3 + \cdots$

$$G(\ell) = \log(Z(\ell)) = \ell M_1 + \frac{1}{2}\ell^2\left(M_2 - M_1^2\right) + \frac{1}{3!}\ell^3\left(M_3 - 3M_2M_1 + 2M_1^3\right) + \cdots$$

- So that $\kappa_n = G^{(n)}(0)$, with $\kappa_1 = M_1$ (the mean), $\kappa_2 = M_2 - M_1^2$ (the variance), $\kappa_3 = M_3 - 3M_2M_1 + 2M_1^3$ (the third cumulant related to the skewness)

More Integration

- Although we have a few tricks, integration is hard.
- Surprisingly integration sometimes is easier when carried out in the complex plane.
- This is a beautiful part of mathematics (due largely to Cauchy)—but beyond the scope of this course.
- Interestingly, also there is an algorithm that allows us to integrate a lot of function. It is sufficiently complicated that you need to write a computer algorithm of considerable complexity to implement it. Most symbolic manipulation packages (e.g. Mathematica) have implemented some part of this algorithm.

Special Functions


- There are integrals with no known closed form solution
- We saw that $\Pi(z) = \int_0^\infty x^z \mathrm{e}^{-x} \mathrm{d}x$ satisfies $\Pi(z) = z\Pi(z-1)$
- For integer n then $\Pi(n) = n!$, but for general z, the integal $\Pi(z)$ can't be written in terms of elementary functions
- We consider $\Pi(z)$ as a special function in its own right
- Although, history has left us with the gamma function instead

$$\Gamma(z) = \int_0^\infty x^{z-1} \mathrm{e}^{-x} \mathrm{d}x = \Pi(z-1)$$

- Other special function defined by integrals exist (e.g. the Bessel , Aire, hypergeometric, elliptic, error functions, . . .)

Outline

1. Defining Integrals
2. Doing Integrals
3. **Changing Variables**



Substitution

- We can make a transformation from x to $u = u(x)$

$$\int_a^b f(x)\mathrm{d}x = \lim_{\delta x_i \to 0} \sum_{i=0}^{n-1} f(x_i)(x_{i+1} - x_i)$$

$$= \lim_{\delta u_i \to 0} \sum_{i=0}^{n-1} f(x(u_i)) \frac{x(u_{i+1}) - x(u_i)}{u_{i+1} - u_i}(u_{i+1} - u_i)$$

$$= \int_{u(a)}^{u(b)} f(x(u)) \frac{\mathrm{d}x(u)}{\mathrm{d}u}\mathrm{d}u$$

  - ⋆ where u_i is such that $x(u_i) = x_i$ or $u_i = u(x_i)$ where $u(x)$ is the inverse of $x(u)$
  - ⋆ using $\lim_{\delta u_i \to 0} \frac{x(u_{i+1}) - x(u_i)}{u_{i+1} - u_i} = \frac{\mathrm{d}x(u_i)}{\mathrm{d}u}$

Substitution in Practice

- A recipe
  1. Choose $u(x)$: often an "inner" function whose derivative also appears in the integrand
  2. Write $\mathrm{d}u = u'(x)\mathrm{d}x$ and replace every x by u
  3. Change the limits: $x = a$ becomes $u = u(a)$
  4. Integrate in u

- Example: $\displaystyle\int_0^1 2x(1+x^2)^3\mathrm{d}x$

$$u = 1 + x^2, \quad \mathrm{d}u = 2x\mathrm{d}x, \quad x : 0 \to 1 \;\Rightarrow\; u : 1 \to 2$$

$$\int_0^1 2x(1+x^2)^3\mathrm{d}x = \int_1^2 u^3\mathrm{d}u = \left[\frac{u^4}{4}\right]_1^2 = \frac{16-1}{4} = \frac{15}{4}$$

More Substitutions

- The limits can swap over: for $\int_0^{\pi/2} \sin(x)\cos^2(x)\mathrm{d}x$ take $u = \cos(x)$, $\mathrm{d}u = -\sin(x)\mathrm{d}x$, and $x : 0 \to \pi/2$ gives $u : 1 \to 0$

$$\int_0^{\pi/2} \sin(x)\cos^2(x)\mathrm{d}x = \int_1^0 -u^2\mathrm{d}u = \int_0^1 u^2\mathrm{d}u = \frac{1}{3}$$

- A linear change of variable stretches $\mathrm{d}x$: with $z = (x-\mu)/\sigma$, $\mathrm{d}x = \sigma\mathrm{d}z$

$$\int_{-\infty}^{\infty} \mathrm{e}^{-(x-\mu)^2/(2\sigma^2)}\mathrm{d}x = \sigma\int_{-\infty}^{\infty} \mathrm{e}^{-z^2/2}\mathrm{d}z$$

- The factor σ is the one-dimensional version of the Jacobian determinant (next slides)

Example of Integration by Substitution

- We consider $I(n) = \int_0^\infty x^n \mathrm{e}^{-x^2/2} \mathrm{d}x$

- Let $u(x) = x^2/2$ or $x(u) = \sqrt{2u}$ so that

$$\frac{\mathrm{d}x(u)}{\mathrm{d}u} = \frac{1}{\sqrt{2u}} \qquad u(0) = 0 \qquad u(\infty) = \infty$$

- Thus

$$I(n) = \int_0^\infty \left(\sqrt{2u}\right)^n \mathrm{e}^{-u} \frac{1}{\sqrt{2u}} \mathrm{d}u$$
$$= 2^{\frac{n-1}{2}} \int_0^\infty u^{\frac{n-1}{2}} \mathrm{e}^{-u} \mathrm{d}u = 2^{\frac{n-1}{2}} \Pi\left(\frac{n-1}{2}\right)$$

- $I(1) = 1$, $I(3) = 2 \times 1! = 2$, $I(5) = 2^2 \times 2! = 8$, but $I(0) = \Pi(-1/2)/\sqrt{2}$, $I(2) = \sqrt{2}\Pi(1/2) = \Pi(-1/2)/\sqrt{2}$

Changing Variables in Multidimensional Space

- When changing variables in many dimensions $\boldsymbol{x} \rightarrow \boldsymbol{u}$ the change of variables involves the Jacobian


$$\int f(\boldsymbol{x})\mathrm{d}\boldsymbol{x} = \int f(\boldsymbol{x}(\boldsymbol{u}))|\det(\mathbf{J})|\mathrm{d}\boldsymbol{u}, \qquad \boldsymbol{J} = \begin{pmatrix} \frac{\partial x_1}{\partial u_1} & \frac{\partial x_1}{\partial u_2} & \cdots & \frac{\partial x_1}{\partial u_n} \\ \frac{\partial x_2}{\partial u_1} & \frac{\partial x_2}{\partial u_2} & \cdots & \frac{\partial x_2}{\partial u_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial u_1} & \frac{\partial x_n}{\partial u_2} & \cdots & \frac{\partial x_n}{\partial u_n} \end{pmatrix}$$

- E.g. transforming from Cartesian coordinates (x,y) to polar coordinates (r,θ) then $x = r\cos(\theta)$ and $y = r\sin(\theta)$

$$|\det(\mathbf{J})| = \left|\det\begin{pmatrix} \frac{\partial r\cos(\theta)}{\partial r} & \frac{\partial r\cos(\theta)}{\partial \theta} \\ \frac{\partial r\sin(\theta)}{\partial r} & \frac{\partial r\sin(\theta)}{\partial \theta} \end{pmatrix}\right| = \left|\det\begin{pmatrix} \cos(\theta) & -r\sin(\theta) \\ \sin(\theta) & r\cos(\theta) \end{pmatrix}\right|$$
$$= r\left(\cos^2(\theta) + \sin^2(\theta)\right) = r$$

- That is, $\mathrm{d}x\mathrm{d}y = r\mathrm{d}r\mathrm{d}\theta$

Change of Variables in Pictures



Summary

- Integration is extra-ordinarily useful as a tool of analysis
- It occurs when you work with probabilities densities for continuous random variables
- Integration is beautiful, but hard—often impossible
- Changing variables turns many hard integrals into easy ones
- Normal distributions, luckily, give integrals that can be done in closed form, although it takes some work: that is the next lecture
- Making friends with integration will give you a super-power that not too many people share