

Name:	Student ID:
-------	-------------

VECTOR SPACE PROBLEM SHEET

This paper asks you to prove some well known results. Although the algebra is easy the proofs are not entirely straightforward. There are marks assigned to the readability of the solution and also how well laid out and explained the steps you make are. (A good proof needs to be easy to follow: you need not comment on trivial algebra, but there should not be steps that are difficult to follow).

This looks very mathematical, but it helps to develop the tools and language that is used to describe machine learning.

1 An inner product $\langle x, y \rangle$ between vectors in a vector space $\mathcal{V}$ satisfies the following properties

- (a) $\langle x, x \rangle \geq 0$ for all $x \in \mathcal{V}$
- (b) $\langle x, x \rangle = 0$ if and only if $x = 0$
- (c) $\langle \alpha x, y \rangle = \alpha \langle x, y \rangle$
- (d) $\langle x, y + z \rangle = \langle x, y \rangle + \langle x, z \rangle$
- (e) $\langle x, y \rangle = \langle y, x \rangle$

The question explores properties of inner products.

$$q(t) = \langle \mathbf{x} + t\mathbf{y}, \mathbf{x} + t\mathbf{y} \rangle$$

Expand $q(t)$ in the form $q(t) = At^2 + 2Bt + C$ to find A , B and C . For $q(t)$ to not change sign its roots (values of t such that $q(t) = 0$) must be complex (i.e. have an imaginary part), or possible have a double root. If there is a value of t such that $q(t) = 0$. Use the standard solutions to a quadratic of the form $q(t) = At^2 + 2Bt + C$ to show that for our $q(t)$ to never become negative then

$$\langle x, y \rangle^2 \leq \langle x, x \rangle \langle y, y \rangle.$$

This is the famous Cauchy-Schwarz inequality written in a very general form. [10 marks]

[illegible]

- (b) From an inner product we can define a norm $\|x\| = \sqrt{\langle x, x \rangle}$. This clearly satisfies non-negativity and linearity. The only non-trivial property to show is that this norm satisfies the triangular inequality. Expand out $\|x + y\|$ and hence show that

$$\|x + y\|^2 = \|x\|^2 + \|y\|^2 + 2\langle x, y \rangle$$

Then use the Cauchy-Schwarz inequality to prove the triangular inequality.

[10 marks]

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There is no handwriting or other markings on the paper.

10

End of question 1

(a) $\frac{\quad}{10}$ (b) $\frac{\quad}{10}$ Total $\frac{\quad}{20}$

2 Random variables, X , Y , etc. form a vector space (i.e. they satisfy properties such as closure under addition and scalar multiplication). Furthermore we can define an inner product between random variables as

$$\langle X, Y \rangle = \mathbb{E}[XY]$$

(i.e. the expectation of the random variable $Z = XY$)

(a) Use the Cauchy-Schwarz inequality to show that

$$\text{Cov}(X,Y)^2 \leq \text{Var}(X)\text{Var}(Y)$$

hence show that the Pearson correlation is between -1 and 1. [10 marks]

[illegible]

(b) Show that for vectors $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ that the inner product

$$\langle \mathbf{x}, \mathbf{y} \rangle = \sum_{i=1}^n w_i x_i y_i$$

satisfies the condition of an inner product provided $w_i > 0$ for all i . Write down the norm and distance induced by this inner product and provide an interpretation of what this distance means.

[10 marks]

10

End of question 2

(a) $\frac{\quad}{10}$	(b) $\frac{\quad}{10}$	Total $\frac{\quad}{20}$
------------------------	------------------------	--------------------------