


Mathematics of Machine Learning


Introduction



Course Details and Topics

Outline

1. **Why are you doing this course?**
2. What we assume that you know



Orientation

- This course addresses three cohorts of students:
 - ★ Third year Artificial Intelligence Students
 - ★ MSc in Machine Learning Students
 - ★ Volunteers
- It is dual coded (AICE3001/AICE6002), with subtle difference in expectations reflected in slightly different evaluation
- Notes are on Moodle and
<https://ecs-vlc.github.io/aice3001/>

Orientation

- This courses addresses three cohorts of students:
 - ★ Third year Artificial Intelligence Students
 - ★ MSc in Machine Learning Students
 - ★ Volunteers
- It is dual coded (AICE3001/AICE6002), with subtle difference in expectations reflected in slightly different evaluation
- Notes are on Moodle and
<https://ecs-vlc.github.io/aice3001/>

Orientation

- This course addresses three cohorts of students:
 - ★ Third year Artificial Intelligence Students
 - ★ MSc in Machine Learning Students
 - ★ Volunteers
- It is dual coded (AICE3001/AICE6002), with subtle difference in expectations reflected in slightly different evaluation
- Notes are on Moodle and
<https://ecs-vlc.github.io/aice3001/>

Why are We Teaching You This?

- In a world where LLMs can do everything, why bother with learning mathematics?

Why are We Teaching You This?

- In a world where LLMs can do everything, why bother with learning mathematics?
- You need to learn to trust your own judgement

Why are We Teaching You This?

- In a world where LLMs can do everything, why bother with learning mathematics?
- You need to learn to trust your own judgement
- LLMs can help us a lot, but if you don't trust yourself you are disempowered

Why are We Teaching You This?

- In a world where LLMs can do everything, why bother with learning mathematics?
- You need to learn to trust your own judgement
- LLMs can help us a lot, but if you don't trust yourself you are disempowered
- This course is aimed at building up trust in your understanding of how machine learning works

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition—we can't cheat
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition—we can't cheat **that much**
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition—we can't cheat that much
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition—we can't cheat that much
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Cracking the Code

- Mathematics is a language that takes time to master
- For some it is easy, for others it will be a struggle
- But given determination everyone can crack the code and understand what the mathematics is saying
- This comes from engaging and working with maths

Cracking the Code

- Mathematics is a language that takes time to master
- For some it is easy, for others it will be a struggle
- But given determination everyone can crack the code and understand what the mathematics is saying
- This comes from engaging and working with maths

Cracking the Code

- Mathematics is a language that takes time to master
- For some it is easy, for others it will be a struggle
- But given determination everyone can crack the code and understand what the mathematics is saying
- This comes from engaging and working with maths

Cracking the Code

- Mathematics is a language that takes time to master
- For some it is easy, for others it will be a struggle
- But given determination everyone can crack the code and understand what the mathematics is saying
- This comes from engaging and working with maths

Cracking the Code

- Mathematics is a language that takes time to master
- For some it is easy, for others it will be a struggle
- But given determination everyone can crack the code and understand what the mathematics is saying
- This comes from engaging and working with maths—this course has tutorials that should be considered compulsory

Cracking the Code

- Mathematics is a language that takes time to master
- For some it is easy, for others it will be a struggle
- But given determination everyone can crack the code and understand what the mathematics is saying
- This comes from engaging and working with maths—this course has tutorials that should be considered compulsory **not because there are marks attached, but because without doing the work you aren't going to crack the code**

Starting from Different Places

- You all start from different places
- Some of you may have a degree in mathematics
- For others it may have been your least favourite subject
- I invite strong mathematicians to become even stronger
- For students starting from a weak base, judge yourselves by your progress, not by comparing yourself to others

Starting from Different Places

- You all start from different places
- Some of you may have a degree in mathematics
- For others it may have been your least favourite subject
- I invite strong mathematicians to become even stronger
- For students starting from a weak base, judge yourselves by your progress, not by comparing yourself to others

Starting from Different Places

- You all start from different places
- Some of you may have a degree in mathematics
- For others it may have been your least favourite subject
- I invite strong mathematicians to become even stronger
- For students starting from a weak base, judge yourselves by your progress, not by comparing yourself to others

Starting from Different Places

- You all start from different places
- Some of you may have a degree in mathematics
- For others it may have been your least favourite subject
- I invite strong mathematicians to become even stronger
- For students starting from a weak base, judge yourselves by your progress, not by comparing yourself to others

Starting from Different Places

- You all start from different places
- Some of you may have a degree in mathematics
- For others it may have been your least favourite subject
- I invite strong mathematicians to become even stronger—the best way to learn is through teaching others
- For students starting from a weak base, judge yourselves by your progress, not by comparing yourself to others

Starting from Different Places

- You all start from different places
- Some of you may have a degree in mathematics
- For others it may have been your least favourite subject
- I invite strong mathematicians to become even stronger—the best way to learn is through teaching others
- For students starting from a weak base, judge yourselves by your progress, not by comparing yourself to others

The Golden Rule of Mathematics

- Understand every step
- When you don't understand a step, stop
- Mathematics uses a lot of short-cuts, which speeds-up calculations

The Golden Rule of Mathematics

- Understand every step
- When you don't understand a step, stop
- Mathematics uses a lot of short-cuts, which speeds-up calculations

The Golden Rule of Mathematics

- Understand every step
- When you don't understand a step, stop, break the problem down
- Mathematics uses a lot of short-cuts, which speeds-up calculations

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, **ask someone**
- Mathematics uses a lot of short-cuts, which speeds-up calculations

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, ask someone
- **Mathematics uses a lot of short-cuts**, which speeds-up calculations

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, ask someone
- Mathematics uses a lot of short-cuts, **which speeds-up calculations**

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, ask someone
- Mathematics uses a lot of short-cuts, which speeds-up calculations, **but you need to be able to justify every step**

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, ask someone
- Mathematics uses a lot of short-cuts, which speeds-up calculations, but you need to be able to justify every step
- It takes bravery to admit you don't understand and insist that you will

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, ask someone
- Mathematics uses a lot of short-cuts, which speeds-up calculations, but you need to be able to justify every step
- It takes bravery to admit you don't understand and insist that you will
- Asking “stupid” questions isn't stupid

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, ask someone
- Mathematics uses a lot of short-cuts, which speeds-up calculations, but you need to be able to justify every step
- It takes bravery to admit you don't understand and insist that you will
- Asking “stupid” questions isn't stupid **it is how we all learn**

The Journey

- We are going to visit lots of aspects of mathematics that have proved fruitful in machine learning
- Some of this will be very concrete, while in other cases by look at a higher level of abstraction we can generalise methods across different application areas
- We will cover, calculus, vector spaces, convexity, optimisation, entropy, probability, statistics and much else besides
- The journey will pay you off, these are the foundations of most of machine learning and will give you super-powers that are hard to acquire without mathematics

The Journey

- We are going to visit lots of aspects of mathematics that have proved fruitful in machine learning
- Some of this will be very concrete, while in other cases by look at a higher level of abstraction we can generalise methods across different application areas
- We will cover, calculus, vector spaces, convexity, optimisation, entropy, probability, statistics and much else besides
- The journey will pay you off, these are the foundations of most of machine learning and will give you super-powers that are hard to acquire without mathematics

The Journey

- We are going to visit lots of aspects of mathematics that have proved fruitful in machine learning
- Some of this will be very concrete, while in other cases by look at a higher level of abstraction we can generalise methods across different application areas
- We will cover, calculus, vector spaces, convexity, optimisation, entropy, probability, statistics and much else besides
- The journey will pay you off, these are the foundations of most of machine learning and will give you super-powers that are hard to acquire without mathematics

The Journey


- We are going to visit lots of aspects of mathematics that have proved fruitful in machine learning
- Some of this will be very concrete, while in other cases by look at a higher level of abstraction we can generalise methods across different application areas
- We will cover, calculus, vector spaces, convexity, optimisation, entropy, probability, statistics and much else besides
- **The journey will pay you off**, these are the foundations of most of machine learning and will give you super-powers that are hard to acquire without mathematics

The Journey

- We are going to visit lots of aspects of mathematics that have proved fruitful in machine learning
- Some of this will be very concrete, while in other cases by look at a higher level of abstraction we can generalise methods across different application areas
- We will cover, calculus, vector spaces, convexity, optimisation, entropy, probability, statistics and much else besides
- The journey will pay you off, these are the foundations of most of machine learning and will give you super-powers that are hard to acquire without mathematics

Outline

1. Why are you doing this course?
2. **What we assume that you know**



What we Assume you Know

- You should all have A-level mathematics and have done some mathematics at Uni
- You should know what a function is, e.g. $f(x) = x^2 + 3$
- You should be able to do simple algebraic manipulations
- And some elementary calculus

$$f'(x) = \frac{d(x^2 + 3)}{dx} = 2x$$

What we Assume you Know

- You should all have A-level mathematics and have done some mathematics at Uni
- You should know what a function is, e.g. $f(x) = x^2 + 3$
- You should be able to do simple algebraic manipulations
- And some elementary calculus

$$f'(x) = \frac{d(x^2 + 3)}{dx} = 2x$$

What we Assume you Know

- You should all have A-level mathematics and have done some mathematics at Uni
- You should know what a function is, e.g. $f(x) = x^2 + 3$
- You should be able to do simple algebraic manipulations
- And some elementary calculus

$$f'(x) = \frac{d(x^2 + 3)}{dx} = 2x$$

What we Assume you Know

- You should all have A-level mathematics and have done some mathematics at Uni
- You should know what a function is, e.g. $f(x) = x^2 + 3$
- You should be able to do simple algebraic manipulations
- And some elementary calculus

$$f'(x) = \frac{d(x^2 + 3)}{dx} = 2x$$

Vectors

- Ordinary vectors (what mathematicians call elements of $\mathbb{R}^n$) are much used in machine learning and should be familiar
- We will use bold symbols to represent **column vectors**, e.g.

$$\mathbf{x} = \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix}$$

- We denote **row vectors** as the **transpose** of column vectors
 $\mathbf{x}^T = (1.2, 2.3, 4.7)$
- We care whether vectors are row or column vectors as it allows us to write equations involving vectors succinctly and unambiguously

Vectors

- Ordinary vectors (what mathematicians call elements of $\mathbb{R}^n$) are much used in machine learning and should be familiar
- We will use bold symbols to represent **column vectors**, e.g.

$$\mathbf{x} = \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix}$$

- We denote **row vectors** as the **transpose** of column vectors
 $\mathbf{x}^T = (1.2, 2.3, 4.7)$
- We care whether vectors are row or column vectors as it allows us to write equations involving vectors succinctly and unambiguously

Vectors

- Ordinary vectors (what mathematicians call elements of $\mathbb{R}^n$) are much used in machine learning and should be familiar
- We will use bold symbols to represent **column vectors**, e.g.

$$\boldsymbol{x} = \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix}$$

- We denote **row vectors** as the **transpose** of column vectors
 $\boldsymbol{x}^T = (1.2, 2.3, 4.7)$
- We care whether vectors are row or column vectors as it allows us to write equations involving vectors succinctly and unambiguously

Vectors

- Ordinary vectors (what mathematicians call elements of $\mathbb{R}^n$) are much used in machine learning and should be familiar
- We will use bold symbols to represent **column vectors**, e.g.

$$\boldsymbol{x} = \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix}$$

- We denote **row vectors** as the **transpose** of column vectors
 $\boldsymbol{x}^T = (1.2, 2.3, 4.7)$
- We care whether vectors are row or column vectors as it allows us to write equations involving vectors succinctly and unambiguously

Matrices

- Matrices are blocks of number that define linear mappings between vectors, e.g. $\mathbf{M} \in \mathbb{R}^{2 \times 3}$

$$\mathbf{M} = \begin{pmatrix} 1.1 & -2.1 & 6.2 \\ 9.3 & 3.1 & -4.4 \end{pmatrix}$$

- An $m \times n$ matrix (i.e. $\mathbf{M} \in \mathbb{R}^{m \times n}$) has m rows and n columns
- An $m \times n$ matrix acting on a vector in $\mathbb{R}^n$ transforms it to $\mathbb{R}^m$ vector, e.g.

$$\mathbf{M}\mathbf{x} = \begin{pmatrix} 1.1 & -2.1 & 6.2 \\ 9.3 & 3.1 & -4.4 \end{pmatrix} \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix} = \begin{pmatrix} 28.63 \\ -2.39 \end{pmatrix}$$

Matrices

- Matrices are blocks of number that define linear mappings between vectors, e.g. $\mathbf{M} \in \mathbb{R}^{2 \times 3}$

$$\mathbf{M} = \begin{pmatrix} 1.1 & -2.1 & 6.2 \\ 9.3 & 3.1 & -4.4 \end{pmatrix}$$

- An $m \times n$ matrix (i.e. $\mathbf{M} \in \mathbb{R}^{m \times n}$) has m rows and n columns
- An $m \times n$ matrix acting on a vector in $\mathbb{R}^n$ transforms it to $\mathbb{R}^m$ vector, e.g.

$$\mathbf{M}\mathbf{x} = \begin{pmatrix} 1.1 & -2.1 & 6.2 \\ 9.3 & 3.1 & -4.4 \end{pmatrix} \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix} = \begin{pmatrix} 28.63 \\ -2.39 \end{pmatrix}$$

Matrices

- Matrices are blocks of number that define linear mappings between vectors, e.g. $\mathbf{M} \in \mathbb{R}^{2 \times 3}$

$$\mathbf{M} = \begin{pmatrix} 1.1 & -2.1 & 6.2 \\ 9.3 & 3.1 & -4.4 \end{pmatrix}$$

- An $m \times n$ matrix (i.e. $\mathbf{M} \in \mathbb{R}^{m \times n}$) has m rows and n columns
- An $m \times n$ matrix acting on a vector in $\mathbb{R}^n$ transforms it to $\mathbb{R}^m$ vector, e.g.

$$\mathbf{M}\mathbf{x} = \begin{pmatrix} 1.1 & -2.1 & 6.2 \\ 9.3 & 3.1 & -4.4 \end{pmatrix} \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix} = \begin{pmatrix} 28.63 \\ -2.39 \end{pmatrix}$$

Composition of Mappings

- Matrices, $\mathbf{A}$, $\mathbf{B}$, etc. represent mappings between spaces
- Provided the matrices are the correct size we can compose these mappings, e.g. if $\mathbf{A} \in \mathbb{R}^{\ell \times m}$ and $\mathbf{B} \in \mathbb{R}^{m \times n}$ then $\mathbf{AB}$ represents the effect of
 - ★ first applying $\mathbf{B}$ to vectors in $\mathbb{R}^n$ to obtain vectors in $\mathbb{R}^m$
 - ★ and then applying $\mathbf{A}$ to get vectors in $\mathbb{R}^\ell$

$$\mathbf{AB}x = \mathbf{A}(\mathbf{B}x)$$

Composition of Mappings

- Matrices, $\mathbf{A}$, $\mathbf{B}$, etc. represent mappings between spaces
- Provided the matrices are the correct size we can compose these mappings, e.g. if $\mathbf{A} \in \mathbb{R}^{\ell \times m}$ and $\mathbf{B} \in \mathbb{R}^{m \times n}$ then $\mathbf{AB}$ represents the effect of
 - ★ first applying $\mathbf{B}$ to vectors in $\mathbb{R}^n$ to obtain vectors in $\mathbb{R}^m$
 - ★ and then applying $\mathbf{A}$ to get vectors in $\mathbb{R}^\ell$

$$\mathbf{AB}x = \mathbf{A}(\mathbf{B}x)$$

Composition of Mappings

- Matrices, $\mathbf{A}$, $\mathbf{B}$, etc. represent mappings between spaces
- Provided the matrices are the correct size we can compose these mappings, e.g. if $\mathbf{A} \in \mathbb{R}^{\ell \times m}$ and $\mathbf{B} \in \mathbb{R}^{m \times n}$ then $\mathbf{AB}$ represents the effect of
 - ★ first applying $\mathbf{B}$ to vectors in $\mathbb{R}^n$ to obtain vectors in $\mathbb{R}^m$
 - ★ and then applying $\mathbf{A}$ to get vectors in $\mathbb{R}^\ell$

$$\mathbf{AB}x = \mathbf{A}(\mathbf{B}x)$$

Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)


Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

The diagram illustrates matrix multiplication using a visual representation of matrices as grids of squares. The first matrix, representing $\mathbf{A}$, is a 5x5 grid with 5 rows and 5 columns. The second matrix, representing $\mathbf{B}$, is a 5x3 grid with 5 rows and 3 columns. An equals sign follows, leading to the resulting matrix $\mathbf{C}$, which is a 5x3 grid with 5 rows and 3 columns. This visualizes the operation $\mathbf{AB} = \mathbf{C}$.


Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

$$\begin{pmatrix} \text{red} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \end{pmatrix} \begin{pmatrix} \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{red} & \text{gray} \end{pmatrix} = \begin{pmatrix} \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \end{pmatrix}$$

Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

The diagram illustrates matrix multiplication using a grid of squares. The first matrix is 4 rows by 5 columns. The top row consists of 5 red squares, while the other three rows consist of 5 gray squares each. The second matrix is 5 rows by 3 columns. The first four rows consist of 3 gray squares each, and the fifth row consists of 3 red squares. An equals sign follows, leading to the resulting matrix, which is 4 rows by 3 columns. The top-right element (row 1, column 3) is a red square, while all other elements are gray squares.

Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

The diagram illustrates matrix multiplication using a grid of squares. The first matrix is a 4x5 grid with gray squares, where the second row, first column square is red. The second matrix is a 5x3 grid with gray squares, where the first, second, third, fourth, and fifth rows, first column squares are red. An equals sign follows, and the resulting matrix is a 4x3 grid with gray squares, where the second row, first column square is red.

Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

The diagram illustrates matrix multiplication using a grid of squares. The first matrix is 4 rows by 5 columns. The second matrix is 5 rows by 3 columns. The result is a 4 rows by 3 columns matrix. The red squares indicate the specific row and column being multiplied to produce a single element in the result matrix.

Gray	Gray	Gray	Gray	Gray	=	Gray	Gray	Gray
Red	Red	Red	Red	Red		Gray	Red	Gray
Gray	Gray	Gray	Gray	Gray		Gray	Gray	Gray
Gray	Gray	Gray	Gray	Gray		Gray	Gray	Gray
Gray	Gray	Gray	Gray	Gray		Gray	Gray	Gray


Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

$$\begin{pmatrix} \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{red} & \text{red} & \text{red} & \text{red} & \text{red} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \end{pmatrix} \begin{pmatrix} \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \end{pmatrix} = \begin{pmatrix} \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \end{pmatrix}$$

Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

The diagram illustrates the matrix multiplication $\mathbf{AB} = \mathbf{C}$ using a grid of squares. The first matrix (A) is 5 rows by 5 columns. The second matrix (B) is 5 rows by 3 columns. The resulting matrix (C) is 5 rows by 3 columns. In matrix A, the third row is highlighted in red. In matrix B, the second column is highlighted in red. In matrix C, the second column is highlighted in red. This visualizes how the dot product of the third row of A and the second column of B results in the second element of the third row of C.


Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

$$\begin{pmatrix} \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{red} & \text{red} & \text{red} & \text{red} & \text{red} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \end{pmatrix} \begin{pmatrix} \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \end{pmatrix} = \begin{pmatrix} \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{gray} \end{pmatrix}$$

Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



The diagram illustrates matrix multiplication using a grid of squares. The first matrix is a 4x5 grid with 4 rows and 5 columns. The bottom row is highlighted in red, while the other three rows are gray. The second matrix is a 5x3 grid with 5 rows and 3 columns. The first column is highlighted in red, while the other two columns are gray. An equals sign follows, leading to the resulting 4x3 matrix. The bottom row of this matrix is highlighted in red, while the other three rows are gray.


Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

The diagram illustrates matrix multiplication using a grid of squares. The first matrix is a 4x5 grid with 4 rows and 5 columns. The bottom row is highlighted in red, while the other three rows are gray. The second matrix is a 5x3 grid with 5 rows and 3 columns. The middle column is highlighted in red, while the other two columns are gray. An equals sign follows, leading to the resulting 4x3 matrix. This matrix has 4 rows and 3 columns. The bottom-middle square (row 4, column 2) is highlighted in red, while the other squares are gray.

Matrix Multiplication


- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



The diagram illustrates matrix multiplication using a visual representation of matrices as grids of squares. The first matrix is a 4x5 grid with 4 rows and 5 columns. The first three rows consist of 5 gray squares each. The fourth row consists of 5 red squares. The second matrix is a 5x3 grid with 5 rows and 3 columns. The first four rows consist of 2 gray squares followed by 1 red square. The fifth row consists of 2 gray squares followed by 1 red square. An equals sign follows, and the resulting matrix is a 4x3 grid with 4 rows and 3 columns. The first three rows consist of 3 gray squares each. The fourth row consists of 2 gray squares followed by 1 red square.

Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



- Matrix multiplication is associative: $(\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC})$

Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

The diagram illustrates matrix multiplication using a grid of squares. The first matrix is 4 rows by 5 columns, with 4 gray squares in each row and 1 red square in each row. The second matrix is 5 rows by 3 columns, with 4 gray squares in each row and 1 red square in each row. An equals sign follows, and the resulting matrix is 4 rows by 3 columns, with 4 gray squares in each row and 1 red square in each row.

- Matrix multiplication is associative: $(\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC})$
- But in general not commutative $\mathbf{AB} \neq \mathbf{BA}$

Transpose

- We often encounter transposes of a matrix $\mathbf{M}^T$ where we have exchanged the rows and columns: $[\mathbf{M}^T]_{ij} = [\mathbf{M}]_{ji}$
- Clear $(\mathbf{M}^T)^T = \mathbf{M}$
- It is simple question of book keeping to show that

$$(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$$

- For a symmetric matrix $\mathbf{M} = \mathbf{M}^T$
- Since 1×1 matrices are always symmetric

$$\mathbf{a}^T \mathbf{b} = (\mathbf{a}^T \mathbf{b})^T = \mathbf{b}^T \mathbf{a}$$

Transpose

- We often encounter transposes of a matrix $\mathbf{M}^T$ where we have exchanged the rows and columns: $[\mathbf{M}^T]_{ij} = [\mathbf{M}]_{ji}$
- Clear $(\mathbf{M}^T)^T = \mathbf{M}$
- It is simple question of book keeping to show that

$$(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$$

- For a symmetric matrix $\mathbf{M} = \mathbf{M}^T$
- Since 1×1 matrices are always symmetric

$$\mathbf{a}^T \mathbf{b} = (\mathbf{a}^T \mathbf{b})^T = \mathbf{b}^T \mathbf{a}$$

Transpose

- We often encounter transposes of a matrix $\mathbf{M}^\top$ where we have exchanged the rows and columns: $[\mathbf{M}^\top]_{ij} = [\mathbf{M}]_{ji}$
- Clear $(\mathbf{M}^\top)^\top = \mathbf{M}$
- It is simple question of book keeping to show that

$$(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$$

- For a symmetric matrix $\mathbf{M} = \mathbf{M}^\top$
- Since 1×1 matrices are always symmetric

$$\mathbf{a}^\top \mathbf{b} = (\mathbf{a}^\top \mathbf{b})^\top = \mathbf{b}^\top \mathbf{a}$$

Transpose

- We often encounter transposes of a matrix $\mathbf{M}^\top$ where we have exchanged the rows and columns: $[\mathbf{M}^\top]_{ij} = [\mathbf{M}]_{ji}$
- Clear $(\mathbf{M}^\top)^\top = \mathbf{M}$
- It is simple question of book keeping to show that

$$(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$$

- For a symmetric matrix $\mathbf{M} = \mathbf{M}^\top$
- Since 1×1 matrices are always symmetric

$$\mathbf{a}^\top \mathbf{b} = (\mathbf{a}^\top \mathbf{b})^\top = \mathbf{b}^\top \mathbf{a}$$

Transpose

- We often encounter transposes of a matrix $\mathbf{M}^T$ where we have exchanged the rows and columns: $[\mathbf{M}^T]_{ij} = [\mathbf{M}]_{ji}$
- Clear $(\mathbf{M}^T)^T = \mathbf{M}$
- It is simple question of book keeping to show that

$$(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$$

- For a symmetric matrix $\mathbf{M} = \mathbf{M}^T$
- Since 1×1 matrices are always symmetric

$$\mathbf{a}^T \mathbf{b} = (\mathbf{a}^T \mathbf{b})^T = \mathbf{b}^T \mathbf{a}$$

Transpose

- We often encounter transposes of a matrix $\mathbf{M}^\top$ where we have exchanged the rows and columns: $[\mathbf{M}^\top]_{ij} = [\mathbf{M}]_{ji}$
- Clear $(\mathbf{M}^\top)^\top = \mathbf{M}$
- It is simple question of book keeping to show that

$$(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$$

- For a symmetric matrix $\mathbf{M} = \mathbf{M}^\top$
- Since 1×1 matrices are always symmetric

$$\begin{aligned} \mathbf{a}^\top \mathbf{b} &= (\mathbf{a}^\top \mathbf{b})^\top = \mathbf{b}^\top \mathbf{a} \\ \mathbf{a}^\top \mathbf{M} \mathbf{b} &= \mathbf{b}^\top \mathbf{M}^\top \mathbf{a} \end{aligned}$$

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix
- Then matrix-vector multiplication can be treated the same as matrix multiplication

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix
- Then matrix-vector multiplication can be treated the same as matrix multiplication

The diagram illustrates matrix-vector multiplication as matrix multiplication. It shows a 4x5 matrix (represented by a grid of 20 gray squares) multiplied by a 5x1 column vector (represented by a single column of 5 gray squares). The result is a 4x1 column vector (represented by a single column of 4 gray squares). The entire expression is enclosed in large parentheses, with an equals sign indicating the result.

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix
- Then matrix-vector multiplication can be treated the same as matrix multiplication

The diagram illustrates matrix-vector multiplication as matrix multiplication. It shows a 4x5 matrix (represented by a grid of 20 squares, with the top row highlighted in red) multiplied by a 5x1 column vector (represented by a single column of 5 red squares). This is followed by an equals sign and a 4x1 column vector (represented by a single column of 4 squares, with the top square highlighted in red). The entire expression is enclosed in large parentheses.

$$\begin{pmatrix} \text{Red} & \text{Red} & \text{Red} & \text{Red} & \text{Red} \\ \text{Gray} & \text{Gray} & \text{Gray} & \text{Gray} & \text{Gray} \\ \text{Gray} & \text{Gray} & \text{Gray} & \text{Gray} & \text{Gray} \\ \text{Gray} & \text{Gray} & \text{Gray} & \text{Gray} & \text{Gray} \end{pmatrix} \begin{pmatrix} \text{Red} \\ \text{Red} \\ \text{Red} \\ \text{Red} \\ \text{Red} \end{pmatrix} = \begin{pmatrix} \text{Red} \\ \text{Gray} \\ \text{Gray} \\ \text{Gray} \end{pmatrix}$$

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix
- Then matrix-vector multiplication can be treated the same as matrix multiplication

The diagram illustrates matrix-vector multiplication as matrix multiplication. On the left, a 5x5 matrix is shown with gray squares in all positions except the second row, which contains five red squares. This matrix is multiplied by a 5x1 column vector, also shown with five red squares. An equals sign follows, and on the right is a 5x1 column vector with four gray squares and one red square in the second position, representing the result of the multiplication.

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix
- Then matrix-vector multiplication can be treated the same as matrix multiplication

The diagram illustrates matrix-vector multiplication as matrix multiplication. It shows a 5x5 matrix (represented by a grid of squares, with the third row highlighted in red) multiplied by a 5x1 column vector (represented by a single column of squares, with the third element highlighted in red). The result is a 5x1 column vector (represented by a single column of squares, with the third element highlighted in red). The equation is shown as:

$$\begin{pmatrix} \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{red} & \text{red} & \text{red} & \text{red} & \text{red} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \end{pmatrix} \begin{pmatrix} \text{red} \\ \text{red} \\ \text{red} \\ \text{red} \\ \text{red} \end{pmatrix} = \begin{pmatrix} \text{gray} \\ \text{gray} \\ \text{red} \\ \text{gray} \\ \text{gray} \end{pmatrix}$$

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix
- Then matrix-vector multiplication can be treated the same as matrix multiplication

The diagram illustrates matrix-vector multiplication as matrix multiplication. It shows a 4x5 matrix (represented by a grid of 20 squares, with the bottom row highlighted in red) multiplied by a 5x1 column vector (represented by a single column of 5 red squares). The result is a 4x1 column vector (represented by a single column of 4 squares, with the bottom square highlighted in red). The entire equation is enclosed in large parentheses.

$$\begin{pmatrix} \text{4x5 matrix} \end{pmatrix} \begin{pmatrix} \text{5x1 column vector} \end{pmatrix} = \begin{pmatrix} \text{4x1 column vector} \end{pmatrix}$$

Inner Products

- The dot or inner-product between two vectors with the same number of elements, can be written $\mathbf{a}^T \mathbf{b}$


Inner Products

- The dot or inner-product between two vectors with the same number of elements, can be written $a^T b$

$$\left(\begin{array}{ccccc} \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare \end{array} \right) = \left(\begin{array}{c} \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \end{array} \right) = \left(\blacksquare \right)$$

Inner Products

- The dot or inner-product between two vectors with the same number of elements, can be written $\mathbf{a}^T \mathbf{b}$



Inner Products

- The dot or inner-product between two vectors with the same number of elements, can be written $a^T b$

$$\left(\begin{array}{ccccc} \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare \end{array} \right) = \left(\begin{array}{c} \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \end{array} \right) = \left(\blacksquare \right)$$

Inner Products

- The dot or inner-product between two vectors with the same number of elements, can be written $\mathbf{a}^T \mathbf{b}$

$$\begin{pmatrix} \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare \end{pmatrix} \begin{pmatrix} \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \end{pmatrix} = \begin{pmatrix} \blacksquare \end{pmatrix}$$

- Inner products encode angles between vectors

$$\mathbf{a}^T \mathbf{b} = \|\mathbf{a}\| \|\mathbf{b}\| \cos(\theta)$$

Outer Products

- We can also define the outer-product between any size vectors,
 ab^T

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4 is represented by a large left parenthesis followed by four gray squares stacked vertically, and a large right parenthesis. This is followed by a row vector of size 3, represented by a large left parenthesis followed by three gray squares in a row, and a large right parenthesis. An equals sign follows, and then a 4x3 matrix is shown. The matrix is represented by a large left parenthesis followed by four columns, each containing three gray squares, and a large right parenthesis. This visualizes the operation ab^T where a is a 4x1 vector and b^T is a 1x3 vector, resulting in a 4x3 matrix.


Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product operation ab^T . On the left, a column vector a of size 4x1 is shown, with its top element highlighted in red. Next to it is a row vector b^T of size 1x3, with its first element highlighted in red. An equals sign follows, leading to a 4x3 matrix on the right. This matrix is the result of the outer product, where each row is a scalar multiple of the vector b^T . The top row of the matrix has its first element highlighted in red, corresponding to the red elements in the input vectors.

Outer Products

- We can also define the outer-product between any size vectors, ab^T



Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4 is shown, with its top element highlighted in red. This is multiplied by a row vector of size 3, with its middle element highlighted in red. The result is a 4x3 matrix where every element in the first column is red, representing the scalar product of the original red elements multiplied by each row of the second vector.

$$\begin{pmatrix} \text{red} \\ \text{gray} \\ \text{gray} \\ \text{gray} \end{pmatrix} \begin{pmatrix} \text{gray} & \text{red} & \text{gray} \end{pmatrix} = \begin{pmatrix} \text{red} & \text{red} & \text{red} \\ \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{red} & \text{gray} \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4 is shown, with its top element highlighted in red and the others in gray. This is multiplied by a row vector of size 3, with its rightmost element highlighted in red and the others in gray. The result is a 4x3 matrix where each row is a scaled version of the row vector, and the top row is highlighted in red to show the contribution of the red element in the column vector.

$$\begin{pmatrix} \text{red} \\ \text{gray} \\ \text{gray} \\ \text{gray} \end{pmatrix} \begin{pmatrix} \text{gray} & \text{gray} & \text{red} \end{pmatrix} = \begin{pmatrix} \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product operation ab^T . On the left, a column vector a of size 4x1 is shown, with its second element highlighted in red. This is multiplied by a row vector b^T of size 1x3, with its first element highlighted in red. The result is a 4x3 matrix where each row is a scalar multiple of the vector b^T , and the first row is identical to b^T because the red element in a is 1. The matrix has 4 rows and 3 columns, with the top row containing one red and two gray squares, and the remaining three rows containing three gray squares each.

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4 is shown, with its second element highlighted in red. This is multiplied by a row vector of size 3, with its second element also highlighted in red. The result is a 4x3 matrix where every element in the second row and second column is red, while all other elements are gray.

$$\begin{pmatrix} \square \\ \color{red}\square \\ \square \\ \square \end{pmatrix} \begin{pmatrix} \square & \color{red}\square & \square \end{pmatrix} = \begin{pmatrix} \square & \square & \square \\ \square & \color{red}\square & \square \\ \square & \square & \square \\ \square & \square & \square \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product operation ab^T . On the left, a column vector a is shown as a vertical stack of four squares, with the second square from the top highlighted in red. This is followed by a row vector b^T shown as a horizontal row of three squares, with the third square from the left highlighted in red. An equals sign follows, leading to the resulting 4x3 matrix. This matrix is represented by a large set of parentheses containing a 4x3 grid of squares. The second row and third column of this grid are highlighted in red, representing the outer product of the red squares from the original vectors.

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product operation ab^T . On the left, a column vector a of size 4x1 is shown, with its third element highlighted in red. This is multiplied by a row vector b^T of size 1x3, with its first element highlighted in red. The result is a 4x3 matrix where each row is a scalar multiple of the vector b^T , and the third row is highlighted in red, corresponding to the red element in a .

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two 4x1 vectors. On the left, a column vector of four gray squares is multiplied by a row vector of three gray squares. The result is a 4x4 matrix where the third row and third column are highlighted in red, representing the outer product of the third elements of the input vectors.

$$\begin{pmatrix} \square \\ \square \\ \square \\ \square \end{pmatrix} \begin{pmatrix} \square & \square & \square \end{pmatrix} = \begin{pmatrix} \square & \square & \square \\ \square & \square & \square \\ \square & \square & \square \\ \square & \square & \square \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4 is shown, with its third element highlighted in red. This is multiplied by a row vector of size 3, with its third element also highlighted in red. The result is a 4x3 matrix where every element in the third row and third column is red, while all other elements are gray.

$$\begin{pmatrix} \square \\ \square \\ \color{red}\square \\ \square \end{pmatrix} \begin{pmatrix} \square & \square & \color{red}\square \end{pmatrix} = \begin{pmatrix} \square & \square & \square \\ \square & \square & \square \\ \color{red}\square & \color{red}\square & \color{red}\square \\ \square & \square & \square \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4 is shown, with the bottom element highlighted in red. This is multiplied by a row vector of size 3, with the first element highlighted in red. The result is a 4x3 matrix where each row is a scalar multiple of the original row vector, and the bottom row is highlighted in red.

$$\begin{pmatrix} \square \\ \square \\ \square \\ \color{red}\square \end{pmatrix} \begin{pmatrix} \color{red}\square & \square & \square \end{pmatrix} = \begin{pmatrix} \square & \square & \square \\ \square & \square & \square \\ \square & \square & \square \\ \color{red}\square & \square & \square \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4 is shown, with its bottom element highlighted in red. This is multiplied by a row vector of size 3, with its middle element highlighted in red. The result is a 4x3 matrix where the bottom row of the original column vector is multiplied by the entire row vector, resulting in the second column of the matrix being red.

$$\begin{pmatrix} \square \\ \square \\ \square \\ \color{red}\square \end{pmatrix} \begin{pmatrix} \square & \color{red}\square & \square \end{pmatrix} = \begin{pmatrix} \square & \square & \square \\ \square & \square & \square \\ \square & \square & \square \\ \square & \color{red}\square & \square \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4 is shown, consisting of four gray squares and one red square at the bottom. This is multiplied by a row vector of size 3, consisting of two gray squares and one red square on the right. The result is a 4x3 matrix on the right, where each element is the product of the corresponding elements from the two input vectors. The bottom-right element is red, while all other elements are gray.

$$\begin{pmatrix} \square \\ \square \\ \square \\ \square \\ \color{red}\square \end{pmatrix} \begin{pmatrix} \square & \square & \color{red}\square \end{pmatrix} = \begin{pmatrix} \square & \square & \square \\ \square & \square & \square \\ \square & \square & \square \\ \square & \square & \square \\ \square & \square & \color{red}\square \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of a column vector and a row vector. On the left, a column vector is shown with four elements: three gray squares and one red square at the bottom. This is multiplied by a row vector with three elements: two gray squares and one red square on the right. The result is a 4x3 matrix on the right, where each element is the product of the corresponding elements from the input vectors. The bottom-right element is red, while all other elements are gray.

$$\begin{pmatrix} \square \\ \square \\ \square \\ \color{red}\square \end{pmatrix} \begin{pmatrix} \square & \square & \color{red}\square \end{pmatrix} = \begin{pmatrix} \square & \square & \square \\ \square & \square & \square \\ \square & \square & \square \\ \square & \square & \color{red}\square \end{pmatrix}$$

- Notice that we need to know whether we are dealing with row or column vectors

Norms

- Norms are non-negative numbers associated with the “size” of a vector
- The norm you should be familiar with is the **Euclidean Norm** or 2-norm

$$\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- We will see there exist other norms (using different ways of measuring size)
- The Euclidean norm is intimately related to the inner-product, we will see other examples of this for different vector spaces

Norms

- Norms are non-negative numbers associated with the “size” of a vector
- The norm you should be familiar with is the **Euclidean Norm** or 2-norm

$$\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- We will see there exist other norms (using different ways of measuring size)
- The Euclidean norm is intimately related to the inner-product, we will see other examples of this for different vector spaces

Norms

- Norms are non-negative numbers associated with the “size” of a vector
- The norm you should be familiar with is the **Euclidean Norm** or 2-norm

$$\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2} = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}$$

- We will see there exist other norms (using different ways of measuring size)
- The Euclidean norm is intimately related to the inner-product, we will see other examples of this for different vector spaces

Norms

- Norms are non-negative numbers associated with the “size” of a vector
- The norm you should be familiar with is the **Euclidean Norm** or 2-norm

$$\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2} = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} = \sqrt{\mathbf{x}^\top \mathbf{x}}$$

- We will see there exist other norms (using different ways of measuring size)
- The Euclidean norm is intimately related to the inner-product, we will see other examples of this for different vector spaces

Norms

- Norms are non-negative numbers associated with the “size” of a vector
- The norm you should be familiar with is the **Euclidean Norm** or 2-norm

$$\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2} = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} = \sqrt{\mathbf{x}^\top \mathbf{x}}$$

- We will see there exist other norms (using different ways of measuring size)
- The Euclidean norm is intimately related to the inner-product, we will see other examples of this for different vector spaces

Norms

- Norms are non-negative numbers associated with the “size” of a vector
- The norm you should be familiar with is the **Euclidean Norm** or 2-norm

$$\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2} = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} = \sqrt{\mathbf{x}^\top \mathbf{x}}$$

- We will see there exist other norms (using different ways of measuring size)
- The Euclidean norm is intimately related to the inner-product, we will see other examples of this for different vector spaces

Matrices as Linear Mappings

- Matrices represent mappings between vectors

$$\mathbf{b} = \mathbf{M}\mathbf{a}$$

- The elements of $\mathbf{b}$ are **linear sums** of the elements of $\mathbf{a}$

$$b_i = \sum_j M_{ij} a_j$$

- Two important consequences of this linearity are for any real c

$$\mathbf{M}(c\mathbf{a}) = c(\mathbf{M}\mathbf{a})$$

Matrices as Linear Mappings

- Matrices represent mappings between vectors

$$\mathbf{b} = \mathbf{M}\mathbf{a}$$

- The elements of $\mathbf{b}$ are **linear sums** of the elements of $\mathbf{a}$

$$b_i = \sum_j M_{ij}a_j = M_{i1}a_1 + M_{i2}a_2 + \cdots + M_{in}a_n$$

- Two important consequences of this linearity are for any real c

$$\mathbf{M}(c\mathbf{a}) = c(\mathbf{M}\mathbf{a})$$

Matrices as Linear Mappings

- Matrices represent mappings between vectors

$$\mathbf{b} = \mathbf{M}\mathbf{a}$$

- The elements of $\mathbf{b}$ are **linear sums** of the elements of $\mathbf{a}$

$$b_i = \sum_j M_{ij}a_j = M_{i1}a_1 + M_{i2}a_2 + \cdots + M_{in}a_n$$

- Two important consequences of this linearity are for any real c

$$\mathbf{M}(c\mathbf{a}) = c(\mathbf{M}\mathbf{a})$$

Matrices as Linear Mappings

- Matrices represent mappings between vectors

$$\mathbf{b} = \mathbf{M}\mathbf{a}$$

- The elements of $\mathbf{b}$ are **linear sums** of the elements of $\mathbf{a}$

$$b_i = \sum_j M_{ij}a_j = M_{i1}a_1 + M_{i2}a_2 + \cdots + M_{in}a_n$$


- Two important consequences of this linearity are for any real c

$$\mathbf{M}(c\mathbf{a}) = c(\mathbf{M}\mathbf{a})$$

$$\mathbf{M}(\mathbf{a} + \mathbf{b}) = (\mathbf{M}\mathbf{a}) + (\mathbf{M}\mathbf{b})$$

Action Of Matrices on Volumes


- For a closed set of points corresponding to a volume, V in $\mathbb{R}^n$, treating each point as a vector, we can ask what happens to the volume after mapping the vector by a matrix



- For square matrices, $\mathbf{M}$, the determinant is equal to $\det(\mathbf{M}) = \pm V'/V$
- The sign tells us if there has been a reflection

Action Of Matrices on Volumes


- For a closed set of points corresponding to a volume, V in $\mathbb{R}^n$, treating each point as a vector, we can ask what happens to the volume after mapping the vector by a matrix



- For square matrices, $\mathbf{M}$, the determinant is equal to $\det(\mathbf{M}) = \pm V'/V$
- The sign tells us if there has been a reflection

Action Of Matrices on Volumes

- For a closed set of points corresponding to a volume, V in $\mathbb{R}^n$, treating each point as a vector, we can ask what happens to the volume after mapping the vector by a matrix



- For square matrices, $\mathbf{M}$, the determinant is equal to $\det(\mathbf{M}) = \pm V'/V$
- The sign tells us if there has been a reflection

Determinants

- A consequence of the fact that $\mathbf{AB}$ denotes the composition, apply mapping $\mathbf{B}$ then apply mapping $\mathbf{A}$

$$\det(\mathbf{AB}) = \det(\mathbf{A}) \times \det(\mathbf{B})$$

- Determinants are only defined for square matrices, as non-square matrices project into different size spaces, where the volumes cannot be compared
- **Singular matrices** projects all points onto a lower dimensional space, so have determinant 0

Determinants

- A consequence of the fact that $\mathbf{AB}$ denotes the composition, apply mapping $\mathbf{B}$ then apply mapping $\mathbf{A}$

$$\det(\mathbf{AB}) = \det(\mathbf{A}) \times \det(\mathbf{B})$$

- Determinants are only defined for square matrices, as non-square matrices project into different size spaces, where the volumes cannot be compared
- **Singular matrices** projects all points onto a lower dimensional space, so have determinant 0

Determinants

- A consequence of the fact that $\mathbf{AB}$ denotes the composition, apply mapping $\mathbf{B}$ then apply mapping $\mathbf{A}$

$$\det(\mathbf{AB}) = \det(\mathbf{A}) \times \det(\mathbf{B})$$

- Determinants are only defined for square matrices, as non-square matrices project into different size spaces, where the volumes cannot be compared
- **Singular matrices** projects all points onto a lower dimensional space, so have determinant 0

Multivariate Functions

- In machine learning we heavily use multivariate functions

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

- Often our functions will be scalar value $f : \mathbb{R}^n \rightarrow \mathbb{R}$
- E.g. a learning machine for regression might be $\hat{f}(\mathbf{x}, \mathbf{w})$
- We say a function f is linear if

$$f(c\mathbf{x}) = cf(\mathbf{x}) \qquad f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$$

Multivariate Functions

- In machine learning we heavily use multivariate functions

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

- Often our functions will be scalar value $f : \mathbb{R}^n \rightarrow \mathbb{R}$

- E.g. a learning machine for regression might be $\hat{f}(\mathbf{x}, \mathbf{w})$

- We say a function f is linear if

$$f(c\mathbf{x}) = cf(\mathbf{x}) \qquad f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$$

Multivariate Functions

- In machine learning we heavily use multivariate functions

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

- Often our functions will be scalar value $f : \mathbb{R}^n \rightarrow \mathbb{R}$

- E.g. a learning machine for regression might be $\hat{f}(\mathbf{x}, \mathbf{w})$

- We say a function f is linear if

$$f(c\mathbf{x}) = cf(\mathbf{x}) \qquad f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$$

Multivariate Functions

- In machine learning we heavily use multivariate functions

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

- Often our functions will be scalar value $f : \mathbb{R}^n \rightarrow \mathbb{R}$
- E.g. a learning machine for regression might be $\hat{f}(\mathbf{x}, \mathbf{w})$ or we might have a loss function $L(\mathbf{w})$
- We say a function f is linear if

$$f(c\mathbf{x}) = cf(\mathbf{x}) \qquad f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$$

Multivariate Functions

- In machine learning we heavily use multivariate functions

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

- Often our functions will be scalar value $f : \mathbb{R}^n \rightarrow \mathbb{R}$
- E.g. a learning machine for regression might be $\hat{f}(\mathbf{x}, \mathbf{w})$ or we might have a loss function $L(\mathbf{w})$
- We say a function f is linear if

$$f(c\mathbf{x}) = cf(\mathbf{x}) \qquad f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$$

Gradients

- The gradient of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ at a point $\mathbf{x}^*$ is defined as the column vector

$$\nabla f(\mathbf{x}^*) = \begin{pmatrix} \frac{\partial f(\mathbf{x}^*)}{\partial x_1} \\ \frac{\partial f(\mathbf{x}^*)}{\partial x_2} \\ \vdots \\ \frac{\partial f(\mathbf{x}^*)}{\partial x_n} \end{pmatrix}$$

- The gradient vector is orthogonal to the contour plane and points in the direction in which the function increases fastest

Gradients

- The gradient of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ at a point $\mathbf{x}^*$ is defined as the column vector

$$\nabla f(\mathbf{x}^*) = \begin{pmatrix} \frac{\partial f(\mathbf{x}^*)}{\partial x_1} \\ \frac{\partial f(\mathbf{x}^*)}{\partial x_2} \\ \vdots \\ \frac{\partial f(\mathbf{x}^*)}{\partial x_n} \end{pmatrix}$$

- The gradient vector is orthogonal to the contour plane and points in the direction in which the function increases fastest

Probability

- We will have a lot to say about probability later in this course
- To reason about events with uncertain outcomes we introduce **random variables**
- Random variables corresponds to objects that we assign numbers to depending on the outcome of a random event
- E.g. we might assign 0 to the outcome of tossing a tail and 1 for a head

Probability

- We will have a lot to say about probability later in this course
- To reason about events with uncertain outcomes we introduce **random variables**
- Random variables corresponds to objects that we assign numbers to depending on the outcome of a random event
- E.g. we might assign 0 to the outcome of tossing a tail and 1 for a head

Probability

- We will have a lot to say about probability later in this course
- To reason about events with uncertain outcomes we introduce **random variables**
- Random variables corresponds to objects that we assign numbers to depending on the outcome of a random event
- E.g. we might assign 0 to the outcome of tossing a tail and 1 for a head

Probability

- We will have a lot to say about probability later in this course
- To reason about events with uncertain outcomes we introduce **random variables**
- Random variables corresponds to objects that we assign numbers to depending on the outcome of a random event
- E.g. we might assign 0 to the outcome of tossing a tail and 1 for a head

Probability

- We will have a lot to say about probability later in this course
- To reason about events with uncertain outcomes we introduce **random variables**
- Random variables corresponds to objects that we assign numbers to depending on the outcome of a random event
- E.g. we might assign 0 to the outcome of tossing a tail and 1 for a head—this is an example of a discrete random variable

Probability

- We will have a lot to say about probability later in this course
- To reason about events with uncertain outcomes we introduce **random variables**
- Random variables corresponds to objects that we assign numbers to depending on the outcome of a random event
- E.g. we might assign 0 to the outcome of tossing a tail and 1 for a head—this is an example of a discrete random variable
- A continuous random variable might be the temperature at 12 noon tomorrow in Hyde park

Probability Distributions

- For discrete events, we can define a **probability mass function** $\mathbb{P}(X = x)$ that the random variable takes the value x
- This is a number between 0 and 1 with $\sum_{x \in \mathcal{X}} \mathbb{P}(X = x) = 1$
- For continuous events the probability of X taking a particular value x is zero
- Instead we consider the probability density $f_X(x)$ which we can define as

$$f_X(x) = \lim_{\delta x \rightarrow 0} \frac{\mathbb{P}(x < X \leq x + \delta x)}{\delta x}$$

Probability Distributions

- For discrete events, we can define a **probability mass function** $\mathbb{P}(X = x)$ that the random variable takes the value x
- This is a number between 0 and 1 with $\sum_{x \in \mathcal{X}} \mathbb{P}(X = x) = 1$
- For continuous events the probability of X taking a particular value x is zero
- Instead we consider the probability density $f_X(x)$ which we can define as

$$f_X(x) = \lim_{\delta x \rightarrow 0} \frac{\mathbb{P}(x < X \leq x + \delta x)}{\delta x}$$

Probability Distributions

- For discrete events, we can define a **probability mass function** $\mathbb{P}(X = x)$ that the random variable takes the value x
- This is a number between 0 and 1 with $\sum_{x \in \mathcal{X}} \mathbb{P}(X = x) = 1$
- For continuous events the probability of X taking a particular value x is zero
- Instead we consider the probability density $f_X(x)$ which we can define as

$$f_X(x) = \lim_{\delta x \rightarrow 0} \frac{\mathbb{P}(x < X \leq x + \delta x)}{\delta x}$$

Probability Distributions

- For discrete events, we can define a **probability mass function** $\mathbb{P}(X = x)$ that the random variable takes the value x
- This is a number between 0 and 1 with $\sum_{x \in \mathcal{X}} \mathbb{P}(X = x) = 1$
- For continuous events the probability of X taking a particular value x is zero
- Instead we consider the probability density $f_X(x)$ which we can define as

$$f_X(x) = \lim_{\delta x \rightarrow 0} \frac{\mathbb{P}(x < X \leq x + \delta x)}{\delta x}$$

Expectations

- One of the most useful quantities in dealing with random numbers is to look at the average value, aka as the **expectation**, for some function $g(X)$ of the random variable

$$\mathbb{E}[g(X)] = \begin{cases} \sum_{x \in \mathcal{X}} \mathbb{P}(X = x) g(x) \\ \int f_X(x) g(x) dx \end{cases}$$

- Expectations are linear operators

$$\mathbb{E}[cg(X)] = c\mathbb{E}[g(x)]$$

$$\mathbb{E}[g(X) + h(X)] = \mathbb{E}[g(X)] + \mathbb{E}[h(X)]$$

- The expectation of a constant is itself $\mathbb{E}[c] = c$

Expectations

- One of the most useful quantities in dealing with random numbers is to look at the average value, aka as the **expectation**, for some function $g(X)$ of the random variable

$$\mathbb{E}[g(X)] = \begin{cases} \sum_{x \in \mathcal{X}} \mathbb{P}(X = x) g(x) \\ \int f_X(x) g(x) dx \end{cases}$$

- Expectations are linear operators

$$\mathbb{E}[cg(X)] = c\mathbb{E}[g(x)]$$

$$\mathbb{E}[g(X) + h(X)] = \mathbb{E}[g(X)] + \mathbb{E}[h(X)]$$

- The expectation of a constant is itself $\mathbb{E}[c] = c$

Expectations

- One of the most useful quantities in dealing with random numbers is to look at the average value, aka as the **expectation**, for some function $g(X)$ of the random variable

$$\mathbb{E}[g(X)] = \begin{cases} \sum_{x \in \mathcal{X}} \mathbb{P}(X = x) g(x) \\ \int f_X(x) g(x) dx \end{cases}$$

- Expectations are linear operators

$$\mathbb{E}[cg(X)] = c\mathbb{E}[g(x)]$$

$$\mathbb{E}[g(X) + h(X)] = \mathbb{E}[g(X)] + \mathbb{E}[h(X)]$$

- The expectation of a constant is itself $\mathbb{E}[c] = c$

Lesson

- I've covered material that we assume you know
- If you don't, you are going to struggle following any machine learning course, so grab a maths book and revise this
- We will revisit most of these topics to gain a better understanding, but you will be lost in the other machine learning modules if you are not familiar with everything I've covered
- There are exercises we will go through in the tutorial session where you can test your proficiency

Lesson

- I've covered material that we assume you know
- If you don't, you are going to struggle following any machine learning course, so grab a maths book and revise this
- We will revisit most of these topics to gain a better understanding, but you will be lost in the other machine learning modules if you are not familiar with everything I've covered
- There are exercises we will go through in the tutorial session where you can test your proficiency

Lesson

- I've covered material that we assume you know
- If you don't, you are going to struggle following any machine learning course, so grab a maths book and revise this
- We will revisit most of these topics to gain a better understanding, but you will be lost in the other machine learning modules if you are not familiar with everything I've covered
- There are exercises we will go through in the tutorial session where you can test your proficiency

Lesson

- I've covered material that we assume you know
- If you don't, you are going to struggle following any machine learning course, so grab a maths book and revise this
- We will revisit most of these topics to gain a better understanding, but you will be lost in the other machine learning modules if you are not familiar with everything I've covered
- There are exercises we will go through in the tutorial session where you can test your proficiency