

## Singular Value Decomposition (SVD)

$$\begin{pmatrix} 0 & X \\ X^T & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \sigma \begin{pmatrix} u \\ v \end{pmatrix}$$

Singular Valued Decomposition, SVD, general linear maps

## Singular Valued Decomposition

- One of the most important results in linear algebra is that any matrix  $X$  can be decomposed as

$$X = USV^T$$

- If  $X$  is a  $m \times n$  matrix then  $U$  is a  $m \times m$  orthogonal matrix and  $V$  is an  $n \times n$  orthogonal matrix i.e.  $UU^T = U^T U = I$ ,  $VV^T = V^T V = I$
- And  $S$  is a  $m \times n$  matrix with non-zero diagonal elements  $S_{ii} = s_i$  known as the **singular values**
- Where did that come from?

## Eigenvectors

- Note that as  $Xv = su$  and  $X^T u = sv$  then

$$X(-v) = (-s)u \quad X^T u = (-s)(-v)$$

if  $\begin{pmatrix} u \\ v \end{pmatrix}$  is an eigenvector of  $B$  with eigenvalue  $s$  then so is  $\begin{pmatrix} u \\ -v \end{pmatrix}$  with eigenvalue  $-s$

- If  $n < m$  then  $X^T X$  is not full rank so some eigenvalues are zero
- As a consequence  $m - n$  vectors exist such that  $Xv = 0$
- The eigenvalues and eigenvectors are

$$n \times \left( s_i, \begin{pmatrix} u_i \\ v_i \end{pmatrix} \right) \quad n \times \left( -s_i, \begin{pmatrix} u_i \\ -v_i \end{pmatrix} \right) \quad m - n \times \left( 0, \begin{pmatrix} 0 \\ v_k \end{pmatrix} \right)$$

## Normalisation Subtlety

$$\begin{pmatrix} 0 & X \\ X^T & 0 \end{pmatrix} = \begin{pmatrix} U & U & 0 \\ V & -V & V_0 \end{pmatrix} \begin{pmatrix} \Sigma & 0 & 0 \\ 0 & -\Sigma & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} U^T & V^T \\ U^T & -V^T \\ 0 & V_0^T \end{pmatrix}$$

- Multiplying out we have

$$X = 2USV^T \quad X^T = 2VSU^T$$

- Now the vectors  $u_i$  and  $v_i$  form an orthogonal set as it satisfy

$$X^T X v = s^2 v \quad X X^T u = s^2 u$$

- But they are not normalised (since  $\begin{pmatrix} u_i \\ v_i \end{pmatrix}$  is normalised). If we define  $\tilde{U} = \sqrt{2}U$  and  $\tilde{V} = \sqrt{2}V$  we find

$$X = \tilde{U} S \tilde{V}^T \quad X^T = \tilde{V} S^T \tilde{U}^T$$

- Singular Value Decomposition
- General Linear Mappings
- Linear Regression Revisited

$$\begin{pmatrix} 0 & X \\ X^T & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \sigma \begin{pmatrix} u \\ v \end{pmatrix}$$

## Eigen Decomposition

- Consider an arbitrary  $n \times m$  matrix  $X$ , and construct the  $(n+m) \times (n+m)$  symmetric matrix,  $B$ ,

$$\begin{pmatrix} 0 & X \\ X^T & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \sigma \begin{pmatrix} u \\ v \end{pmatrix}$$

$\begin{pmatrix} u \\ v \end{pmatrix}$  is an eigenvector of  $B$  with eigenvalue  $s$

- We observe that

$$Xv = su \quad X^T u = sv \\ X^T X v = s X^T u = s^2 v \quad X X^T u = s X v = s^2 u$$

## Matrix Decomposition

- Stacking the eigenvectors into a matrix

$$\begin{pmatrix} 0 & X \\ X^T & 0 \end{pmatrix} = \begin{pmatrix} U & U & 0 \\ V & -V & V_0 \end{pmatrix} = \begin{pmatrix} U & U & 0 \\ V & -V & V_0 \end{pmatrix} \begin{pmatrix} \Sigma & 0 & 0 \\ 0 & -\Sigma & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

- Since the vectors  $\begin{pmatrix} u_i \\ v_i \end{pmatrix}$  are eigenvectors of a symmetric matrix they form an orthogonal matrix if they are normalised.
- Multiply on the right by the transpose of the orthogonal matrix

$$\begin{pmatrix} 0 & X \\ X^T & 0 \end{pmatrix} = \begin{pmatrix} U & U & 0 \\ V & -V & V_0 \end{pmatrix} \begin{pmatrix} \Sigma & 0 & 0 \\ 0 & -\Sigma & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} U^T & V^T \\ U^T & -V^T \\ 0 & V_0^T \end{pmatrix}$$

## SVD

- Any matrix,  $X$ , can be written as  $X = USV^T$

- $U, V$  are orthogonal matrices
- $S = \text{diag}(s_1, s_2, \dots, s_n)$

- $s_i$  can always be chosen to be positive and are known as **singular values**
- Singular value decomposition applies to both square and non-square matrices—they describe general linear mappings

## Finding SVD

- Most libraries will compute the SVD for you
- They can do this by choosing the smaller of two matrices  $\mathbf{X}\mathbf{X}^T$  and  $\mathbf{X}^T\mathbf{X}$  and then compute the eigenvalues
- The singular values are the square root of the eigenvalues (notice that  $\mathbf{X}\mathbf{X}^T$  and  $\mathbf{X}^T\mathbf{X}$  are both positive semi-definite so the eigenvalues will be non-negative)
- It can compute the  $\mathbf{U}$  matrix or  $\mathbf{V}$  matrix by multiplying through by  $\mathbf{X}$  or  $\mathbf{X}^T$  ( $\mathbf{U} = \mathbf{X}\mathbf{V}\mathbf{S}^{-1}$  and  $\mathbf{V} = \mathbf{X}^T\mathbf{U}\mathbf{S}^{-1}$ )
- In practice to perform PCA most people subtract the mean from their data and then perform SVD

Adam Prügel-Bennett AICE3001/6002 Mathematics of Machine Learning <https://ecs-vlc.github.io/aice3001/>


## Outline

- Singular Value Decomposition
- General Linear Mappings
- Linear Regression Revisited

$$\begin{pmatrix} 0 & \mathbf{X} \\ \mathbf{X}^T & 0 \end{pmatrix} \begin{pmatrix} \mathbf{u} \\ \mathbf{v} \end{pmatrix} = \sigma \begin{pmatrix} \mathbf{u} \\ \mathbf{v} \end{pmatrix}$$

## Matrices

$$\mathbf{M} = \begin{pmatrix} -0.45 & 1.9 \\ -0.77 & -0.025 \end{pmatrix} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T = \begin{pmatrix} \cos(-175) & \sin(-175) \\ -\sin(-175) & \cos(-175) \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 0 & 0.75 \end{pmatrix} \begin{pmatrix} \cos(75) & \sin(75) \\ -\sin(75) & \cos(75) \end{pmatrix}$$



Adam Prügel-Bennett AICE3001/6002 Mathematics of Machine Learning <https://ecs-vlc.github.io/aice3001/>

## Non-Square Matrices

- When the matrices are non-square then the matrix of singular value matrix will either
  - Squash some directions to zero
  - Introduce new dimensions orthogonal to the vector

$$\begin{pmatrix} \mathbf{X} \\ \mathbf{X}^T \end{pmatrix} = \begin{pmatrix} \mathbf{U} \\ \mathbf{V} \end{pmatrix} \begin{pmatrix} \mathbf{S} \\ \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{V}^T \\ \mathbf{U}^T \end{pmatrix}$$

- The rank of an arbitrary matrix is the number of non-zero singular values (also number of linearly independent rows or columns)

Adam Prügel-Bennett AICE3001/6002 Mathematics of Machine Learning <https://ecs-vlc.github.io/aice3001/>

## Economical Forms of SVD

- Often the rows or columns of the orthogonal matrices  $\mathbf{U}$  and  $\mathbf{V}$  that are not associated with a singular value are ignored

$$\begin{pmatrix} \mathbf{X} \\ \mathbf{X}^T \end{pmatrix} = \begin{pmatrix} \mathbf{U} \\ \mathbf{V} \end{pmatrix} \begin{pmatrix} \mathbf{S} \\ \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{V}^T \\ \mathbf{U}^T \end{pmatrix}$$

- In Matlab these are obtained using

```
>> [U, S, V] = svd(X)
>> [U, S, V] = svd(X, 'econ')
```

Adam Prügel-Bennett AICE3001/6002 Mathematics of Machine Learning <https://ecs-vlc.github.io/aice3001/>

## General Matrix

- Recall that we can compute the SVD for any matrix,  $\mathbf{X}$
- As matrices describe the most general linear mapping

$$\mathbf{v} \rightarrow \mathcal{T}[\mathbf{v}] = \mathbf{X}\mathbf{v}$$

- We can use SVD to understand any linear mapping
- Thus any linear mapping can be seen as a rotation followed by a squashing or expansion independently in each coordinate followed by another rotation

Adam Prügel-Bennett AICE3001/6002 Mathematics of Machine Learning <https://ecs-vlc.github.io/aice3001/>

## Determinants

- The determinant,  $|\mathbf{M}|$  of a matrix  $\mathbf{M}$  is defined for square matrices
- It describes the change in volume under the mapping
- Now for any two matrices  $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$
- Thus  $|\mathbf{M}| = |\mathbf{U}||\mathbf{S}||\mathbf{V}^T|$
- For an orthogonal matrix  $|\mathbf{U}| = \pm 1$  since  $\mathbf{U}\mathbf{U}^T = \mathbf{I} \Rightarrow |\mathbf{U}\mathbf{U}^T| = |\mathbf{I}| \Rightarrow |\mathbf{U}||\mathbf{U}^T| = 1$  or  $|\mathbf{U}|^2 = 1$
- Thus

$$|\mathbf{M}| = \pm |\mathbf{S}| = \pm \prod_i s_i$$

Adam Prügel-Bennett AICE3001/6002 Mathematics of Machine Learning <https://ecs-vlc.github.io/aice3001/>

## Duality Revisited

- If  $\mathbf{X} = \mathbf{U}\mathbf{S}\mathbf{V}^T$  then

$$\begin{aligned} \mathbf{C} &= \mathbf{X}\mathbf{X}^T & \mathbf{D} &= \mathbf{X}^T\mathbf{X} \\ &= \mathbf{U}\mathbf{S}\mathbf{V}^T\mathbf{V}\mathbf{S}^T\mathbf{U}^T & &= \mathbf{V}\mathbf{S}^T\mathbf{U}^T\mathbf{U}\mathbf{S}\mathbf{V}^T \\ &= \mathbf{U}(\mathbf{S}\mathbf{S}^T)\mathbf{U}^T & &= \mathbf{V}(\mathbf{S}^T\mathbf{S})\mathbf{V}^T \end{aligned}$$

- If  $\mathbf{X}$  is an  $p \times m$  matrix then  $\mathbf{S}\mathbf{S}^T$  is a  $p \times p$  diagonal matrix with elements  $S_{ii}^2 = s_i^2$
- $\mathbf{S}^T\mathbf{S}$  is an  $m \times m$  matrix with elements  $S_{ii}^2 = s_i^2$
- $\mathbf{U}$  and  $\mathbf{V}$  are matrices of eigenvectors for  $\mathbf{C}$  and  $\mathbf{D}$
- The eigenvalues are  $\lambda_i = S_{ii}^2 = s_i^2$

Adam Prügel-Bennett AICE3001/6002 Mathematics of Machine Learning <https://ecs-vlc.github.io/aice3001/>

## SS<sup>T</sup> and S<sup>T</sup>S

$$S = \begin{pmatrix} s_1 & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & s_2 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & s_m & 0 & 0 \cdots & 0 \end{pmatrix}$$

$$S^T S = \begin{pmatrix} s_1^2 & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & s_2^2 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & s_m^2 & 0 & 0 \cdots & 0 \\ 0 & 0 & \cdots & 0 & 0 & 0 \cdots & 0 \\ \vdots & \vdots & \cdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & 0 & 0 \cdots & 0 \end{pmatrix}$$

$$S S^T = \begin{pmatrix} s_1^2 & 0 & \cdots & 0 \\ 0 & s_2^2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & s_m^2 \end{pmatrix}$$

## Outline

1. Singular Value Decomposition
2. General Linear Mappings
3. **Linear Regression Revisited**

$$\begin{pmatrix} 0 & X \\ X^T & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \sigma \begin{pmatrix} u \\ v \end{pmatrix}$$

## Having A Go

- It's really easy to verify this in MATLAB or OCTAVE

```
>> X = rand(3,2)
>> [U, S, V] = svd(X)
>> U*S*V'
>> U(:,1)'*U(:,2)
>> U'*U
>> U*U'
>> [Ua,L] = eig(X*X')
>> S*S'
```

- Test yourself!

## Linear Regression

- Given a set of data  $\mathcal{D} = \{(x_i, y_i) | k = 1, 2, \dots, m\}$
- In linear regression we try to fit a linear model

$$f(x|w) = x^T w$$

- Which we fit by minimising the squared error loss

$$L(w) = \sum_{k=1}^m (f(x_i|w) - y_i)^2$$

## Matrix Form

- In matrix form we write  $L(w) = \|Xw - y\|^2$

$$X = \begin{pmatrix} x_1^T \\ x_2^T \\ \vdots \\ x_m^T \end{pmatrix} \quad y = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{pmatrix}$$

- Then  $\nabla L(w^*) = 0$  implies

$$w^* = (X^T X)^{-1} X^T y = X^+ y$$

- This is known as the pseudo-inverse

## Pseudo-Inverse of S

$$S^T S = \begin{pmatrix} s_1^2 & 0 & \cdots & 0 \\ 0 & s_2^2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & s_p^2 \end{pmatrix} \quad (S^T S)^{-1} = \begin{pmatrix} s_1^{-2} & 0 & \cdots & 0 \\ 0 & s_2^{-2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & s_p^{-2} \end{pmatrix}$$

$$S^+ = (S^T S)^{-1} S^T = \begin{pmatrix} s_1^{-1} & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & s_2^{-1} & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & 0 & s_3^{-1} & \cdots & 0 & 0 \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & s_p^{-1} & 0 & 0 \cdots & 0 \end{pmatrix}$$

## Using SVD

- Using  $X = U S V^T$  then

$$\begin{aligned} X^+ &= (X^T X)^{-1} X^T \\ &= (V S^T S V^T)^{-1} V S^T U^T \\ &= V (S^T S)^{-1} V^T V S^T U^T \\ &= V (S^T S)^{-1} S^T U^T = V S^+ U^T \end{aligned}$$

- If  $m > p$

$$X^T = \begin{pmatrix} | & | & | & | & | & | & | & | \end{pmatrix}, S^T = \begin{pmatrix} s_1 & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & s_2 & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & 0 & s_3 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & s_p & 0 & 0 \cdots & 0 \end{pmatrix}$$

## III-Conditioned Data Matrix

- Recall that

$$w^* = X^+ y = V S^+ U^T y$$

- If any of the singular values of X are small then  $S^+$  will magnify components in that direction
- Any errors in the target  $y$  will be magnified
- This leads to poor weights

## Regularisation

- Consider linear regression with a regulariser

$$\begin{aligned}\mathcal{L}(w) &= \|Xw - y\|^2 + \eta \|w\|^2 \\ &= w^T (X^T X + \eta I) w - 2w^T X^T y + y^T y\end{aligned}$$

- Thus


$$\nabla \mathcal{L}(w) = 2(X^T X + \eta I)w - 2X^T y$$

- and  $\nabla \mathcal{L}(w^*) = 0$  gives

$$w^* = (X^T X + \eta I)^{-1} X^T y$$

## Effect of Regularisation

- Without regularisation if  $s_i = 0$  the problem would be ill-posed (even  $S^+$  does not exist since  $s_i^{-1}$  would be ill defined) and if  $s_i$  is small then  $S^+$  is ill conditioned
- Using  $\hat{S}^+ = (S^T S + \eta I)^{-1} S^T$  instead of  $S^+$  then



- Regularisation makes the machine much more stable (reduces the variance)

## Regularisation Continued

- Using  $X = U S V^T$

$$\begin{aligned}w^* &= (X^T X + \eta I)^{-1} X^T y \\ &= V (S^T S + \eta I)^{-1} S^T U^T y\end{aligned}$$

- where

$$(S^T S + \eta I)^{-1} S^T = \begin{pmatrix} \frac{s_1}{s_1^2 + \eta} & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & \frac{s_2}{s_2^2 + \eta} & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & 0 & \frac{s_3}{s_3^2 + \eta} & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \frac{s_p}{s_p^2 + \eta} & 0 & 0 & \cdots & 0 \end{pmatrix}$$

## Summary

- Any matrix can be decomposed as  $X = U S V^T$  where
  - $U$  and  $V$  are orthogonal (rotation matrices)
  - $S = \text{diag}(s_1, \dots, s_n)$  is a diagonal matrix of positive singular values
- This describes the most general linear transform
- The transform exploits the duality between  $XX^T$  and  $X^T X$
- In linear regression the pseudo-inverse involves the reciprocal of the singular values, which can lead to poor generalisation
- Regularisation improves the conditioning of the “inverse” matrix