


Mathematics of Machine Learning


Introduction



Course Details and Topics

Outline

1. **Why are you doing this course?**
2. What we assume that you know



Orientation

- This courses addresses three cohorts of students:
 - ★ Third year Artificial Intelligence Students
 - ★ MSc in Machine Learning Students
 - ★ Volunteers
- It is dual coded (AICE3001/AICE6002), with subtle difference in expectations reflected in slightly different evaluation
- Notes are on Moodle and

<https://ecs-vlc.github.io/aice3001/>

Orientation

- This courses addresses three cohorts of students:
 - ★ Third year Artificial Intelligence Students
 - ★ MSc in Machine Learning Students
 - ★ Volunteers
- It is dual coded (AICE3001/AICE6002), with subtle difference in expectations reflected in slightly different evaluation
- Notes are on Moodle and
<https://ecs-vlc.github.io/aice3001/>

Orientation

- This courses addresses three cohorts of students:
 - ★ Third year Artificial Intelligence Students
 - ★ MSc in Machine Learning Students
 - ★ Volunteers
- It is dual coded (AICE3001/AICE6002), with subtle difference in expectations reflected in slightly different evaluation
- Notes are on Moodle and
<https://ecs-vlc.github.io/aice3001/>

Why are We Teaching You This?

- In a world where LLMs can do everything, why bother with learning mathematics?

Why are We Teaching You This?

- In a world where LLMs can do everything, why bother with learning mathematics?
- You need to learn to trust your own judgement

Why are We Teaching You This?

- In a world where LLMs can do everything, why bother with learning mathematics?
- You need to learn to trust your own judgement
- LLMs can help us a lot, but if you don't trust yourself you are disempowered

Why are We Teaching You This?

- In a world where LLMs can do everything, why bother with learning mathematics?
- You need to learn to trust your own judgement
- LLMs can help us a lot, but if you don't trust yourself you are disempowered
- This course is aimed at building up trust in your understanding of how machine learning works

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition—we can't cheat
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition—we can't cheat **that much**
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition—we can't cheat that much
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Why Mathematics?

- Learning machines are notoriously difficult to understand
- Mathematics allows us to reason rigorously
- Posing question mathematically and seeking mathematical solutions provide allow us to build a strong intuition—we can't cheat that much
- Mathematics also allow us to describe the world precisely
- It also provides an arsenal of tools that turn out to be hugely useful

Cracking the Code

- Mathematics is a language that takes time to master
- For some it is easy, for others it will be a struggle
- But given determination everyone can crack the code and understand what the mathematics is saying
- This comes from engaging and working with maths

Cracking the Code

- Mathematics is a language that takes time to master
- For some it is easy, for others it will be a struggle
- But given determination everyone can crack the code and understand what the mathematics is saying
- This comes from engaging and working with maths

Cracking the Code

- Mathematics is a language that takes time to master
- For some it is easy, for others it will be a struggle
- But given determination everyone can crack the code and understand what the mathematics is saying
- This comes from engaging and working with maths

Cracking the Code

- Mathematics is a language that takes time to master
- For some it is easy, for others it will be a struggle
- But given determination everyone can crack the code and understand what the mathematics is saying
- This comes from engaging and working with maths

Cracking the Code

- Mathematics is a language that takes time to master
- For some it is easy, for others it will be a struggle
- But given determination everyone can crack the code and understand what the mathematics is saying
- This comes from engaging and working with maths—this course has tutorials that should be considered compulsory

Cracking the Code

- Mathematics is a language that takes time to master
- For some it is easy, for others it will be a struggle
- But given determination everyone can crack the code and understand what the mathematics is saying
- This comes from engaging and working with maths—this course has tutorials that should be considered compulsory **not because there are marks attached, but because without doing the work you aren't going to crack the code**

Starting from Different Places

- You all start from different places
- Some of you may have a degree in mathematics
- For others it may have been your least favourite subject
- I invite strong mathematicians to become even stronger
- For students starting from a weak base, judge yourselves by your progress, not by comparing yourself to others

Starting from Different Places

- You all start from different places
- Some of you may have a degree in mathematics
- For others it may have been your least favourite subject
- I invite strong mathematicians to become even stronger
- For students starting from a weak base, judge yourselves by your progress, not by comparing yourself to others

Starting from Different Places

- You all start from different places
- Some of you may have a degree in mathematics
- For others it may have been your least favourite subject
- I invite strong mathematicians to become even stronger
- For students starting from a weak base, judge yourselves by your progress, not by comparing yourself to others

Starting from Different Places

- You all start from different places
- Some of you may have a degree in mathematics
- For others it may have been your least favourite subject
- I invite strong mathematicians to become even stronger
- For students starting from a weak base, judge yourselves by your progress, not by comparing yourself to others

Starting from Different Places

- You all start from different places
- Some of you may have a degree in mathematics
- For others it may have been your least favourite subject
- I invite strong mathematicians to become even stronger—the best way to learn is through teaching others
- For students starting from a weak base, judge yourselves by your progress, not by comparing yourself to others

Starting from Different Places

- You all start from different places
- Some of you may have a degree in mathematics
- For others it may have been your least favourite subject
- I invite strong mathematicians to become even stronger—the best way to learn is through teaching others
- For students starting from a weak base, judge yourselves by your progress, not by comparing yourself to others

The Golden Rule of Mathematics

- Understand every step
- When you don't understand a step, stop
- Mathematics uses a lot of short-cuts, which speeds-up calculations

The Golden Rule of Mathematics

- Understand every step
- When you don't understand a step, stop
- Mathematics uses a lot of short-cuts, which speeds-up calculations

The Golden Rule of Mathematics

- Understand every step
- When you don't understand a step, stop, break the problem down
- Mathematics uses a lot of short-cuts, which speeds-up calculations

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, **ask someone**
- Mathematics uses a lot of short-cuts, which speeds-up calculations

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, ask someone
- **Mathematics uses a lot of short-cuts**, which speeds-up calculations

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, ask someone
- Mathematics uses a lot of short-cuts, **which speeds-up calculations**

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, ask someone
- Mathematics uses a lot of short-cuts, which speeds-up calculations, but you need to be able to justify every step

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, ask someone
- Mathematics uses a lot of short-cuts, which speeds-up calculations, but you need to be able to justify every step
- It takes bravery to admit you don't understand and insist that you will

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, ask someone
- Mathematics uses a lot of short-cuts, which speeds-up calculations, but you need to be able to justify every step
- It takes bravery to admit you don't understand and insist that you will
- Asking “stupid” questions isn't stupid

The Golden Rule of Mathematics

- **Understand every step**
- When you don't understand a step, stop, break the problem down, ask someone
- Mathematics uses a lot of short-cuts, which speeds-up calculations, but you need to be able to justify every step
- It takes bravery to admit you don't understand and insist that you will
- Asking “stupid” questions isn't stupid **it is how we all learn**

The Journey

- We are going to visit lots of aspects of mathematics that have proved fruitful in machine learning
- Some of this will be very concrete, while in other cases by look at a higher level of abstraction we can generalise methods across different application areas
- We will cover, calculus, vector spaces, convexity, optimisation, entropy, probability, statistics and much else besides
- The journey will pay you off, these are the foundations of most of machine learning and will give you super-powers that are hard to acquire without mathematics

The Journey

- We are going to visit lots of aspects of mathematics that have proved fruitful in machine learning
- Some of this will be very concrete, while in other cases by look at a higher level of abstraction we can generalise methods across different application areas
- We will cover, calculus, vector spaces, convexity, optimisation, entropy, probability, statistics and much else besides
- The journey will pay you off, these are the foundations of most of machine learning and will give you super-powers that are hard to acquire without mathematics

The Journey

- We are going to visit lots of aspects of mathematics that have proved fruitful in machine learning
- Some of this will be very concrete, while in other cases by look at a higher level of abstraction we can generalise methods across different application areas
- We will cover, calculus, vector spaces, convexity, optimisation, entropy, probability, statistics and much else besides
- The journey will pay you off, these are the foundations of most of machine learning and will give you super-powers that are hard to acquire without mathematics

The Journey


- We are going to visit lots of aspects of mathematics that have proved fruitful in machine learning
- Some of this will be very concrete, while in other cases by look at a higher level of abstraction we can generalise methods across different application areas
- We will cover, calculus, vector spaces, convexity, optimisation, entropy, probability, statistics and much else besides
- **The journey will pay you off**, these are the foundations of most of machine learning and will give you super-powers that are hard to acquire without mathematics

The Journey

- We are going to visit lots of aspects of mathematics that have proved fruitful in machine learning
- Some of this will be very concrete, while in other cases by look at a higher level of abstraction we can generalise methods across different application areas
- We will cover, calculus, vector spaces, convexity, optimisation, entropy, probability, statistics and much else besides
- The journey will pay you off, these are the foundations of most of machine learning and will give you super-powers that are hard to acquire without mathematics

Outline

1. Why are you doing this course?
2. **What we assume that you know**



What we Assume you Know

- You should all have A-level mathematics and have done some mathematics at Uni
- You should know what a function is, e.g. $f(x) = x^2 + 3$
- You should be able to do simple algebraic manipulations
- And some elementary calculus

$$f'(x) = \frac{d(x^2 + 3)}{dx} = 2x$$

What we Assume you Know

- You should all have A-level mathematics and have done some mathematics at Uni
- You should know what a function is, e.g. $f(x) = x^2 + 3$
- You should be able to do simple algebraic manipulations
- And some elementary calculus

$$f'(x) = \frac{d(x^2 + 3)}{dx} = 2x$$

What we Assume you Know

- You should all have A-level mathematics and have done some mathematics at Uni
- You should know what a function is, e.g. $f(x) = x^2 + 3$
- You should be able to do simple algebraic manipulations
- And some elementary calculus

$$f'(x) = \frac{d(x^2 + 3)}{dx} = 2x$$

What we Assume you Know

- You should all have A-level mathematics and have done some mathematics at Uni
- You should know what a function is, e.g. $f(x) = x^2 + 3$
- You should be able to do simple algebraic manipulations
- And some elementary calculus

$$f'(x) = \frac{d(x^2 + 3)}{dx} = 2x$$

Vectors

- Ordinary vectors (what mathematicians call elements of $\mathbb{R}^n$) are much used in machine learning and should be familiar
- We will use bold symbols to represent **column vectors**, e.g.

$$\mathbf{x} = \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix}$$

- We denote **row vectors** as the **transpose** of column vectors
 $\mathbf{x}^T = (1.2, 2.3, 4.7)$
- We care whether vectors are row or column vectors as it allows us to write equations involving vectors succinctly and unambiguously

Vectors

- Ordinary vectors (what mathematicians call elements of $\mathbb{R}^n$) are much used in machine learning and should be familiar
- We will use bold symbols to represent **column vectors**, e.g.

$$\mathbf{x} = \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix}$$

- We denote **row vectors** as the **transpose** of column vectors
 $\mathbf{x}^T = (1.2, 2.3, 4.7)$
- We care whether vectors are row or column vectors as it allows us to write equations involving vectors succinctly and unambiguously

Vectors

- Ordinary vectors (what mathematicians call elements of $\mathbb{R}^n$) are much used in machine learning and should be familiar
- We will use bold symbols to represent **column vectors**, e.g.

$$\mathbf{x} = \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix}$$

- We denote **row vectors** as the **transpose** of column vectors
 $\mathbf{x}^T = (1.2, 2.3, 4.7)$
- We care whether vectors are row or column vectors as it allows us to write equations involving vectors succinctly and unambiguously

Vectors

- Ordinary vectors (what mathematicians call elements of $\mathbb{R}^n$) are much used in machine learning and should be familiar
- We will use bold symbols to represent **column vectors**, e.g.

$$\mathbf{x} = \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix}$$

- We denote **row vectors** as the **transpose** of column vectors
 $\mathbf{x}^T = (1.2, 2.3, 4.7)$
- We care whether vectors are row or column vectors as it allows us to write equations involving vectors succinctly and unambiguously

Matrices

- Matrices are blocks of number that define linear mappings between vectors, e.g. $\mathbf{M} \in \mathbb{R}^{2 \times 3}$

$$\mathbf{M} = \begin{pmatrix} 1.1 & -2.1 & 6.2 \\ 9.3 & 3.1 & -4.4 \end{pmatrix}$$

- An $m \times n$ matrix (i.e. $\mathbf{M} \in \mathbb{R}^{m \times n}$) has m rows and n columns
- An $m \times n$ matrix acting on a vector in $\mathbb{R}^n$ transforms it to $\mathbb{R}^m$ vector, e.g.

$$\mathbf{M}\mathbf{x} = \begin{pmatrix} 1.1 & -2.1 & 6.2 \\ 9.3 & 3.1 & -4.4 \end{pmatrix} \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix} = \begin{pmatrix} 28.63 \\ -2.39 \end{pmatrix}$$

Matrices

- Matrices are blocks of number that define linear mappings between vectors, e.g. $\mathbf{M} \in \mathbb{R}^{2 \times 3}$

$$\mathbf{M} = \begin{pmatrix} 1.1 & -2.1 & 6.2 \\ 9.3 & 3.1 & -4.4 \end{pmatrix}$$

- An $m \times n$ matrix (i.e. $\mathbf{M} \in \mathbb{R}^{m \times n}$) has m rows and n columns
- An $m \times n$ matrix acting on a vector in $\mathbb{R}^n$ transforms it to $\mathbb{R}^m$ vector, e.g.

$$\mathbf{M}\mathbf{x} = \begin{pmatrix} 1.1 & -2.1 & 6.2 \\ 9.3 & 3.1 & -4.4 \end{pmatrix} \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix} = \begin{pmatrix} 28.63 \\ -2.39 \end{pmatrix}$$

Matrices

- Matrices are blocks of number that define linear mappings between vectors, e.g. $\mathbf{M} \in \mathbb{R}^{2 \times 3}$

$$\mathbf{M} = \begin{pmatrix} 1.1 & -2.1 & 6.2 \\ 9.3 & 3.1 & -4.4 \end{pmatrix}$$

- An $m \times n$ matrix (i.e. $\mathbf{M} \in \mathbb{R}^{m \times n}$) has m rows and n columns
- An $m \times n$ matrix acting on a vector in $\mathbb{R}^n$ transforms it to $\mathbb{R}^m$ vector, e.g.

$$\mathbf{M}\mathbf{x} = \begin{pmatrix} 1.1 & -2.1 & 6.2 \\ 9.3 & 3.1 & -4.4 \end{pmatrix} \begin{pmatrix} 1.2 \\ 2.3 \\ 4.7 \end{pmatrix} = \begin{pmatrix} 28.63 \\ -2.39 \end{pmatrix}$$

Composition of Mappings

- Matrices, $\mathbf{A}$, $\mathbf{B}$, etc. represent mappings between spaces
- Provided the matrices are the correct size we can compose these mappings, e.g. if $\mathbf{A} \in \mathbb{R}^{\ell \times m}$ and $\mathbf{B} \in \mathbb{R}^{m \times n}$ then $\mathbf{AB}$ represents the effect of
 - ★ first applying $\mathbf{B}$ to vectors in $\mathbb{R}^n$ to obtain vectors in $\mathbb{R}^m$
 - ★ and then applying $\mathbf{A}$ to get vectors in $\mathbb{R}^\ell$

$$\mathbf{AB}x = \mathbf{A}(\mathbf{B}x)$$

Composition of Mappings

- Matrices, $\mathbf{A}$, $\mathbf{B}$, etc. represent mappings between spaces
- Provided the matrices are the correct size we can compose these mappings, e.g. if $\mathbf{A} \in \mathbb{R}^{\ell \times m}$ and $\mathbf{B} \in \mathbb{R}^{m \times n}$ then $\mathbf{AB}$ represents the effect of
 - ★ first applying $\mathbf{B}$ to vectors in $\mathbb{R}^n$ to obtain vectors in $\mathbb{R}^m$
 - ★ and then applying $\mathbf{A}$ to get vectors in $\mathbb{R}^\ell$

$$\mathbf{AB}x = \mathbf{A}(\mathbf{B}x)$$

Composition of Mappings

- Matrices, $\mathbf{A}$, $\mathbf{B}$, etc. represent mappings between spaces
- Provided the matrices are the correct size we can compose these mappings, e.g. if $\mathbf{A} \in \mathbb{R}^{\ell \times m}$ and $\mathbf{B} \in \mathbb{R}^{m \times n}$ then $\mathbf{AB}$ represents the effect of
 - ★ first applying $\mathbf{B}$ to vectors in $\mathbb{R}^n$ to obtain vectors in $\mathbb{R}^m$
 - ★ and then applying $\mathbf{A}$ to get vectors in $\mathbb{R}^\ell$

$$\mathbf{AB}x = \mathbf{A}(\mathbf{B}x)$$

Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)


Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

The diagram illustrates matrix multiplication using a visual representation of matrices as grids of squares. On the left, a 5x5 matrix (A) is shown with 5 rows and 5 columns of squares. This is followed by a 5x3 matrix (B) with 5 rows and 3 columns of squares. An equals sign follows, leading to a 5x3 matrix (C) with 5 rows and 3 columns of squares. The visual representation shows that the product of a 5x5 matrix and a 5x3 matrix is a 5x3 matrix.


Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)




Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

$$\begin{pmatrix} \text{red} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \end{pmatrix} \begin{pmatrix} \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{red} & \text{gray} \end{pmatrix} = \begin{pmatrix} \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \end{pmatrix}$$


Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

The diagram illustrates matrix multiplication using a grid of squares. The first matrix is 4 rows by 5 columns. The second matrix is 5 rows by 3 columns. The result is a 4 rows by 3 columns matrix. The red squares indicate the specific row and column being multiplied to produce a single element in the result matrix.

Gray	Gray	Gray	Gray	Gray	=	Gray	Gray	Gray
Red	Red	Red	Red	Red		Gray	Red	Gray
Gray	Gray	Gray	Gray	Gray		Gray	Gray	Gray
Gray	Gray	Gray	Gray	Gray		Gray	Gray	Gray
Gray	Gray	Gray	Gray	Gray		Gray	Gray	Gray


Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

$$\begin{pmatrix} \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{red} & \text{red} & \text{red} & \text{red} & \text{red} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \end{pmatrix} \begin{pmatrix} \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{red} \end{pmatrix} = \begin{pmatrix} \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \end{pmatrix}$$

Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

The diagram illustrates the matrix multiplication $\mathbf{AB} = \mathbf{C}$ using a grid of squares. The first matrix (A) is 5 rows by 5 columns. The second matrix (B) is 5 rows by 3 columns. The result matrix (C) is 5 rows by 3 columns. The red squares indicate the non-zero elements in each matrix.

Matrix A (5x5)	Matrix B (5x3)	Result Matrix C (5x3)
Gray	Gray	Gray
Gray	Gray	Gray
Red	Red	Red
Gray	Gray	Gray
Gray	Gray	Gray


Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

The diagram illustrates matrix multiplication using a grid of squares. The first matrix is a 4x5 grid with 4 rows and 5 columns. The third row is highlighted in red, while all other squares are gray. This is followed by a 5x3 grid with 5 rows and 3 columns. The third column is highlighted in red, while all other squares are gray. An equals sign follows, leading to a 4x3 grid with 4 rows and 3 columns. The third column is highlighted in red, while all other squares are gray. This represents the result of the matrix multiplication.


Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)




Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)

The diagram illustrates matrix multiplication using a grid of squares. The first matrix is a 4x5 grid with 4 rows and 5 columns. The first three rows are all gray squares, and the fourth row consists of five red squares. The second matrix is a 5x3 grid with 5 rows and 3 columns. The first four rows are all gray squares, and the fifth row consists of three red squares. An equals sign follows, and the resulting matrix is a 4x3 grid with 4 rows and 3 columns. The first three rows are all gray squares, and the fourth row consists of two gray squares followed by one red square.

Matrix Multiplication


- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



- Matrix multiplication is associative: $(\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC})$

Matrix Multiplication

- The combined mapping $\mathbf{AB}$ can be represented by an $\ell \times n$ matrix $\mathbf{C}$ which we can compute using matrix multiplication (rows times columns)



- Matrix multiplication is associative: $(\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC})$
- But in general not commutative $\mathbf{AB} \neq \mathbf{BA}$

Transpose

- We often encounter transposes of a matrix $\mathbf{M}^T$ where we have exchanged the rows and columns: $[\mathbf{M}^T]_{ij} = [\mathbf{M}]_{ji}$
- Clear $(\mathbf{M}^T)^T = \mathbf{M}$
- It is simple question of book keeping to show that

$$(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$$

- For a symmetric matrix $\mathbf{M} = \mathbf{M}^T$
- Since 1×1 matrices are always symmetric

$$\mathbf{a}^T \mathbf{b} = (\mathbf{a}^T \mathbf{b})^T = \mathbf{b}^T \mathbf{a}$$

Transpose

- We often encounter transposes of a matrix $\mathbf{M}^T$ where we have exchanged the rows and columns: $[\mathbf{M}^T]_{ij} = [\mathbf{M}]_{ji}$
- Clear $(\mathbf{M}^T)^T = \mathbf{M}$
- It is simple question of book keeping to show that

$$(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$$

- For a symmetric matrix $\mathbf{M} = \mathbf{M}^T$
- Since 1×1 matrices are always symmetric

$$\mathbf{a}^T \mathbf{b} = (\mathbf{a}^T \mathbf{b})^T = \mathbf{b}^T \mathbf{a}$$

Transpose

- We often encounter transposes of a matrix $\mathbf{M}^T$ where we have exchanged the rows and columns: $[\mathbf{M}^T]_{ij} = [\mathbf{M}]_{ji}$
- Clear $(\mathbf{M}^T)^T = \mathbf{M}$
- It is simple question of book keeping to show that

$$(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$$

- For a symmetric matrix $\mathbf{M} = \mathbf{M}^T$
- Since 1×1 matrices are always symmetric

$$\mathbf{a}^T \mathbf{b} = (\mathbf{a}^T \mathbf{b})^T = \mathbf{b}^T \mathbf{a}$$

Transpose

- We often encounter transposes of a matrix $\mathbf{M}^\top$ where we have exchanged the rows and columns: $[\mathbf{M}^\top]_{ij} = [\mathbf{M}]_{ji}$
- Clear $(\mathbf{M}^\top)^\top = \mathbf{M}$
- It is simple question of book keeping to show that

$$(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$$

- For a symmetric matrix $\mathbf{M} = \mathbf{M}^\top$
- Since 1×1 matrices are always symmetric

$$\mathbf{a}^\top \mathbf{b} = (\mathbf{a}^\top \mathbf{b})^\top = \mathbf{b}^\top \mathbf{a}$$

Transpose

- We often encounter transposes of a matrix $\mathbf{M}^\top$ where we have exchanged the rows and columns: $[\mathbf{M}^\top]_{ij} = [\mathbf{M}]_{ji}$
- Clear $(\mathbf{M}^\top)^\top = \mathbf{M}$
- It is simple question of book keeping to show that

$$(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$$

- For a symmetric matrix $\mathbf{M} = \mathbf{M}^\top$
- Since 1×1 matrices are always symmetric

$$\mathbf{a}^\top \mathbf{b} = (\mathbf{a}^\top \mathbf{b})^\top = \mathbf{b}^\top \mathbf{a}$$

Transpose

- We often encounter transposes of a matrix $\mathbf{M}^\top$ where we have exchanged the rows and columns: $[\mathbf{M}^\top]_{ij} = [\mathbf{M}]_{ji}$
- Clear $(\mathbf{M}^\top)^\top = \mathbf{M}$
- It is simple question of book keeping to show that

$$(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$$

- For a symmetric matrix $\mathbf{M} = \mathbf{M}^\top$
- Since 1×1 matrices are always symmetric

$$\begin{aligned} \mathbf{a}^\top \mathbf{b} &= (\mathbf{a}^\top \mathbf{b})^\top = \mathbf{b}^\top \mathbf{a} \\ \mathbf{a}^\top \mathbf{M} \mathbf{b} &= \mathbf{b}^\top \mathbf{M}^\top \mathbf{a} \end{aligned}$$

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix
- Then matrix-vector multiplication can be treated the same as matrix multiplication

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix
- Then matrix-vector multiplication can be treated the same as matrix multiplication

The diagram illustrates matrix-vector multiplication as matrix multiplication. On the left, a 4x5 matrix is represented by a large left parenthesis followed by a 4x5 grid of small gray squares, followed by a large right parenthesis. This is multiplied by a 5x1 column vector, represented by a large left parenthesis followed by a vertical column of five small gray squares, followed by a large right parenthesis. An equals sign follows, and then a 4x1 column vector is shown, represented by a large left parenthesis followed by a vertical column of four small gray squares, followed by a large right parenthesis.

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix
- Then matrix-vector multiplication can be treated the same as matrix multiplication

The diagram illustrates matrix-vector multiplication as matrix multiplication. It shows a 4x5 matrix (with 5 red squares in the top row and 4 gray squares in each of the other three rows) multiplied by a 5x1 column vector (with 5 red squares). This is equated to a 4x1 column vector (with 1 red square in the top row and 3 gray squares in the other three rows).

$$\begin{pmatrix} \text{red} & \text{red} & \text{red} & \text{red} & \text{red} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \end{pmatrix} \begin{pmatrix} \text{red} \\ \text{red} \\ \text{red} \\ \text{red} \\ \text{red} \end{pmatrix} = \begin{pmatrix} \text{red} \\ \text{gray} \\ \text{gray} \\ \text{gray} \end{pmatrix}$$

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix
- Then matrix-vector multiplication can be treated the same as matrix multiplication

The diagram illustrates matrix-vector multiplication as matrix multiplication. It shows a 5x5 matrix (with 1 red square in the second row) multiplied by a 5x1 column vector (all red squares) equals a 5x1 column vector (with 1 red square in the second row).

$$\begin{pmatrix} \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{red} & \text{red} & \text{red} & \text{red} & \text{red} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} & \text{gray} & \text{gray} \end{pmatrix} \begin{pmatrix} \text{red} \\ \text{red} \\ \text{red} \\ \text{red} \\ \text{red} \end{pmatrix} = \begin{pmatrix} \text{gray} \\ \text{red} \\ \text{gray} \\ \text{gray} \\ \text{gray} \end{pmatrix}$$

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix
- Then matrix-vector multiplication can be treated the same as matrix multiplication

The diagram illustrates matrix-vector multiplication as matrix multiplication. On the left, a 5x5 matrix is shown with 4 gray squares and 1 red square in each row. This matrix is multiplied by a 5x1 column vector, which is represented as a single column containing 5 red squares. An equals sign follows, and on the right, the resulting 5x1 column vector is shown, containing 4 gray squares and 1 red square. This visualizes the process of multiplying each row of the matrix by the column vector and summing the results to form the new column vector.

Matrix-Vector Multiplication

- We can treat a column vector as a $n \times 1$ matrix
- Then matrix-vector multiplication can be treated the same as matrix multiplication

The diagram illustrates matrix-vector multiplication as matrix multiplication. It shows a 5x5 matrix (represented by a grid of 25 squares, with the bottom row highlighted in red) multiplied by a 5x1 column vector (represented by a single column of 5 red squares). The result is a 5x1 column vector (represented by a single column of 5 squares, with the bottom square highlighted in red). The entire equation is enclosed in large parentheses.


$$\begin{pmatrix} \text{5x5 grid of squares} \end{pmatrix} \begin{pmatrix} \text{5x1 column of red squares} \end{pmatrix} = \begin{pmatrix} \text{5x1 column of squares} \end{pmatrix}$$

Inner Products

- The dot or inner-product between two vectors with the same number of elements, can be written $a^T b$


Inner Products

- The dot or inner-product between two vectors with the same number of elements, can be written $a^T b$


$$\left(\begin{array}{ccccc} \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare \end{array} \right) \left(\begin{array}{c} \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \end{array} \right) = \left(\blacksquare \right)$$


Inner Products

- The dot or inner-product between two vectors with the same number of elements, can be written $\mathbf{a}^T \mathbf{b}$




Inner Products

- The dot or inner-product between two vectors with the same number of elements, can be written $a^T b$


$$\left(\begin{array}{ccccc} \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare \end{array} \right) = \left(\begin{array}{c} \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \end{array} \right) = \left(\blacksquare \right)$$

Inner Products

- The dot or inner-product between two vectors with the same number of elements, can be written $\mathbf{a}^T \mathbf{b}$


$$\left(\begin{array}{ccccc} \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare \end{array} \right) \left(\begin{array}{c} \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \end{array} \right) = \left(\blacksquare \right)$$

- Inner products encode angles between vectors

$$\mathbf{a}^T \mathbf{b} = \|\mathbf{a}\| \|\mathbf{b}\| \cos(\theta)$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product operation ab^T . On the left, a column vector a of size 4x1 is represented by a large left parenthesis followed by four gray squares stacked vertically, and a large right parenthesis. This is multiplied by a row vector b^T of size 1x3, represented by a large left parenthesis followed by three gray squares, and a large right parenthesis. An equals sign follows, leading to the resulting 4x3 matrix. This matrix is shown as a large left parenthesis followed by four rows, each containing three gray squares, and a large right parenthesis.


Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product operation ab^T using square blocks to represent matrix elements. On the left, a column vector a of size 4x1 is shown, with its top element in red and the others in gray. This is multiplied by a row vector b^T of size 1x3, also with its first element in red and the others in gray. An equals sign follows, leading to a 4x3 matrix on the right. Each row of this matrix corresponds to one of the four elements of a . The first row contains one red square followed by two gray squares, representing the first row of a multiplied by the entire row vector b^T . The subsequent three rows each contain three gray squares, representing the other rows of a multiplied by b^T .

Outer Products

- We can also define the outer-product between any size vectors, ab^T



Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4x1 is shown, with its top element highlighted in red. This is multiplied by a row vector of size 1x3, also with its middle element highlighted in red. The result is a 4x3 matrix where every element in the first column is red, representing the scaling of the row vector by the scalar value of the first element of the column vector.

$$\begin{pmatrix} \text{red} \\ \text{gray} \\ \text{gray} \\ \text{gray} \end{pmatrix} \begin{pmatrix} \text{gray} & \text{red} & \text{gray} \end{pmatrix} = \begin{pmatrix} \text{red} & \text{red} & \text{red} \\ \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{red} & \text{gray} \\ \text{gray} & \text{red} & \text{gray} \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector is shown with four elements: a red square at the top, followed by three gray squares. This is multiplied by a row vector with three elements: two gray squares followed by a red square. The result is a 4x3 matrix on the right, where each row is a scaled version of the row vector. The first row has a red square in the third column, while the other three rows have gray squares in all three columns.

$$\begin{pmatrix} \text{red} \\ \text{gray} \\ \text{gray} \\ \text{gray} \end{pmatrix} \begin{pmatrix} \text{gray} & \text{gray} & \text{red} \end{pmatrix} = \begin{pmatrix} \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product operation ab^T . On the left, a column vector a of size 4x1 is shown, with its second element highlighted in red. This is multiplied by a row vector b^T of size 1x3, with its first element highlighted in red. The result is a 4x3 matrix where each row is a scalar multiple of the vector b^T , and the first column is a scalar multiple of the vector a . The red element in the first row, first column of the resulting matrix corresponds to the product of the red elements in the input vectors.

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product operation ab^T using square blocks to represent elements of vectors and matrices. On the left, a column vector a is shown as a vertical stack of four squares, with the second square from the top highlighted in red. To its right is a row vector b^T shown as a horizontal row of three squares, with the middle square highlighted in red. An equals sign follows, leading to the resulting 4x3 matrix. This matrix is represented by a large set of parentheses containing a 4x3 grid of squares. The second row and third column of this grid are highlighted in red, showing that every element in the resulting matrix is the product of the red element from the first vector and the red element from the second vector.

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4 is shown, with its second element highlighted in red. This is multiplied by a row vector of size 3, with its third element highlighted in red. The result is a 4x3 matrix where the third column is red and all other elements are gray, representing the outer product of the original vectors.

$$\begin{pmatrix} \text{gray} \\ \text{red} \\ \text{gray} \\ \text{gray} \end{pmatrix} \begin{pmatrix} \text{gray} & \text{gray} & \text{red} \end{pmatrix} = \begin{pmatrix} \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{red} \\ \text{gray} & \text{gray} & \text{gray} \\ \text{gray} & \text{gray} & \text{gray} \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product operation ab^T . On the left, a column vector a of size 4x1 is shown, with its third element highlighted in red. This is multiplied by a row vector b^T of size 1x3, with its first element highlighted in red. The result is a 4x3 matrix where each row is a scalar multiple of the vector b^T , and the third row is highlighted in red because it is the product of the red element in a and the red element in b^T .

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4 is shown, with its third element highlighted in red. This is multiplied by a row vector of size 3, also with its third element highlighted in red. The result is a 4x3 matrix where the third row and third column are highlighted in red, representing the outer product of the original vectors.

$$\begin{pmatrix} \square \\ \square \\ \color{red}\square \\ \square \end{pmatrix} \begin{pmatrix} \square & \color{red}\square & \square \end{pmatrix} = \begin{pmatrix} \square & \square & \square \\ \square & \square & \square \\ \color{red}\square & \color{red}\square & \color{red}\square \\ \square & \square & \square \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector is represented by a large left parenthesis followed by four gray squares, with the third square highlighted in red, and a large right parenthesis. This is followed by a row vector represented by a large left parenthesis followed by three squares (two gray, one red), and a large right parenthesis. An equals sign follows, leading to a 4x3 matrix. The matrix is enclosed in large parentheses and contains four rows and three columns of squares. The third row and third column are highlighted in red, representing the outer product of the red elements from the original vectors.

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4 is shown, with the bottom element highlighted in red. This is multiplied by a row vector of size 3, with the first element highlighted in red. The result is a 4x3 matrix on the right, where each row is a scaled version of the row vector, and the first column is a scaled version of the column vector. The red squares indicate the contribution of the highlighted elements in the input vectors to the resulting matrix.

$$\begin{pmatrix} \square \\ \square \\ \square \\ \color{red}\square \end{pmatrix} \begin{pmatrix} \color{red}\square & \square & \square \end{pmatrix} = \begin{pmatrix} \square & \square & \square \\ \square & \square & \square \\ \square & \square & \square \\ \color{red}\square & \square & \square \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector a of size 4x1 is shown, with its bottom element highlighted in red. This is multiplied by a row vector b^T of size 1x3, with its middle element highlighted in red. The result is a 4x3 matrix on the right, where the bottom-middle element is red, indicating it is the product of the red elements from the input vectors. All other elements in the matrix are gray.

$$\begin{pmatrix} \square \\ \square \\ \square \\ \color{red}\square \end{pmatrix} \begin{pmatrix} \square & \color{red}\square & \square \end{pmatrix} = \begin{pmatrix} \square & \square & \square \\ \square & \square & \square \\ \square & \square & \square \\ \square & \color{red}\square & \square \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of two vectors. On the left, a column vector of size 4 is shown, consisting of three gray squares and one red square at the bottom. This is multiplied by a row vector of size 3, consisting of two gray squares and one red square on the right. The result is a 4x3 matrix on the right, where each element is the product of the corresponding elements from the two input vectors. The bottom-right element is red, while all other elements are gray.

$$\begin{pmatrix} \square \\ \square \\ \square \\ \color{red}\square \end{pmatrix} \begin{pmatrix} \square & \square & \color{red}\square \end{pmatrix} = \begin{pmatrix} \square & \square & \square \\ \square & \square & \square \\ \square & \square & \square \\ \square & \square & \color{red}\square \end{pmatrix}$$

Outer Products

- We can also define the outer-product between any size vectors, ab^T

The diagram illustrates the outer product of a column vector and a row vector. On the left, a column vector is shown with four elements: three gray squares and one red square at the bottom. This is multiplied by a row vector with three elements: two gray squares and one red square on the right. The result is a 4x3 matrix on the right, where each element is the product of the corresponding elements from the input vectors. The red square in the bottom-right corner of the matrix indicates the product of the red squares from the input vectors.

- Notice that we need to know whether we are dealing with row or column vectors

Norms

- Norms are non-negative numbers associated with the “size” of a vector
- The norm you should be familiar with is the **Euclidean Norm** or 2-norm

$$\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- We will see there exist other norms (using different ways of measuring size)
- The Euclidean norm is intimately related to the inner-product, we will see other examples of this for different vector spaces

Norms

- Norms are non-negative numbers associated with the “size” of a vector
- The norm you should be familiar with is the **Euclidean Norm** or 2-norm

$$\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- We will see there exist other norms (using different ways of measuring size)
- The Euclidean norm is intimately related to the inner-product, we will see other examples of this for different vector spaces

Norms

- Norms are non-negative numbers associated with the “size” of a vector
- The norm you should be familiar with is the **Euclidean Norm** or 2-norm

$$\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2} = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}$$

- We will see there exist other norms (using different ways of measuring size)
- The Euclidean norm is intimately related to the inner-product, we will see other examples of this for different vector spaces

Norms

- Norms are non-negative numbers associated with the “size” of a vector
- The norm you should be familiar with is the **Euclidean Norm** or 2-norm

$$\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2} = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} = \sqrt{\mathbf{x}^\top \mathbf{x}}$$

- We will see there exist other norms (using different ways of measuring size)
- The Euclidean norm is intimately related to the inner-product, we will see other examples of this for different vector spaces

Norms

- Norms are non-negative numbers associated with the “size” of a vector
- The norm you should be familiar with is the **Euclidean Norm** or 2-norm

$$\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2} = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} = \sqrt{\mathbf{x}^\top \mathbf{x}}$$

- We will see there exist other norms (using different ways of measuring size)
- The Euclidean norm is intimately related to the inner-product, we will see other examples of this for different vector spaces

Norms

- Norms are non-negative numbers associated with the “size” of a vector
- The norm you should be familiar with is the **Euclidean Norm** or 2-norm

$$\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2} = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} = \sqrt{\mathbf{x}^\top \mathbf{x}}$$

- We will see there exist other norms (using different ways of measuring size)
- The Euclidean norm is intimately related to the inner-product, we will see other examples of this for different vector spaces

Matrices as Linear Mappings

- Matrices represent mappings between vectors

$$\mathbf{b} = \mathbf{M}\mathbf{a}$$

- The elements of $\mathbf{b}$ are **linear sums** of the elements of $\mathbf{a}$

$$b_i = \sum_j M_{ij} a_j$$

- Two important consequences of this linearity are for any real c

$$\mathbf{M}(c\mathbf{a}) = c(\mathbf{M}\mathbf{a})$$

Matrices as Linear Mappings

- Matrices represent mappings between vectors

$$\mathbf{b} = \mathbf{M}\mathbf{a}$$

- The elements of $\mathbf{b}$ are **linear sums** of the elements of $\mathbf{a}$

$$b_i = \sum_j M_{ij}a_j = M_{i1}a_1 + M_{i2}a_2 + \cdots + M_{in}a_n$$

- Two important consequences of this linearity are for any real c

$$\mathbf{M}(c\mathbf{a}) = c(\mathbf{M}\mathbf{a})$$

Matrices as Linear Mappings

- Matrices represent mappings between vectors

$$\mathbf{b} = \mathbf{M}\mathbf{a}$$

- The elements of $\mathbf{b}$ are **linear sums** of the elements of $\mathbf{a}$

$$b_i = \sum_j M_{ij}a_j = M_{i1}a_1 + M_{i2}a_2 + \cdots + M_{in}a_n$$

- Two important consequences of this linearity are for any real c

$$\mathbf{M}(c\mathbf{a}) = c(\mathbf{M}\mathbf{a})$$

Matrices as Linear Mappings

- Matrices represent mappings between vectors

$$\mathbf{b} = \mathbf{M}\mathbf{a}$$

- The elements of $\mathbf{b}$ are **linear sums** of the elements of $\mathbf{a}$

$$b_i = \sum_j M_{ij}a_j = M_{i1}a_1 + M_{i2}a_2 + \cdots + M_{in}a_n$$


- Two important consequences of this linearity are for any real c

$$\mathbf{M}(c\mathbf{a}) = c(\mathbf{M}\mathbf{a})$$

$$\mathbf{M}(\mathbf{a} + \mathbf{b}) = (\mathbf{M}\mathbf{a}) + (\mathbf{M}\mathbf{b})$$

Action Of Matrices on Volumes


- For a closed set of points corresponding to a volume, V in $\mathbb{R}^n$, treating each point as a vector, we can ask what happens to the volume after mapping the vector by a matrix



- For square matrices, $\mathbf{M}$, the determinant is equal to $\det(\mathbf{M}) = \pm V'/V$
- The sign tells us if there has been a reflection

Action Of Matrices on Volumes


- For a closed set of points corresponding to a volume, V in $\mathbb{R}^n$, treating each point as a vector, we can ask what happens to the volume after mapping the vector by a matrix



- For square matrices, $\mathbf{M}$, the determinant is equal to $\det(\mathbf{M}) = \pm V'/V$
- The sign tells us if there has been a reflection

Action Of Matrices on Volumes

- For a closed set of points corresponding to a volume, V in $\mathbb{R}^n$, treating each point as a vector, we can ask what happens to the volume after mapping the vector by a matrix



- For square matrices, $\mathbf{M}$, the determinant is equal to $\det(\mathbf{M}) = \pm V'/V$
- The sign tells us if there has been a reflection

Determinants

- A consequence of the fact that $\mathbf{AB}$ denotes the composition, apply mapping $\mathbf{B}$ then apply mapping $\mathbf{A}$

$$\det(\mathbf{AB}) = \det(\mathbf{A}) \times \det(\mathbf{B})$$

- Determinants are only defined for square matrices, as non-square matrices project into different size spaces, where the volumes cannot be compared
- **Singular matrices** projects all points onto a lower dimensional space, so have determinant 0

Determinants

- A consequence of the fact that $\mathbf{AB}$ denotes the composition, apply mapping $\mathbf{B}$ then apply mapping $\mathbf{A}$

$$\det(\mathbf{AB}) = \det(\mathbf{A}) \times \det(\mathbf{B})$$

- Determinants are only defined for square matrices, as non-square matrices project into different size spaces, where the volumes cannot be compared
- **Singular matrices** projects all points onto a lower dimensional space, so have determinant 0

Determinants

- A consequence of the fact that $\mathbf{AB}$ denotes the composition, apply mapping $\mathbf{B}$ then apply mapping $\mathbf{A}$

$$\det(\mathbf{AB}) = \det(\mathbf{A}) \times \det(\mathbf{B})$$

- Determinants are only defined for square matrices, as non-square matrices project into different size spaces, where the volumes cannot be compared
- **Singular matrices** projects all points onto a lower dimensional space, so have determinant 0

Multivariate Functions

- In machine learning we heavily use multivariate functions

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

- Often our functions will be scalar value $f : \mathbb{R}^n \rightarrow \mathbb{R}$
- E.g. a learning machine for regression might be $\hat{f}(\mathbf{x}, \mathbf{w})$
- We say a function f is linear if

$$f(c\mathbf{x}) = cf(\mathbf{x}) \qquad f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$$

Multivariate Functions

- In machine learning we heavily use multivariate functions

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

- Often our functions will be scalar value $f : \mathbb{R}^n \rightarrow \mathbb{R}$

- E.g. a learning machine for regression might be $\hat{f}(\mathbf{x}, \mathbf{w})$

- We say a function f is linear if

$$f(c\mathbf{x}) = cf(\mathbf{x}) \qquad f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$$

Multivariate Functions

- In machine learning we heavily use multivariate functions

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

- Often our functions will be scalar value $f : \mathbb{R}^n \rightarrow \mathbb{R}$

- E.g. a learning machine for regression might be $\hat{f}(\mathbf{x}, \mathbf{w})$

- We say a function f is linear if

$$f(c\mathbf{x}) = cf(\mathbf{x}) \qquad f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$$

Multivariate Functions

- In machine learning we heavily use multivariate functions

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

- Often our functions will be scalar value $f : \mathbb{R}^n \rightarrow \mathbb{R}$
- E.g. a learning machine for regression might be $\hat{f}(\mathbf{x}, \mathbf{w})$ or we might have a loss function $L(\mathbf{w})$
- We say a function f is linear if

$$f(c\mathbf{x}) = cf(\mathbf{x}) \qquad f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$$

Multivariate Functions

- In machine learning we heavily use multivariate functions

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

- Often our functions will be scalar value $f : \mathbb{R}^n \rightarrow \mathbb{R}$
- E.g. a learning machine for regression might be $\hat{f}(\mathbf{x}, \mathbf{w})$ or we might have a loss function $L(\mathbf{w})$
- We say a function f is linear if

$$f(c\mathbf{x}) = cf(\mathbf{x}) \qquad f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$$

Gradients

- The gradient of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ at a point $\mathbf{x}^*$ is defined as the column vector

$$\nabla f(\mathbf{x}) = \begin{pmatrix} \frac{\partial f(\mathbf{x})}{\partial x_1} \\ \frac{\partial f(\mathbf{x})}{\partial x_2} \\ \vdots \\ \frac{\partial f(\mathbf{x})}{\partial x_n} \end{pmatrix}$$

- The gradient vector is orthogonal to the contour plane and points in the direction that the function increases in fastest

Gradients

- The gradient of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ at a point $\mathbf{x}^*$ is defined as the column vector

$$\nabla f(\mathbf{x}) = \begin{pmatrix} \frac{\partial f(\mathbf{x})}{\partial x_1} \\ \frac{\partial f(\mathbf{x})}{\partial x_2} \\ \vdots \\ \frac{\partial f(\mathbf{x})}{\partial x_n} \end{pmatrix}$$

- The gradient vector is orthogonal to the contour plane and points in the direction that the function increases in fastest

Probability

- We will have a lot to say about probability later in this course
- To reason about events with uncertain outcomes we introduce **random variables**
- Random variables corresponds to objects that we assign numbers to depending on the outcome of a random event
- E.g. we might assign 0 to the outcome of tossing a tail and 1 for a head

Probability

- We will have a lot to say about probability later in this course
- To reason about events with uncertain outcomes we introduce **random variables**
- Random variables corresponds to objects that we assign numbers to depending on the outcome of a random event
- E.g. we might assign 0 to the outcome of tossing a tail and 1 for a head

Probability

- We will have a lot to say about probability later in this course
- To reason about events with uncertain outcomes we introduce **random variables**
- Random variables corresponds to objects that we assign numbers to depending on the outcome of a random event
- E.g. we might assign 0 to the outcome of tossing a tail and 1 for a head

Probability

- We will have a lot to say about probability later in this course
- To reason about events with uncertain outcomes we introduce **random variables**
- Random variables corresponds to objects that we assign numbers to depending on the outcome of a random event
- E.g. we might assign 0 to the outcome of tossing a tail and 1 for a head

Probability

- We will have a lot to say about probability later in this course
- To reason about events with uncertain outcomes we introduce **random variables**
- Random variables corresponds to objects that we assign numbers to depending on the outcome of a random event
- E.g. we might assign 0 to the outcome of tossing a tail and 1 for a head—this is an example of a discrete random variable

Probability

- We will have a lot to say about probability later in this course
- To reason about events with uncertain outcomes we introduce **random variables**
- Random variables corresponds to objects that we assign numbers to depending on the outcome of a random event
- E.g. we might assign 0 to the outcome of tossing a tail and 1 for a head—this is an example of a discrete random variable
- A continuous random variable might be the temperature at 12 noon tomorrow in Hyde park

Probability Distributions

- For discrete events, we can define a **probability mass function** $\mathbb{P}(X = x)$ that the random variable takes the value x
- This is a number between 0 and 1 with $\sum_{x \in \mathcal{X}} \mathbb{P}(X = x) = 1$
- For continuous events the probability of X taking a particular value x is zero
- Instead we consider the probability density $f_X(x)$ which we can define as

$$f_X(x) = \lim_{\delta x \rightarrow 0} \frac{\mathbb{P}(x < X \leq x + \delta x)}{\delta x}$$

Probability Distributions

- For discrete events, we can define a **probability mass function** $\mathbb{P}(X = x)$ that the random variable takes the value x
- This is a number between 0 and 1 with $\sum_{x \in \mathcal{X}} \mathbb{P}(X = x) = 1$
- For continuous events the probability of X taking a particular value x is zero
- Instead we consider the probability density $f_X(x)$ which we can define as

$$f_X(x) = \lim_{\delta x \rightarrow 0} \frac{\mathbb{P}(x < X \leq x + \delta x)}{\delta x}$$

Probability Distributions

- For discrete events, we can define a **probability mass function** $\mathbb{P}(X = x)$ that the random variable takes the value x
- This is a number between 0 and 1 with $\sum_{x \in \mathcal{X}} \mathbb{P}(X = x) = 1$
- For continuous events the probability of X taking a particular value x is zero
- Instead we consider the probability density $f_X(x)$ which we can define as

$$f_X(x) = \lim_{\delta x \rightarrow 0} \frac{\mathbb{P}(x < X \leq x + \delta x)}{\delta x}$$

Probability Distributions

- For discrete events, we can define a **probability mass function** $\mathbb{P}(X = x)$ that the random variable takes the value x
- This is a number between 0 and 1 with $\sum_{x \in \mathcal{X}} \mathbb{P}(X = x) = 1$
- For continuous events the probability of X taking a particular value x is zero
- Instead we consider the probability density $f_X(x)$ which we can define as

$$f_X(x) = \lim_{\delta x \rightarrow 0} \frac{\mathbb{P}(x < X \leq x + \delta x)}{\delta x}$$

Expectations

- One of the most useful quantities in dealing with random numbers is to look at the average value, aka as the **expectation**, for some function $g(X)$ of the random variable

$$\mathbb{E}[g(X)] = \begin{cases} \sum_{x \in \mathcal{X}} \mathbb{P}(X = x) g(x) \\ \int f_X(x) g(x) dx \end{cases}$$

- Expectations are linear operators

$$\mathbb{E}[cg(X)] = c\mathbb{E}[g(x)]$$

$$\mathbb{E}[g(X) + h(X)] = \mathbb{E}[g(X)] + \mathbb{E}[h(X)]$$

- The expectation of a constant is itself $\mathbb{E}[c] = c$

Expectations

- One of the most useful quantities in dealing with random numbers is to look at the average value, aka as the **expectation**, for some function $g(X)$ of the random variable

$$\mathbb{E}[g(X)] = \begin{cases} \sum_{x \in \mathcal{X}} \mathbb{P}(X = x) g(x) \\ \int f_X(x) g(x) dx \end{cases}$$

- Expectations are linear operators

$$\mathbb{E}[cg(X)] = c\mathbb{E}[g(x)]$$

$$\mathbb{E}[g(X) + h(X)] = \mathbb{E}[g(X)] + \mathbb{E}[h(X)]$$

- The expectation of a constant is itself $\mathbb{E}[c] = c$

Expectations

- One of the most useful quantities in dealing with random numbers is to look at the average value, aka as the **expectation**, for some function $g(X)$ of the random variable

$$\mathbb{E}[g(X)] = \begin{cases} \sum_{x \in \mathcal{X}} \mathbb{P}(X = x) g(x) \\ \int f_X(x) g(x) dx \end{cases}$$

- Expectations are linear operators

$$\mathbb{E}[cg(X)] = c\mathbb{E}[g(x)]$$

$$\mathbb{E}[g(X) + h(X)] = \mathbb{E}[g(X)] + \mathbb{E}[h(X)]$$

- The expectation of a constant is itself $\mathbb{E}[c] = c$

Lesson

- I've covered material that we assume you know
- If you don't, you are going to struggle following any machine learning course, so grab a maths book and revise this
- We will revisit most of these topics to gain a better understanding, but you will be lost in the other machine learning modules if you are not familiar with everything I've covered
- There are exercises we will go through in the tutorial session where you can test your proficiency

Lesson

- I've covered material that we assume you know
- If you don't, you are going to struggle following any machine learning course, so grab a maths book and revise this
- We will revisit most of these topics to gain a better understanding, but you will be lost in the other machine learning modules if you are not familiar with everything I've covered
- There are exercises we will go through in the tutorial session where you can test your proficiency

Lesson

- I've covered material that we assume you know
- If you don't, you are going to struggle following any machine learning course, so grab a maths book and revise this
- We will revisit most of these topics to gain a better understanding, but you will be lost in the other machine learning modules if you are not familiar with everything I've covered
- There are exercises we will go through in the tutorial session where you can test your proficiency

Lesson

- I've covered material that we assume you know
- If you don't, you are going to struggle following any machine learning course, so grab a maths book and revise this
- We will revisit most of these topics to gain a better understanding, but you will be lost in the other machine learning modules if you are not familiar with everything I've covered
- There are exercises we will go through in the tutorial session where you can test your proficiency