Computational Biology

Lecture 15: *Cell Differentiation*



Random Boolean networks, stem cells, epigenetics

Outline

1. **Cell Differentiation**
2. The Life of an Organism
3. Epigenetic



Cell Specialisation

- Unlike unicellular organisms most multi-cellular organisms exhibit different types of cells
- By differentiating, cells can be much better adapted to a particular task
- In humans there are around 200 different types of cells
- Each cell contains the same DNA so cell differentiation involves reading a different set of instructions

Cell Specialisation

- Unlike unicellular organisms most multi-cellular organisms exhibit different types of cells
- By differentiating, cells can be much better adapted to a particular task
- In humans there are around 200 different types of cells
- Each cell contains the same DNA so cell differentiation involves reading a different set of instructions

Cell Specialisation

- Unlike unicellular organisms most multi-cellular organisms exhibit different types of cells
- By differentiating, cells can be much better adapted to a particular task
- In humans there are around 200 different types of cells
- Each cell contains the same DNA so cell differentiation involves reading a different set of instructions

Cell Specialisation

- Unlike unicellular organisms most multi-cellular organisms exhibit different types of cells
- By differentiating, cells can be much better adapted to a particular task
- In humans there are around 200 different types of cells
- Each cell contains the same DNA so cell differentiation involves reading a different set of instructions

Cell Functions

- Cell function is determined by the expression of different genes
- The expression of genes are determined by the transcription factors (proteins that interact with the promoter sites either exciting or inhibiting the transcription of a gene)
- Cells can respond to their environment by the interaction between signalling molecules and proteins
- Thus, each gene type will respond differently to its environment

Cell Functions

- Cell function is determined by the expression of different genes
- The expression of genes are determined by the transcription factors (proteins that interact with the promoter sites either exciting or inhibiting the transcription of a gene)
- Cells can respond to their environment by the interaction between signalling molecules and proteins
- Thus, each gene type will respond differently to its environment

Cell Functions

- Cell function is determined by the expression of different genes
- The expression of genes are determined by the transcription factors (proteins that interact with the promoter sites either exciting or inhibiting the transcription of a gene)
- Cells can respond to their environment by the interaction between signalling molecules and proteins
- Thus, each gene type will respond differently to its environment

Cell Functions

- Cell function is determined by the expression of different genes
- The expression of genes are determined by the transcription factors (proteins that interact with the promoter sites either exciting or inhibiting the transcription of a gene)
- Cells can respond to their environment by the interaction between signalling molecules and proteins
- Thus, each gene type will respond differently to its environment

Regulation by Signal



Regulation by Signal



Regulation by Signal



Regulation by Signal



Regulation by Signal



Regulation by Signal



Regulation by Signal



Regulation by Signal



Regulation by Signal



Regulation by Signal



Regulation by Signal



Regulation by Signal



Regulation by Signal



Regulation by Signal



Regulation by Signal



Gene Regulation Networks

- Proteins control each other's expression forming complex gene regulation networks
- The pattern of interaction determines the type of cell
- Once the type is determine cells overwhelming stay in that type

Gene Regulation Networks

- Proteins control each other's expression forming complex gene regulation networks
- The pattern of interaction determines the type of cell
- Once the type is determine cells overwhelming stay in that type

Gene Regulation Networks

- Proteins control each other's expression forming complex gene regulation networks
- The pattern of interaction determines the type of cell
- Once the type is determine cells overwhelming stay in that type

Large Gene Regulation Network



Random Boolean Networks

k	$f_k(F,F)$	$f_k(F,T)$	$f_k(T,F)$	$f_k(T,T)$
0	F	F	F	F
1	T	F	F	F
2	F	T	F	F
3	T	T	F	F
4	F	F	T	F
5	T	F	T	F
6	F	T	T	F
7	T	T	T	F
8	F	F	F	T
9	T	F	F	T
10	F	T	F	T
11	T	T	F	T
12	F	F	T	T
13	T	F	T	T
14	F	T	T	T
15	T	T	T	T

- To understand gene dynamics researchers (lead by Stuart Kauffman) considered the dynamics of random Boolean networks
- Each node is connected to k_{in} other nodes uniformly at random
- The nodes perform a randomly selected Boolean functions

Random Boolean Networks

k	$f_k(F,F)$	$f_k(F,T)$	$f_k(T,F)$	$f_k(T,T)$
0	F	F	F	F
1	T	F	F	F
2	F	T	F	F
3	T	T	F	F
4	F	F	T	F
5	T	F	T	F
6	F	T	T	F
7	T	T	T	F
8	F	F	F	T
9	T	F	F	T
10	F	T	F	T
11	T	T	F	T
12	F	F	T	T
13	T	F	T	T
14	F	T	T	T
15	T	T	T	T

- To understand gene dynamics researchers (lead by Stuart Kauffman) considered the dynamics of random Boolean networks
- Each node is connected to k_{in} other nodes uniformly at random
- The nodes perform a randomly selected Boolean functions

Random Boolean Networks

k	$f_k(F,F)$	$f_k(F,T)$	$f_k(T,F)$	$f_k(T,T)$
0	F	F	F	F
1	T	F	F	F
2	F	T	F	F
3	T	T	F	F
4	F	F	T	F
5	T	F	T	F
6	F	T	T	F
7	T	T	T	F
8	F	F	F	T
9	T	F	F	T
10	F	T	F	T
11	T	T	F	T
12	F	F	T	T
13	T	F	T	T
14	F	T	T	T
15	T	T	T	T

- To understand gene dynamics researchers (lead by Stuart Kauffman) considered the dynamics of random Boolean networks
- Each node is connected to k_{in} other nodes uniformly at random
- The nodes perform a randomly selected Boolean functions

Dynamics of RBN



Dynamics of RBN



Dynamics of RBN



Dynamics of RBN



Dynamics of RBN



Dynamics of RBN



Dynamics of RBN



Dynamics of RBN



Attractors



Phase Transition

- The behaviour of random Boolean networks shows two phases depending on the number of connections and the randomness of the functions
  - ⋆ For sparsely connected networks (in-degree ≤ 2) there are a small number of attractors which are reached after a small number of iterations
  - ⋆ Otherwise there are an exponential number of attractors which take exponential time to reach—the "chaotic" phase
- It has been argued that biology lives on the "edge of chaos" where there are only a few attractors, but the network is adaptable

Phase Transition

- The behaviour of random Boolean networks shows two phases depending on the number of connections and the randomness of the functions
  - ⋆ For sparsely connected networks (in-degree ≤ 2) there are a small number of attractors which are reached after a small number of iterations
  - ⋆ Otherwise there are an exponential number of attractors which take exponential time to reach—the "chaotic" phase
- It has been argued that biology lives on the "edge of chaos" where there are only a few attractors, but the network is adaptable

Phase Transition

- The behaviour of random Boolean networks shows two phases depending on the number of connections and the randomness of the functions
  - ⋆ For sparsely connected networks (in-degree ≤ 2) there are a small number of attractors which are reached after a small number of iterations
  - ⋆ Otherwise there are an exponential number of attractors which take exponential time to reach—the "chaotic" phase
- It has been argued that biology lives on the "edge of chaos" where there are only a few attractors, but the network is adaptable

Phase Transition

- The behaviour of random Boolean networks shows two phases depending on the number of connections and the randomness of the functions
  - ⋆ For sparsely connected networks (in-degree ≤ 2) there are a small number of attractors which are reached after a small number of iterations
  - ⋆ Otherwise there are an exponential number of attractors which take exponential time to reach—the "chaotic" phase
- It has been argued that biology lives on the "edge of chaos" where there are only a few attractors, but the network is adaptable

Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Dynamics of RBN in Chaotic Phase



Real Biology

- Of course, biological systems are more complex
- It is very difficult to turn one type of cell into another
- When cells divide they faithfully remember what type they are
- There are influences beyond the current protein interactions that affect a cell's type (epigenetics)

Real Biology

- Of course, biological systems are more complex
- It is very difficult to turn one type of cell into another
- When cells divide they faithfully remember what type they are
- There are influences beyond the current protein interactions that affect a cell's type (epigenetics)

Real Biology

- Of course, biological systems are more complex
- It is very difficult to turn one type of cell into another
- When cells divide they faithfully remember what type they are
- There are influences beyond the current protein interactions that affect a cell's type (epigenetics)

Real Biology

- Of course, biological systems are more complex
- It is very difficult to turn one type of cell into another
- When cells divide they faithfully remember what type they are
- There are influences beyond the current protein interactions that affect a cell's type (epigenetics)

Outline

1. Cell Differentiation
2. **The Life of an Organism**
3. Epigenetic



In the Beginning

- The life cycle of multi-cellular organisms are complex
- They begin life as a single embryo cell
- This cell will divide to form all the specialised cells of the organism (and the placenta)
- The embryo cell can make any other cell and is known as totipotent
- The embryo grows through cell division and quickly differentiates into cells of the organism and cells of the placenta

In the Beginning

- The life cycle of multi-cellular organisms are complex
- They begin life as a single embryo cell
- This cell will divide to form all the specialised cells of the organism (and the placenta)
- The embryo cell can make any other cell and is known as totipotent
- The embryo grows through cell division and quickly differentiates into cells of the organism and cells of the placenta

In the Beginning

- The life cycle of multi-cellular organisms are complex
- They begin life as a single embryo cell
- This cell will divide to form all the specialised cells of the organism (and the placenta)
- The embryo cell can make any other cell and is known as totipotent
- The embryo grows through cell division and quickly differentiates into cells of the organism and cells of the placenta

In the Beginning

- The life cycle of multi-cellular organisms are complex
- They begin life as a single embryo cell
- This cell will divide to form all the specialised cells of the organism (and the placenta)
- The embryo cell can make any other cell and is known as totipotent
- The embryo grows through cell division and quickly differentiates into cells of the organism and cells of the placenta

In the Beginning

- The life cycle of multi-cellular organisms are complex
- They begin life as a single embryo cell
- This cell will divide to form all the specialised cells of the organism (and the placenta)
- The embryo cell can make any other cell and is known as totipotent
- The embryo grows through cell division and quickly differentiates into cells of the organism and cells of the placenta

Embryonic Stem Cells

- The cells in the organism are no longer able to form a placenta, but can produce any other cell in the body
- These **embryonic stem cells** are said to be **pluripotent**
- The embryonic stem cells can change to more specialised stem cells which are said to be *multipotent*
- These will eventually specialise to the normal cells of the body known as the **soma**

Embryonic Stem Cells

- The cells in the organism are no longer able to form a placenta, but can produce any other cell in the body
- These **embryonic stem cells** are said to be **pluripotent**
- The embryonic stem cells can change to more specialised stem cells which are said to be *multipotent*
- These will eventually specialise to the normal cells of the body known as the **soma**

Embryonic Stem Cells

- The cells in the organism are no longer able to form a placenta, but can produce any other cell in the body
- These **embryonic stem cells** are said to be **pluripotent**
- The embryonic stem cells can change to more specialised stem cells which are said to be *multipotent*
- These will eventually specialise to the normal cells of the body known as the **soma**

Embryonic Stem Cells

- The cells in the organism are no longer able to form a placenta, but can produce any other cell in the body
- These **embryonic stem cells** are said to be **pluripotent**
- The embryonic stem cells can change to more specialised stem cells which are said to be *multipotent*
- These will eventually specialise to the normal cells of the body known as the **soma**

Specialisation



Gilbert 9th Edition Fig. 1.7

Somatic vs. Germline cell lineages

Causes of Specialisation





- There are a number of routes to specialisation
- During the development of the embryo some transcription factors aggregate, at cell division this leads to differentiation in the cell function
- Other cells are known to specialise depending on signalling molecules that occur at receptors

Causes of Specialisation





- There are a number of routes to specialisation
- During the development of the embryo some transcription factors aggregate, at cell division this leads to differentiation in the cell function
- Other cells are known to specialise depending on signalling molecules that occur at receptors

Causes of Specialisation





- There are a number of routes to specialisation
- During the development of the embryo some transcription factors aggregate, at cell division this leads to differentiation in the cell function
- Other cells are known to specialise depending on signalling molecules that occur at receptors

Outline

1. Cell Differentiation
2. The Life of an Organism
3. **Epigenetic**



Cell Division

- Cells in the body constantly divide
- They all carry the same DNA code
- But, the DNA is often modified by methylation and acetylation
- These can be facilitated, e.g. by proteins called DNA $methyltransferase$ (DNMT)

Cell Division

- Cells in the body constantly divide
- They all carry the same DNA code
- But, the DNA is often modified by methylation and acetylation
- These can be facilitated, e.g. by proteins called DNA $methyltransferase$ (DNMT)

Cell Division

- Cells in the body constantly divide
- They all carry the same DNA code
- But, the DNA is often modified by methylation and acetylation
- These can be facilitated, e.g. by proteins called DNA $methyltransferase$ (DNMT)

Cell Division

- Cells in the body constantly divide
- They all carry the same DNA code
- But, the DNA is often modified by methylation and acetylation
- These can be facilitated, e.g. by proteins called *DNA methyltransferase* (DNMT)

DNA Methylation



Histone Acetylation



Epigenetics

- Acetylation suppresses the expression of genes
- DNA methylation can suppress or activate the expression of genes
- Stem cells have more methylated sites than somatic (normal) cells, showing that this is an important mechanism for ensuring cell specialisation is maintained during cell division
- The methylation and acetylation is modified by proteins
- Epigenetics can be seen as another component in a complex Boolean network

Epigenetics

- Acetylation suppresses the expression of genes
- DNA methylation can suppress or activate the expression of genes
- Stem cells have more methylated sites than somatic (normal) cells, showing that this is an important mechanism for ensuring cell specialisation is maintained during cell division
- The methylation and acetylation is modified by proteins
- Epigenetics can be seen as another component in a complex Boolean network

Epigenetics

- Acetylation suppresses the expression of genes
- DNA methylation can suppress or activate the expression of genes
- Stem cells have more methylated sites than somatic (normal) cells, showing that this is an important mechanism for ensuring cell specialisation is maintained during cell division
- The methylation and acetylation is modified by proteins
- Epigenetics can be seen as another component in a complex Boolean network

Epigenetics

- Acetylation suppresses the expression of genes
- DNA methylation can suppress or activate the expression of genes
- Stem cells have more methylated sites than somatic (normal) cells, showing that this is an important mechanism for ensuring cell specialisation is maintained during cell division
- The methylation and acetylation is modified by proteins
- Epigenetics can be seen as another component in a complex Boolean network

Epigenetics

- Acetylation suppresses the expression of genes
- DNA methylation can suppress or activate the expression of genes
- Stem cells have more methylated sites than somatic (normal) cells, showing that this is an important mechanism for ensuring cell specialisation is maintained during cell division
- The methylation and acetylation is modified by proteins
- Epigenetics can be seen as another component in a complex Boolean network

Conclusion

- Cell differentiation is a complex biological process that is essential to multi-cellular organism
- Simplified models of Boolean networks suggests that a small number of attractors might naturally occur in an interacting network
- Whether this "explains" multi-cellular organisms or their origin is a matter of debate
- Specialisation happens through a hierarchical series of steps
- Epigenetics seem to be an important component in determining which proteins are expressed

Conclusion

- Cell differentiation is a complex biological process that is essential to multi-cellular organism
- Simplified models of Boolean networks suggests that a small number of attractors might naturally occur in an interacting network
- Whether this "explains" multi-cellular organisms or their origin is a matter of debate
- Specialisation happens through a hierarchical series of steps
- Epigenetics seem to be an important component in determining which proteins are expressed

Conclusion

- Cell differentiation is a complex biological process that is essential to multi-cellular organism
- Simplified models of Boolean networks suggests that a small number of attractors might naturally occur in an interacting network
- Whether this "explains" multi-cellular organisms or their origin is a matter of debate
- Specialisation happens through a hierarchical series of steps
- Epigenetics seem to be an important component in determining which proteins are expressed

Conclusion

- Cell differentiation is a complex biological process that is essential to multi-cellular organism
- Simplified models of Boolean networks suggests that a small number of attractors might naturally occur in an interacting network
- Whether this "explains" multi-cellular organisms or their origin is a matter of debate
- Specialisation happens through a hierarchical series of steps
- Epigenetics seem to be an important component in determining which proteins are expressed

Conclusion

- Cell differentiation is a complex biological process that is essential to multi-cellular organism
- Simplified models of Boolean networks suggests that a small number of attractors might naturally occur in an interacting network
- Whether this "explains" multi-cellular organisms or their origin is a matter of debate
- Specialisation happens through a hierarchical series of steps
- Epigenetics seem to be an important component in determining which proteins are expressed