Computational Biology

Lecture 11: *Population Genetics*



Fisher-Wright model, ODEs, Markov chains.

Outline

1. **Population Genetics**
2. Facts of Life
3. Infinite Population Model
4. Finite Population Model



Evolution and Molecular Biology

- Evolution is the theory that underpins our understanding of biology
- Molecular biology has open up a whole new perspective on evolution as we see similar proteins in related species
- This evolution ancestry helps us understand molecular biology as similar proteins usually share the same function
- However we can also use molecular biology to help understand evolution—more of this when we discuss phylogenetic tree
- But how does evolution work?

Evolutionary Population Dynamics

- Since as least the 1920's researchers have tried to model evolution
- The field is known as **population genetics**
- Much of it was developed before we understood the mechanisms of heredity
- It is therefore rather abstracted from biology
- However it was enormously influential to our understanding of evolution

Outline

1. Population Genetics
2. **Facts of Life**
3. Infinite Population Model
4. Finite Population Model



Fisher-Wright Model

- The classic model of population genetics is the Fisher-Wright (or Wright-Fisher) model developed independently by Fisher and Wright around 1930
- It is important because it laid the theoretical framework for many of the prejudices held today
- In its simplest form it is a model of what happens at a single locus (site) under selection, mutation and crossover



Locus and Alleles

- The locus can be thought of as the location of a gene
- The type of gene at a site will differ between individuals (e.g. one for smooth peas and one for wrinkled peas)
- The different types of genes are known as **alleles**
- We assume that genes can be modified from one allele to another by mutation
- Alleles are also exchanged during sexual reproduction (crossover)

Fitness

- Fitness has a technical meaning in population genetics; it is a measure of the expected number of offspring an individual produces
- If one individual is 10% fitter than another it will (in expectation) produce 10% more offspring
- Of course, the individual may get flattened by a car before it reaches sexual maturity and never produce any offspring
- Thus fitness measures potential to reproduce

Epistatic Interaction

- Many genes interact (we now know they often code for proteins that directly interact chemically in many cases)
- Thus the fitness of an allele will typically depend on other alleles at other locations
- This interaction is known as **epistasis**
- To keep things simple in the Fisher-Wright model we assume no epistasis
- There exists more complex models which treat epistasis between two or more loci, but these rapidly become rather complex

Sex

- Sex is surprisingly ubiquitous in higher organisms
- Few higher organisms seem to reproduce asexually (although even if they do, many sometimes also reproduce sexually)
- Asexual higher organisms all would have produced sexually sometime in their past
- Why sex is surprising is that if females produce asexually (producing clones) then they breed at twice the rate of mixed couples—hence they are twice as fit
- Sexual reproduction has to give at least a two fold advantage in fitness on a very short time scale to survive

Linkage Equilibrium

- As well as no epistasis the model assumes lots of crossover (compared to selection and mutation)
- This is not unrealistic in many cases
- Under this assumption we can treat the loci independently and concentrate on a single locus
- This is known as **linkage equilibrium**
- We therefore only need to consider the proportion of the different alleles at the locus
- In an asexual population, the alleles at different sites are coupled and the situation is much harder (in fact, unsolved)

Diploidy

- Most organisms are diploids, that is they carry two copies of DNA; one from their father and one from their mother
- Like sex, it is not altogether clear what advantage diploidy brings
- Clearly, diploids can tolerate mutations that make one set of genes dysfunctional
- But, this also allows detrimental mutations to build up in a genome
- Incidentally, mating between close relatives is likely to create nonviable offspring because it is likely to combine two dysfunctional alleles

Haploidy

- To keep things simple we are going to assume each individual only carries one set of genes
- Organisms that do this are said to be **haploidy**
- There are various organisms (e.g. male bees, wasps, ants, many fungi) that are haploid
- Generalising to diploid organisms is just fiddly—it mainly involves adding in factors of two every so often, but it does not substantially change things
- Incidentally many plants are polypoids having more than two copies of each gene

Gender

- Life tends also to be gendered
- This possibly just reflects that finding a mate consumes energy and by specialisation some energy can be saved
- However, this is another complicating factor, which adds little to the model
- I will assume a **monoecious** population, i.e. one with only one gender, although it still mates
- These are all simplifying assumptions, but it allows us to concentrate on the role of selection, mutation and crossover

Outline

1. Population Genetics
2. Facts of Life
3. **Infinite Population Model**
4. Finite Population Model



Single Locus Model

- We consider the case where the allele can take just two states A and B
- We call B the mutant
- We make the following assumptions
  - B has a selective advantage of $1 + s$
  - A mutates to B with a forward mutation rate of u
  - B mutates to A with a backward mutation rate of v
  - The population size, P, is unchanged from generation to generation

Many Seeds

- We consider the population changing from one generation to the next by a three stage process
  - Each individual produces a very large number of seeds in proportional to its fitness (selective advantage)
  - The seeds are mutated according to the mutation probabilities
  - We select a sample of P of the seeds independently to form the next generation
- Initially we assume P is infinite and consider only the proportion of mutants in the population

Dynamics

- Assume we start with n mutants
- After selection we have a proportion p_s of mutant seeds

$$p_s = \frac{(1+s)n}{(1+s)n + P - n} = \frac{(1+s)n}{P + s\,n}$$

- After mutation we have a proportion p_{sm} of mutants

$$p_{sm} = (1-v)p_s + u(1-p_s) = \frac{(1-v)(1+s)n + u(P-n)}{P + s\,n}$$

Infinite Population Approximation

- If we assume an infinite population then this will be the proportion of individuals at the next generation

$$p(t+1) = \frac{(1-v)(1+s)p(t) + u\big(1-p(t)\big)}{1 + s\,p(t)}$$

- If the effect of selection and mutation is small then we can approximate the dynamics by a differential equation

$$\frac{\mathrm{d}p}{\mathrm{d}t} \approx p(t+1) - p(t) = \frac{\big(u - (u+v-s+u\,s)p(t) + s\,p^2(t)\big)}{1 + s\,p(t)}$$

Infinite Population Prediction



$s = 0.1,\ u = v = 0.001$

Difficulties of Infinite Populations

- In an infinite population a mutation (even if it is very unlikely) will always occur in a small fraction of individuals
- Selection can rapidly amplify the mutants if they are fitter—note fitter individuals initial grow exponentially

$$(1+s)^t = \mathrm{e}^{t\,\log(1+s)} \approx \mathrm{e}^{s\,t}$$

- If the mutation rate is small then in finite populations the mutant may take a very long time to appear
- Thus infinite populations can have far more rapid take over times than is experience in large, but finite, populations

Outline

1. Population Genetics
2. Facts of Life
3. Infinite Population Model
4. **Finite Population Model**



Finite Populations

- To model finite populations, we have to draw a sample of P individuals from the seeds
- The probability of drawing $n' = n(t+1)$ mutants given a proportion $p_{sm}(n(t))$ of mutants in the seed population is given by a binomial distribution

$$\mathbb{P}(n') = \binom{P}{n'} p_{sm}^{n'} \left(1 - p_{sm}\right)^{P-n'}$$

- (Recall that p_{sm} is the proportion of mutant seeds after selection and mutation and depends on $n(t)$)
- We can simulate the evolution of a population, by drawing a random binomial deviate at each generation

Finite Population Run



$s = 0.1,\ u = v = 0.001$

Markov Models

- The evolving population can be modelled as a Markov chain
- The population at time t depends only on the population at time $t-1$
- It is *conditionally independent* of the population at earlier times *given* the population at time $t-1$
- We define $p_n(t) = \mathbb{P}(n(t) = n)$ to be the probability that there are n mutants at time t
- Let $W(n \to n')$ be the transition probability of a population with n mutants going to a population of n' mutants in the next generation

$$p_{n'}(t+1) = \sum_{n=0}^{P} W(n \to n') p_n(t)$$

Matrix Formulation

- Denote the *probability vector* $(p_0(t), p_1(t), \ldots, p_P(t))$ by $\boldsymbol{p}(t)$
- Let $\mathbf{W}$ be the *transition matrix* with elements $W_{n'n} = W(n \to n')$
- The dynamics can be written in matrix form

$$\boldsymbol{p}(t) = \mathbf{W}\boldsymbol{p}(t-1) = \mathbf{W}\left(\mathbf{W}\boldsymbol{p}(t-2)\right) = \mathbf{W}\mathbf{W}\cdots\mathbf{W}\boldsymbol{p}(0) = \mathbf{W}^t\boldsymbol{p}(0)$$

- Where $\boldsymbol{p}(t=0)$ encodes our *initial conditions*
- If initially $n(t=0) = 0$ then $p_0(t=0) = 1$ and $p_i(t=0) = 0, \forall i > 0$

Markov Analysis

- In our problem the transition matrix, $\mathbf{W}$, has elements

$$W_{n'n} = \binom{P}{n'} \left(p_{sm}(n)\right)^{n'} \left(1 - p_{sm}(n)\right)^{P-n'}$$

- where $p_{sm}(n)$ is given by

$$p_{sm}(n) = \frac{(1-v)(1+s)n + u(P-n)}{P + s\,n}$$

- This provides an exact formulation of a finite population

Example of Markov Analysis



Mean Number of Mutants



Problems with Markov Chain Analysis

- The Markov Chain analysis is exact and provides a full picture of the probability of the population being in a certain state
- Unfortunately, the transition matrix $W(n \to n')$ is of size the number of states squared
- The number of states for a population of size P is $P + 1$
- For a large population writing the transition matrix becomes infeasible
- Even when it is feasible we don't get much intuition on changing the parameters

Conclusion

- The Fisher-Wright model provides a simple model undergoing selection, mutation and crossover
- It allows us to explore the effects of selection pressure and mutation rate
- We can solve the infinite population model directly or by solving an ordinary differential equation (ODE)
- Unfortunately infinite population models don't always capture the behaviour of finite populations even when the population is very large
- We can simulate finite population models
- For small populations we can use Markov models