Lecture 13: *Phylogenetic Trees*



Trees of Life, UPGMA

Outline

1. **Importance of Phylogeny**
2. Distance Based Trees



Giant Panda

- Is a giant panda a bear or a raccoon?
- When they were first discovered in Europe scientists couldn't decide—they had features of both
- They were named after red pandas that are raccoons
- This was only resolved in 1985 when Steven O'Brien and colleagues used sequence matching from 500 000 nucleotides to finally produce a definitive evolutionary tree

Giant Panda

- Is a giant panda a bear or a raccoon?
- When they were first discovered in Europe scientists couldn't decide—they had features of both
- They were named after red pandas that are raccoons
- This was only resolved in 1985 when Steven O'Brien and colleagues used sequence matching from 500 000 nucleotides to finally produce a definitive evolutionary tree

Giant Panda

- Is a giant panda a bear or a raccoon?
- When they were first discovered in Europe scientists couldn't decide—they had features of both
- They were named after red pandas that are raccoons
- This was only resolved in 1985 when Steven O'Brien and colleagues used sequence matching from 500 000 nucleotides to finally produce a definitive evolutionary tree

Giant Panda

- Is a giant panda a bear or a raccoon?
- When they were first discovered in Europe scientists couldn't decide—they had features of both
- They were named after red pandas that are raccoons
- This was only resolved in 1985 when Steven O'Brien and colleagues used sequence matching from 500 000 nucleotides to finally produce a definitive evolutionary tree

The Tree of Bears



Out of Africa

- In the mid 1980s Allan Wilson and colleagues studied the evolutionary tree of human mitrochondria and decided that all races outside Africa shared a common ancestor who lived in Africa 200 000 years ago

- Hominids had left Africa much earlier (e.g. Neanderthals) so this theory postulated that modern humans would have displaced all existing hominids outside of Africa

Out of Africa

- In the mid 1980s Allan Wilson and colleagues studied the evolutionary tree of human mitrochondria and decided that all races outside Africa shared a common ancestor who lived in Africa 200 000 years ago

- Hominids had left Africa much earlier (e.g. Neanderthals) so this theory postulated that modern humans would have displaced all existing hominids outside of Africa

Modern Humans



By NordNordWest - File:Spreading homo sapiens ru.svg by Urutseg, Public Domain, https://commons.wikimedia.org/w/index.php?curid=34697001

Mitrochondrial DNA

- Mitrochondria are the powerhouse of cells
- They would have originally been an independent organism that somehow got captured by eukaryotic cells
- They contain their own DNA
- They are inherited through the mother's ova
- They therefore reproduce asexually

Mitrochondrial DNA

- Mitrochondria are the powerhouse of cells
- They would have originally been an independent organism that somehow got captured by eukaryotic cells
- They contain their own DNA
- They are inherited through the mother's ova
- They therefore reproduce asexually

Mitrochondrial DNA

- Mitrochondria are the powerhouse of cells
- They would have originally been an independent organism that somehow got captured by eukaryotic cells
- They contain their own DNA
- They are inherited through the mother's ova
- They therefore reproduce asexually

Mitrochondrial DNA

- Mitrochondria are the powerhouse of cells
- They would have originally been an independent organism that somehow got captured by eukaryotic cells
- They contain their own DNA
- They are inherited through the mother's ova
- They therefore reproduce asexually

Mitrochondrial DNA

- Mitrochondria are the powerhouse of cells
- They would have originally been an independent organism that somehow got captured by eukaryotic cells
- They contain their own DNA
- They are inherited through the mother's ova
- They therefore reproduce asexually

Asexual Inheritance

- Asexual inheritance is much easier to study
- Populations will share a common ancestor because of
  - ⋆ Natural coalescence
  - ⋆ Going through a bottle neck (e.g. leaving Africa)
- Natural coalescence happens by chance, but will be enhanced by selection

Asexual Inheritance

- Asexual inheritance is much easier to study
- Populations will share a common ancestor because of
  - ⋆ Natural coalescence
  - ⋆ Going through a bottle neck (e.g. leaving Africa)
- Natural coalescence happens by chance, but will be enhanced by selection

Asexual Inheritance

- Asexual inheritance is much easier to study
- Populations will share a common ancestor because of
  - ⋆ Natural coalescence
  - ⋆ Going through a bottle neck (e.g. leaving Africa)
- Natural coalescence happens by chance, but will be enhanced by selection

Asexual Inheritance

- Asexual inheritance is much easier to study
- Populations will share a common ancestor because of
  - ⋆ Natural coalescence
  - ⋆ Going through a bottle neck (e.g. leaving Africa)
- Natural coalescence happens by chance, but will be enhanced by selection

Coalescence

Coalescence



Coalescence



Coalescence



Coalescence



Coalescence

Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



Coalescence



- If you go far enough back in time an asexual population will have a common ancestor

Cross-Breeding

- Didn't modern man cross-breed with other hominids?
- This was unknown even 10 years ago
- However, the numbers were not large and due to coalescence (the coalescence time being small because there was not huge cross-breeding) all traces of Neanderthal mitrochondrial DNA vanished
- The story is very different for the rest of our DNA
- Normal DNA evolves sexually

Cross-Breeding

- Didn't modern man cross-breed with other hominids?
- This was unknown even 10 years ago
- However, the numbers were not large and due to coalescence (the coalescence time being small because there was not huge cross-breeding) all traces of Neanderthal mitrochondrial DNA vanished
- The story is very different for the rest of our DNA
- Normal DNA evolves sexually

Cross-Breeding

- Didn't modern man cross-breed with other hominids?
- This was unknown even 10 years ago, but of course they did
- However, the numbers were not large and due to coalescence (the coalescence time being small because there was not huge cross-breeding) all traces of Neanderthal mitrochondrial DNA vanished
- The story is very different for the rest of our DNA
- Normal DNA evolves sexually

Cross-Breeding

- Didn't modern man cross-breed with other hominids?
- This was unknown even 10 years ago, but of course they did
- However, the numbers were not large and due to coalescence (the coalescence time being small because there was not huge cross-breeding) all traces of Neanderthal mitrochondrial DNA vanished
- The story is very different for the rest of our DNA
- Normal DNA evolves sexually

Cross-Breeding

- Didn't modern man cross-breed with other hominids?
- This was unknown even 10 years ago, but of course they did
- However, the numbers were not large and due to coalescence (the coalescence time being small because there was not huge cross-breeding) all traces of Neanderthal mitrochondrial DNA vanished
- The story is very different for the rest of our DNA
- Normal DNA evolves sexually

Cross-Breeding

- Didn't modern man cross-breed with other hominids?
- This was unknown even 10 years ago, but of course they did
- However, the numbers were not large and due to coalescence (the coalescence time being small because there was not huge cross-breeding) all traces of Neanderthal mitrochondrial DNA vanished
- The story is very different for the rest of our DNA
- Normal DNA evolves sexually

The Selfish Gene

- With crossover you cannot build a tree of life as you inherit from two parents
- However, each gene (at least if it doesn't get disrupted by crossover) can be mapped to its own ancestral tree
- Because there are lots of genes it is much more likely for genes from different origins to survive (particularly if they have different fitness)
- In fact, we carry DNA that clearly came from Neanderthal and a few other early hominids
- These genes seem to have been selected for

The Selfish Gene

- With crossover you cannot build a tree of life as you inherit from two parents
- However, each gene (at least if it doesn't get disrupted by crossover) can be mapped to its own ancestral tree
- Because there are lots of genes it is much more likely for genes from different origins to survive (particularly if they have different fitness)
- In fact, we carry DNA that clearly came from Neanderthal and a few other early hominids
- These genes seem to have been selected for

The Selfish Gene

- With crossover you cannot build a tree of life as you inherit from two parents
- However, each gene (at least if it doesn't get disrupted by crossover) can be mapped to its own ancestral tree
- Because there are lots of genes it is much more likely for genes from different origins to survive (particularly if they have different fitness)
- In fact, we carry DNA that clearly came from Neanderthal and a few other early hominids
- These genes seem to have been selected for

The Selfish Gene

- With crossover you cannot build a tree of life as you inherit from two parents
- However, each gene (at least if it doesn't get disrupted by crossover) can be mapped to its own ancestral tree
- Because there are lots of genes it is much more likely for genes from different origins to survive (particularly if they have different fitness)
- In fact, we carry DNA that clearly came from Neanderthal and a few other early hominids
- These genes seem to have been selected for

The Selfish Gene

- With crossover you cannot build a tree of life as you inherit from two parents
- However, each gene (at least if it doesn't get disrupted by crossover) can be mapped to its own ancestral tree
- Because there are lots of genes it is much more likely for genes from different origins to survive (particularly if they have different fitness)
- In fact, we carry DNA that clearly came from Neanderthal and a few other early hominids
- These genes seem to have been selected for

Horizontal Gene Transfer

- Although bacteria and archaea reproduce asexually they regularly exchange plasmids
- These are loops of DNA
- This is the primary cause of antibiotic resistance
- The tree of life is not quite a tree



By Dr. Smets and perhaps others - Barth F. Smets, Ph.D., with permission, Attribution, https://commons.wikimedia.org/w/index.php?curid=5720886

Horizontal Gene Transfer

- Although bacteria and archaea reproduce asexually they regularly exchange plasmids
- These are loops of DNA
- This is the primary cause of antibiotic resistance
- The tree of life is not quite a tree



By Dr. Smets and perhaps others - Barth F. Smets, Ph.D., with permission, Attribution, https://commons.wikimedia.org/w/index.php?curid=5720886

Horizontal Gene Transfer

- Although bacteria and archaea reproduce asexually they regularly exchange plasmids
- These are loops of DNA
- This is the primary cause of antibiotic resistance
- The tree of life is not quite a tree



By Dr. Smets and perhaps others - Barth F. Smets, Ph.D., with permission, Attribution, https://commons.wikimedia.org/w/index.php?curid=5720886

Horizontal Gene Transfer

- Although bacteria and archaea reproduce asexually they regularly exchange plasmids
- These are loops of DNA
- This is the primary cause of antibiotic resistance
- The tree of life is not quite a tree



By Dr. Smets and perhaps others - Barth F. Smets, Ph.D., with permission, Attribution, https://commons.wikimedia.org/w/index.php?curid=5720886

Ligers and Tigons

- Most higher animals form species where individuals between species don't inter-breed
- In captivity ligers are crosses between male lions and female tigers and tigons are crosses between male tigers and female lions
- In the wild lions and tigers no longer overlap, but in the past there was some report of ligers (which are typically enormous)
- Like most cross-breeds they have reduced fertility (one sex is often inviable or sterile) so that cross breeds die out
- Because of this lack of significant cross-breeding we can model most of evolution at a species level as a tree

Ligers and Tigons

- Most higher animals form species where individuals between species don't inter-breed very much
- In captivity ligers are crosses between male lions and female tigers and tigons are crosses between male tigers and female lions
- In the wild lions and tigers no longer overlap, but in the past there was some report of ligers (which are typically enormous)
- Like most cross-breeds they have reduced fertility (one sex is often inviable or sterile) so that cross breeds die out
- Because of this lack of significant cross-breeding we can model most of evolution at a species level as a tree

Ligers and Tigons

- Most higher animals form species where individuals between species don't inter-breed very much
- In captivity ligers are crosses between male lions and female tigers and tigons are crosses between male tigers and female lions
- In the wild lions and tigers no longer overlap, but in the past there was some report of ligers (which are typically enormous)
- Like most cross-breeds they have reduced fertility (one sex is often inviable or sterile) so that cross breeds die out
- Because of this lack of significant cross-breeding we can model most of evolution at a species level as a tree

Ligers and Tigons

- Most higher animals form species where individuals between species don't inter-breed very much
- In captivity ligers are crosses between male lions and female tigers and tigons are crosses between male tigers and female lions
- In the wild lions and tigers no longer overlap, but in the past there was some report of ligers (which are typically enormous)
- Like most cross-breeds they have reduced fertility (one sex is often inviable or sterile) so that cross breeds die out
- Because of this lack of significant cross-breeding we can model most of evolution at a species level as a tree

Ligers and Tigons

- Most higher animals form species where individuals between species don't inter-breed very much
- In captivity ligers are crosses between male lions and female tigers and tigons are crosses between male tigers and female lions
- In the wild lions and tigers no longer overlap, but in the past there was some report of ligers (which are typically enormous)
- Like most cross-breeds they have reduced fertility (one sex is often inviable or sterile) so that cross breeds die out
- Because of this lack of significant cross-breeding we can model most of evolution at a species level as a tree

Ligers and Tigons

- Most higher animals form species where individuals between species don't inter-breed very much
- In captivity ligers are crosses between male lions and female tigers and tigons are crosses between male tigers and female lions
- In the wild lions and tigers no longer overlap, but in the past there was some report of ligers (which are typically enormous)
- Like most cross-breeds they have reduced fertility (one sex is often inviable or sterile) so that cross breeds die out
- Because of this lack of significant cross-breeding we can model most of evolution at a species level as a tree

Evolution and HIV

- We often think of evolution as happening over geological time
- But the HIV virus evolved over decades from a disease that infected African primates to one that infects man
- Furthermore it evolves over months and years to evade our immune system
- Tracking these changes can be important in understanding and controlling the disease
- The virus evolves asexually so we can build an evolution tree

Evolution and HIV

- We often think of evolution as happening over geological time
- But the HIV virus evolved over decades from a disease that infected African primates to one that infects man
- Furthermore it evolves over months and years to evade our immune system
- Tracking these changes can be important in understanding and controlling the disease
- The virus evolves asexually so we can build an evolution tree

Evolution and HIV

- We often think of evolution as happening over geological time
- But the HIV virus evolved over decades from a disease that infected African primates to one that infects man
- Furthermore it evolves over months and years to evade our immune system
- Tracking these changes can be important in understanding and controlling the disease
- The virus evolves asexually so we can build an evolution tree

Evolution and HIV

- We often think of evolution as happening over geological time
- But the HIV virus evolved over decades from a disease that infected African primates to one that infects man
- Furthermore it evolves over months and years to evade our immune system
- Tracking these changes can be important in understanding and controlling the disease
- The virus evolves asexually so we can build an evolution tree

Evolution and HIV

- We often think of evolution as happening over geological time
- But the HIV virus evolved over decades from a disease that infected African primates to one that infects man
- Furthermore it evolves over months and years to evade our immune system
- Tracking these changes can be important in understanding and controlling the disease
- The virus evolves asexually so we can build an evolution tree

Evolution of Cancer

- Cells even in higher organisms predominantly reproduce asexually (they continuously divide)
- Over time the genome can get disrupted which if we are unlucky can grow into tumours
- Cancer cells accumulate mutations, some of which will die out others of which will replicate faster that other cells
- Understanding the development of cancer is often crucial to its treatment

Evolution of Cancer

- Cells even in higher organisms predominantly reproduce asexually (they continuously divide)
- Over time the genome can get disrupted which if we are unlucky can grow into tumours
- Cancer cells accumulate mutations, some of which will die out others of which will replicate faster that other cells
- Understanding the development of cancer is often crucial to its treatment

Evolution of Cancer

- Cells even in higher organisms predominantly reproduce asexually (they continuously divide)
- Over time the genome can get disrupted which if we are unlucky can grow into tumours
- Cancer cells accumulate mutations, some of which will die out others of which will replicate faster that other cells
- Understanding the development of cancer is often crucial to its treatment

Evolution of Cancer

- Cells even in higher organisms predominantly reproduce asexually (they continuously divide)
- Over time the genome can get disrupted which if we are unlucky can grow into tumours
- Cancer cells accumulate mutations, some of which will die out others of which will replicate faster that other cells
- Understanding the development of cancer is often crucial to its treatment

Orthologues and Paralogues

- Genes that diverge due to speciation are known as orthologues
- Genes that diverge due to gene duplication are known as paralogues
- If we are interested in understanding the relationship between species (e.g. pandas) we have to make sure we choose orthologues
- But if we are interested in the different types of haemoglobins then we have to look at paralogues

Orthologues and Paralogues

- Genes that diverge due to speciation are known as orthologues
- Genes that diverge due to gene duplication are known as paralogues
- If we are interested in understanding the relationship between species (e.g. pandas) we have to make sure we choose orthologues
- But if we are interested in the different types of haemoglobins then we have to look at paralogues

Orthologues and Paralogues

- Genes that diverge due to speciation are known as orthologues
- Genes that diverge due to gene duplication are known as paralogues
- If we are interested in understanding the relationship between species (e.g. pandas) we have to make sure we choose orthologues
- But if we are interested in the different types of haemoglobins then we have to look at paralogues

Orthologues and Paralogues

- Genes that diverge due to speciation are known as orthologues
- Genes that diverge due to gene duplication are known as paralogues
- If we are interested in understanding the relationship between species (e.g. pandas) we have to make sure we choose orthologues
- But if we are interested in the different types of haemoglobins then we have to look at paralogues

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Phylogenetic Tree of Globin Gene



Evolutionary history of the globin genes



© 2005 Encyclopædia Britannica, Inc.

Three Camps

- Although you would think that evolutionary trees should be pretty well understood, there is a huge debate about how to do this right
- There are three bitter rival camps in the field
  - ⋆ Evolutionary Taxonomy
  - ⋆ Phenetics or numerical taxonomy
  - ⋆ Cladistics
- If you are interested you can delve in the philosophical differences, but we will ignore them

Three Camps

- Although you would think that evolutionary trees should be pretty well understood, there is a huge debate about how to do this right
- There are three bitter rival camps in the field
  - ⋆ Evolutionary Taxonomy
  - ⋆ Phenetics or numerical taxonomy
  - ⋆ Cladistics
- If you are interested you can delve in the philosophical differences, but we will ignore them

Three Camps

- Although you would think that evolutionary trees should be pretty well understood, there is a huge debate about how to do this right
- There are three bitter rival camps in the field
  - ⋆ Evolutionary Taxonomy
  - ⋆ Phenetics or numerical taxonomy
  - ⋆ Cladistics
- If you are interested you can delve in the philosophical differences, but we will ignore them

Outline

1. Importance of Phylogeny
2. **Distance Based Trees**



Building Phylogenetic Trees

- Building evolutionary trees (also known as phylogenetic trees) based on some measure of evolutionary distance is scientifically important
- Traditionally morphological features where used, but today sequence edit distance is usually used
- There are often many plausible tree that can explain the data
- Efficient methods that find the best evolutionary trees still don't exist—current algorithms are compromises

Building Phylogenetic Trees

- Building evolutionary trees (also known as phylogenetic trees) based on some measure of evolutionary distance is scientifically important
- Traditionally morphological features where used, but today sequence edit distance is usually used
- There are often many plausible tree that can explain the data
- Efficient methods that find the best evolutionary trees still don't exist—current algorithms are compromises

Building Phylogenetic Trees

- Building evolutionary trees (also known as phylogenetic trees) based on some measure of evolutionary distance is scientifically important
- Traditionally morphological features where used, but today sequence edit distance is usually used
- There are often many plausible tree that can explain the data
- Efficient methods that find the best evolutionary trees still don't exist—current algorithms are compromises

Building Phylogenetic Trees

- Building evolutionary trees (also known as phylogenetic trees) based on some measure of evolutionary distance is scientifically important
- Traditionally morphological features where used, but today sequence edit distance is usually used
- There are often many plausible tree that can explain the data
- Efficient methods that find the best evolutionary trees still don't exist—current algorithms are compromises

The Molecular Clock

- It is a nice picture to see the accumulation of mutations acts as a clock telling us when species separated
- Unfortunately, mutations are random and strongly affected by selection pressure
- Also mutations are often very complex, involve a duplication of a stretch of DNA
- A lot of care is needed to use mutations as a molecular clock, including using as large sequences as possible

The Molecular Clock

- It is a nice picture to see the accumulation of mutations acts as a clock telling us when species separated
- Unfortunately, mutations are random and strongly affected by selection pressure
- Also mutations are often very complex, involve a duplication of a stretch of DNA
- A lot of care is needed to use mutations as a molecular clock, including using as large sequences as possible

The Molecular Clock

- It is a nice picture to see the accumulation of mutations acts as a clock telling us when species separated
- Unfortunately, mutations are random and strongly affected by selection pressure
- Also mutations are often very complex, involve a duplication of a stretch of DNA
- A lot of care is needed to use mutations as a molecular clock, including using as large sequences as possible

The Molecular Clock

- It is a nice picture to see the accumulation of mutations acts as a clock telling us when species separated
- Unfortunately, mutations are random and strongly affected by selection pressure
- Also mutations are often very complex, involve a duplication of a stretch of DNA
- A lot of care is needed to use mutations as a molecular clock, including using as large sequences as possible

Two Types of Trees

- There are two main types of evolutionary trees
  - ⋆ **Distance-based trees**, where we have a table of "distances" between species and we wish to find a tree explaining these distances
  - ⋆ **Sequence based trees**, where we are given a set of sequences and we want to find a tree with a minimum number of mutations per link (maximum parsimony) that explains the data

Two Types of Trees

- There are two main types of evolutionary trees
  - ⋆ **Distance-based trees**, where we have a table of "distances" between species and we wish to find a tree explaining these distances
  - ⋆ **Sequence based trees**, where we are given a set of sequences and we want to find a tree with a minimum number of mutations per link (maximum parsimony) that explains the data

Ultrametric Trees

- We consider binary trees

- If we had a perfect molecular clock then evolutionary trees to existing species will have an ultrametric structure



	A	B	C	D	E
A	0	3	5	9	9
B	3	0	5	9	9
C	5	5	0	9	9
D	9	9	9	0	6
E	9	9	9	6	0

- For any three nodes i, j and k with distances d_{ij}, d_{ik} and d_{jk} two distances will be identical, and one will be smaller than these

Ultrametric Trees

- We consider binary trees

- If we had a perfect molecular clock then evolutionary trees to existing species will have an ultrametric structure



	A	B	C	D	E
A	0	3	5	9	9
B	3	0	5	9	9
C	5	5	0	9	9
D	9	9	9	0	6
E	9	9	9	6	0

- For any three nodes i, j and k with distances d_{ij}, d_{ik} and d_{jk} two distances will be identical, and one will be smaller than these

Ultrametric Trees

- We consider binary trees

- If we had a perfect molecular clock then evolutionary trees to existing species will have an ultrametric structure



	A	B	C	D	E
A	0	3	5	9	9
B	3	0	5	9	9
C	5	5	0	9	9
D	9	9	9	0	6
E	9	9	9	6	0

- For any three nodes i, j and k with distances d_{ij}, d_{ik} and d_{jk} two distances will be identical, and one will be smaller than these

Real Trees

- Because we cannot accurately measure divergence time, in practice it is very unlikely that the table of distances can be modelled with an ultrametric tree
- We can however build an ultrametric tree through a clustering algorithm known as **Unweighted Pair Group Method using Arithmetic averaging** UPGMA
- This is an easy algorithm where we build a tree from the leaves upwards
- We select the two closest subtrees remove all the nodes and replace the two joined subtree with a new subtree

Real Trees

- Because we cannot accurately measure divergence time, in practice it is very unlikely that the table of distances can be modelled with an ultrametric tree
- We can however build an ultrametric tree through a clustering algorithm known as **Unweighted Pair Group Method using Arithmetic averaging** UPGMA
- This is an easy algorithm where we build a tree from the leaves upwards
- We select the two closest subtrees remove all the nodes and replace the two joined subtree with a new subtree

Real Trees

- Because we cannot accurately measure divergence time, in practice it is very unlikely that the table of distances can be modelled with an ultrametric tree
- We can however build an ultrametric tree through a clustering algorithm known as **Unweighted Pair Group Method using Arithmetic averaging** UPGMA
- This is an easy algorithm where we build a tree from the leaves upwards
- We select the two closest subtrees remove all the nodes and replace the two joined subtree with a new subtree

Real Trees

- Because we cannot accurately measure divergence time, in practice it is very unlikely that the table of distances can be modelled with an ultrametric tree
- We can however build an ultrametric tree through a clustering algorithm known as **Unweighted Pair Group Method using Arithmetic averaging** UPGMA
- This is an easy algorithm where we build a tree from the leaves upwards
- We select the two closest subtrees remove all the nodes and replace the two joined subtree with a new subtree

UPGMA



UPGMA



UPGMA



UPGMA



UPGMA



UPGMA



UPGMA





Distance Calculation for UPGMA

- The UPGMA algorithm starts from a table of distances

- The distance between two clusters $\mathcal{C}_k$ and $\mathcal{C}_l$ is given by

$$d_{kl} = \frac{1}{|\mathcal{C}_k|\,|\mathcal{C}_l|} \sum_{\substack{p \in \mathcal{C}_k \\ q \in \mathcal{C}_l}} d_{pq}$$

- If we join two clusters so that $\mathcal{C}_k = \mathcal{C}_i \cup \mathcal{C}_j$ then the distance to a cluster $\mathcal{C}_l$ is

$$d_{kl} = \frac{d_{il}\,|\mathcal{C}_i| + d_{jl}\,|\mathcal{C}_j|}{|\mathcal{C}_i| + |\mathcal{C}_j|}$$

- If the distances are not ultrametric then UPGMA can give poor trees

Distance Calculation for UPGMA

- The UPGMA algorithm starts from a table of distances

- The distance between two clusters $\mathcal{C}_k$ and $\mathcal{C}_l$ is given by

$$d_{kl} = \frac{1}{|\mathcal{C}_k|\,|\mathcal{C}_l|} \sum_{\substack{p \in \mathcal{C}_k \\ q \in \mathcal{C}_l}} d_{pq}$$

- If we join two clusters so that $\mathcal{C}_k = \mathcal{C}_i \cup \mathcal{C}_j$ then the distance to a cluster $\mathcal{C}_l$ is

$$d_{kl} = \frac{d_{il}\,|\mathcal{C}_i| + d_{jl}\,|\mathcal{C}_j|}{|\mathcal{C}_i| + |\mathcal{C}_j|}$$

- If the distances are not ultrametric then UPGMA can give poor trees

Distance Calculation for UPGMA

- The UPGMA algorithm starts from a table of distances

- The distance between two clusters $\mathcal{C}_k$ and $\mathcal{C}_l$ is given by

$$d_{kl} = \frac{1}{|\mathcal{C}_k|\,|\mathcal{C}_l|} \sum_{\substack{p \in \mathcal{C}_k \\ q \in \mathcal{C}_l}} d_{pq}$$

- If we join two clusters so that $\mathcal{C}_k = \mathcal{C}_i \cup \mathcal{C}_j$ then the distance to a cluster $\mathcal{C}_l$ is

$$d_{kl} = \frac{d_{il}\,|\mathcal{C}_i| + d_{jl}\,|\mathcal{C}_j|}{|\mathcal{C}_i| + |\mathcal{C}_j|}$$

- If the distances are not ultrametric then UPGMA can give poor trees

Distance Calculation for UPGMA

- The UPGMA algorithm starts from a table of distances

- The distance between two clusters $\mathcal{C}_k$ and $\mathcal{C}_l$ is given by

$$d_{kl} = \frac{1}{|\mathcal{C}_k|\,|\mathcal{C}_l|} \sum_{\substack{p \in \mathcal{C}_k \\ q \in \mathcal{C}_l}} d_{pq}$$

- If we join two clusters so that $\mathcal{C}_k = \mathcal{C}_i \cup \mathcal{C}_j$ then the distance to a cluster $\mathcal{C}_l$ is

$$d_{kl} = \frac{d_{il}\,|\mathcal{C}_i| + d_{jl}\,|\mathcal{C}_j|}{|\mathcal{C}_i| + |\mathcal{C}_j|}$$

- If the distances are not ultrametric then UPGMA can give poor trees

Weakness of UPGMA

- Note that organism 5 and 7 are quite close, but UPGMA distorts this

Weakness of UPGMA

- Note that organism 5 and 7 are quite close, but UPGMA distorts this



Weakness of UPGMA

- Note that organism 5 and 7 are quite close, but UPGMA distorts this



Weakness of UPGMA

- Note that organism 5 and 7 are quite close, but UPGMA distorts this



Weakness of UPGMA

- Note that organism 5 and 7 are quite close, but UPGMA distorts this



Weakness of UPGMA

- Note that organism 5 and 7 are quite close, but UPGMA distorts this



Weakness of UPGMA

- Note that organism 5 and 7 are quite close, but UPGMA distorts this



Reflection on Ultrametricity

- UPGMA imposes an ultrametric structure on the derived evolutionary trees
- If mutations accumulated at a constant rate then evolutionary trees should have an ultrametric structure
- Unfortunately mutations experience selection and in some cases they are very non-trivial (genome copy mutations)
- There are stretches of "junk" DNA where mutations appear to accumulate at a reasonable constant rate, but these are not well studied or stable regions

Reflection on Ultrametricity

- UPGMA imposes an ultrametric structure on the derived evolutionary trees
- If mutations accumulated at a constant rate then evolutionary trees should have an ultrametric structure
- Unfortunately mutations experience selection and in some cases they are very non-trivial (genome copy mutations)
- There are stretches of "junk" DNA where mutations appear to accumulate at a reasonable constant rate, but these are not well studied or stable regions

Reflection on Ultrametricity

- UPGMA imposes an ultrametric structure on the derived evolutionary trees
- If mutations accumulated at a constant rate then evolutionary trees should have an ultrametric structure
- Unfortunately mutations experience selection and in some cases they are very non-trivial (genome copy mutations)
- There are stretches of "junk" DNA where mutations appear to accumulate at a reasonable constant rate, but these are not well studied or stable regions

Reflection on Ultrametricity

- UPGMA imposes an ultrametric structure on the derived evolutionary trees
- If mutations accumulated at a constant rate then evolutionary trees should have an ultrametric structure
- Unfortunately mutations experience selection and in some cases they are very non-trivial (genome copy mutations)
- There are stretches of "junk" DNA where mutations appear to accumulate at a reasonable constant rate, but these are not well studied or stable regions

Summary

- Evolutionary trees provide very important scientific insights
- Building evolutionary trees is non-trivial
- If the distances are approximately ultrametric then UPGMA will generate a good tree
- Need to look at other ways of building evolutionary trees

Summary

- Evolutionary trees provide very important scientific insights
- Building evolutionary trees is non-trivial
- If the distances are approximately ultrametric then UPGMA will generate a good tree
- Need to look at other ways of building evolutionary trees

Summary

- Evolutionary trees provide very important scientific insights
- Building evolutionary trees is non-trivial
- If the distances are approximately ultrametric then UPGMA will generate a good tree
- Need to look at other ways of building evolutionary trees

Summary

- Evolutionary trees provide very important scientific insights
- Building evolutionary trees is non-trivial
- If the distances are approximately ultrametric then UPGMA will generate a good tree
- Need to look at other ways of building evolutionary trees