


# Advanced Machine Learning


## *Differential Calculus*



*Differentiation, product and chain rules, vectors and matrices*

# Outline

1. **Why Calculus?**
2. Differentiation
3. Vector and Matrix Calculus



# Why Calculus?

- Calculus is a fundamental tool of mathematical analysis
- In machine learning differentiation is fundamental tool in optimisation
- Integration is an essential tool in taking expectations over continuous distributions
- Both differentiation and integration crop up elsewhere

# Why Calculus?

- Calculus is a fundamental tool of mathematical analysis
- In machine learning differentiation is fundamental tool in optimisation
- Integration is an essential tool in taking expectations over continuous distributions
- Both differentiation and integration crop up elsewhere

# Why Calculus?

- Calculus is a fundamental tool of mathematical analysis
- In machine learning differentiation is fundamental tool in optimisation
- Integration is an essential tool in taking expectations over continuous distributions
- Both differentiation and integration crop up elsewhere

# Why Calculus?

- Calculus is a fundamental tool of mathematical analysis
- In machine learning differentiation is fundamental tool in optimisation
- Integration is an essential tool in taking expectations over continuous distributions
- Both differentiation and integration crop up elsewhere

# Why Calculus?

- Calculus is a fundamental tool of mathematical analysis
- In machine learning differentiation is fundamental tool in optimisation
- Integration is an essential tool in taking expectations over continuous distributions
- Both differentiation and integration crop up elsewhere
- This material will not be examined explicitly

# Why Calculus?

- Calculus is a fundamental tool of mathematical analysis
- In machine learning differentiation is fundamental tool in optimisation
- Integration is an essential tool in taking expectations over continuous distributions
- Both differentiation and integration crop up elsewhere
- This material will not be examined explicitly, but I assume elsewhere that you can do calculus



# Back to Basics

- You have all done A-level maths so should be familiar with the rules of calculus
- But, it is easy to forget the rules and sometimes we use quite sophisticated tricks
- Although the sophisticated tricks really speed up calculations, it pays to be able to understand where these tricks come from

# Back to Basics


- You have all done A-level maths so should be familiar with the rules of calculus
- But, it is easy to forget the rules and sometimes we use quite sophisticated tricks
- Although the sophisticated tricks really speed up calculations, it pays to be able to understand where these tricks come from

# Back to Basics


- You have all done A-level maths so should be familiar with the rules of calculus
- But, it is easy to forget the rules and sometimes we use quite sophisticated tricks
- Although the sophisticated tricks really speed up calculations, it pays to be able to understand where these tricks come from

# Outline


1. Why Calculus?
2. **Differentiation**
3. Vector and Matrix Calculus




# Differentiation




# Differentiation



# Differentiation




# Differentiation





# Differentiation



# Polynomials

- $f(x) = x^2$

$$\frac{dx^2}{dx} = \lim_{\epsilon \rightarrow 0} \frac{(x + \epsilon)^2 - x^2}{\epsilon}$$

# Polynomials

- $f(x) = x^2$

$$\frac{dx^2}{dx} = \lim_{\epsilon \rightarrow 0} \frac{(x + \epsilon)^2 - x^2}{\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{(x^2 + 2\epsilon x + \epsilon^2) - x^2}{\epsilon}$$

# Polynomials

- $f(x) = x^2$

$$\begin{aligned}\frac{dx^2}{dx} &= \lim_{\epsilon \rightarrow 0} \frac{(x + \epsilon)^2 - x^2}{\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{(x^2 + 2\epsilon x + \epsilon^2) - x^2}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} 2x + \epsilon\end{aligned}$$

# Polynomials

- $f(x) = x^2$

$$\begin{aligned}\frac{dx^2}{dx} &= \lim_{\epsilon \rightarrow 0} \frac{(x + \epsilon)^2 - x^2}{\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{(x^2 + 2\epsilon x + \epsilon^2) - x^2}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} 2x + \epsilon = 2x\end{aligned}$$

# Polynomials

- $f(x) = x^2$

$$\begin{aligned}\frac{dx^2}{dx} &= \lim_{\epsilon \rightarrow 0} \frac{(x + \epsilon)^2 - x^2}{\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{(x^2 + 2\epsilon x + \epsilon^2) - x^2}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} 2x + \epsilon = 2x\end{aligned}$$

- $(x + \epsilon)^n = (x + \epsilon)(x + \epsilon) \cdots (x + \epsilon)$

# Polynomials

- $f(x) = x^2$

$$\begin{aligned}\frac{dx^2}{dx} &= \lim_{\epsilon \rightarrow 0} \frac{(x + \epsilon)^2 - x^2}{\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{(x^2 + 2\epsilon x + \epsilon^2) - x^2}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} 2x + \epsilon = 2x\end{aligned}$$

- $(x + \epsilon)^n = (x + \epsilon)(x + \epsilon) \cdots (x + \epsilon) = x^n + n\epsilon x^{n-1} + O(\epsilon^2)$

# Polynomials

- $f(x) = x^2$

$$\begin{aligned}\frac{dx^2}{dx} &= \lim_{\epsilon \rightarrow 0} \frac{(x + \epsilon)^2 - x^2}{\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{(x^2 + 2\epsilon x + \epsilon^2) - x^2}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} 2x + \epsilon = 2x\end{aligned}$$

- $(x + \epsilon)^n = (x + \epsilon)(x + \epsilon) \cdots (x + \epsilon) = x^n + n\epsilon x^{n-1} + O(\epsilon^2)$

$$\frac{dx^n}{dx} = \lim_{\epsilon \rightarrow 0} \frac{(x + \epsilon)^n - x^n}{\epsilon}$$



# Polynomials

- $f(x) = x^2$

$$\begin{aligned}\frac{dx^2}{dx} &= \lim_{\epsilon \rightarrow 0} \frac{(x + \epsilon)^2 - x^2}{\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{(x^2 + 2\epsilon x + \epsilon^2) - x^2}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} 2x + \epsilon = 2x\end{aligned}$$

- $(x + \epsilon)^n = (x + \epsilon)(x + \epsilon) \cdots (x + \epsilon) = x^n + n\epsilon x^{n-1} + O(\epsilon^2)$

$$\frac{dx^n}{dx} = \lim_{\epsilon \rightarrow 0} \frac{(x + \epsilon)^n - x^n}{\epsilon} = \lim_{\epsilon \rightarrow 0} nx^{n-1} + O(\epsilon)$$

# Polynomials

- $f(x) = x^2$

$$\begin{aligned}\frac{dx^2}{dx} &= \lim_{\epsilon \rightarrow 0} \frac{(x + \epsilon)^2 - x^2}{\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{(x^2 + 2\epsilon x + \epsilon^2) - x^2}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} 2x + \epsilon = 2x\end{aligned}$$

- $(x + \epsilon)^n = (x + \epsilon)(x + \epsilon) \cdots (x + \epsilon) = x^n + n\epsilon x^{n-1} + O(\epsilon^2)$

$$\frac{dx^n}{dx} = \lim_{\epsilon \rightarrow 0} \frac{(x + \epsilon)^n - x^n}{\epsilon} = \lim_{\epsilon \rightarrow 0} nx^{n-1} + O(\epsilon) = nx^{n-1}$$

# Linearity of derivatives

- Note that  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$  (from the definition of  $f'(x)$ )

$$\frac{d(a f(x) + b g(x))}{dx} = \lim_{\epsilon \rightarrow 0} \frac{(a f(x + \epsilon) + b g(x + \epsilon)) - (a f(x) + b g(x))}{\epsilon}$$

# Linearity of derivatives

- Note that  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$  (from the definition of  $f'(x)$ )

$$\frac{d(a f(x) + b g(x))}{dx} = \lim_{\epsilon \rightarrow 0} \frac{(a f(x + \epsilon) + b g(x + \epsilon)) - (a f(x) + b g(x))}{\epsilon}$$

# Linearity of derivatives

- Note that  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$  (from the definition of  $f'(x)$ )

$$\begin{aligned} \frac{d(a f(x) + b g(x))}{dx} &= \lim_{\epsilon \rightarrow 0} \frac{(a f(x + \epsilon) + b g(x + \epsilon)) - (a f(x) + b g(x))}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} \frac{a \epsilon f'(x) + b \epsilon g'(x) + O(\epsilon^2)}{\epsilon} \end{aligned}$$

# Linearity of derivatives

- Note that  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$  (from the definition of  $f'(x)$ )

$$\begin{aligned}\frac{d(a f(x) + b g(x))}{dx} &= \lim_{\epsilon \rightarrow 0} \frac{(a f(x + \epsilon) + b g(x + \epsilon)) - (a f(x) + b g(x))}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} \frac{a \epsilon f'(x) + b \epsilon g'(x) + O(\epsilon^2)}{\epsilon} \\ &= a f'(x) + b g'(x)\end{aligned}$$


# Linearity of derivatives

- Note that  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$  (from the definition of  $f'(x)$ )

$$\begin{aligned}\frac{d(a f(x) + b g(x))}{dx} &= \lim_{\epsilon \rightarrow 0} \frac{(a f(x + \epsilon) + b g(x + \epsilon)) - (a f(x) + b g(x))}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} \frac{a \epsilon f'(x) + b \epsilon g'(x) + O(\epsilon^2)}{\epsilon} \\ &= a f'(x) + b g'(x)\end{aligned}$$


- **Differentiation is a linear operation!**

# Linearity in Pictures







# Linearity in Pictures




# Linearity in Pictures




# Linearity in Pictures




# Linearity in Pictures




# Linearity in Pictures




# Linearity in Pictures



# Linearity in Pictures



# Linearity in Pictures





# Product Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- If  $h(x) = f(x)g(x)$

# Product Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- If  $h(x) = f(x)g(x)$

$$h'(x) = \lim_{\epsilon \rightarrow 0} \frac{f(x + \epsilon)g(x + \epsilon) - f(x)g(x)}{\epsilon}$$

# Product Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- If  $h(x) = f(x)g(x)$

$$\begin{aligned} h'(x) &= \lim_{\epsilon \rightarrow 0} \frac{f(x + \epsilon)g(x + \epsilon) - f(x)g(x)}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} \frac{(f(x) + \epsilon f'(x) + O(\epsilon^2))(g(x) + \epsilon g'(x) + O(\epsilon^2)) - f(x)g(x)}{\epsilon} \end{aligned}$$

# Product Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- If  $h(x) = f(x)g(x)$

$$\begin{aligned} h'(x) &= \lim_{\epsilon \rightarrow 0} \frac{f(x + \epsilon)g(x + \epsilon) - f(x)g(x)}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} \frac{(f(x) + \epsilon f'(x) + O(\epsilon^2))(g(x) + \epsilon g'(x) + O(\epsilon^2)) - f(x)g(x)}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} \frac{\epsilon(f'(x)g(x) + f(x)g'(x)) + O(\epsilon^2)}{\epsilon} \end{aligned}$$

# Product Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- If  $h(x) = f(x)g(x)$

$$\begin{aligned} h'(x) &= \lim_{\epsilon \rightarrow 0} \frac{f(x + \epsilon)g(x + \epsilon) - f(x)g(x)}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} \frac{(f(x) + \epsilon f'(x) + O(\epsilon^2))(g(x) + \epsilon g'(x) + O(\epsilon^2)) - f(x)g(x)}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} \frac{\epsilon(f'(x)g(x) + f(x)g'(x)) + O(\epsilon^2)}{\epsilon} = f'(x)g(x) + f(x)g'(x) \end{aligned}$$

# Product Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- If  $h(x) = f(x)g(x)$

$$\begin{aligned} h'(x) &= \lim_{\epsilon \rightarrow 0} \frac{f(x + \epsilon)g(x + \epsilon) - f(x)g(x)}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} \frac{(f(x) + \epsilon f'(x) + O(\epsilon^2))(g(x) + \epsilon g'(x) + O(\epsilon^2)) - f(x)g(x)}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} \frac{\epsilon(f'(x)g(x) + f(x)g'(x)) + O(\epsilon^2)}{\epsilon} = f'(x)g(x) + f(x)g'(x) \end{aligned}$$

- This is the **product rule**

# Chain Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- Let  $h(x) = f(g(x))$
- Then

$$h(x + \epsilon) = f(g(x + \epsilon))$$

# Chain Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- Let  $h(x) = f(g(x))$
- Then

$$h(x + \epsilon) = f(g(x + \epsilon))$$



# Chain Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- Let  $h(x) = f(g(x))$
- Then

$$h(x + \epsilon) = f(g(x + \epsilon)) = f(g(x) + \epsilon g'(x) + O(\epsilon^2))$$

# Chain Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- Let  $h(x) = f(g(x))$
- Then

$$\begin{aligned} h(x + \epsilon) &= f(g(x + \epsilon)) = f(g(x) + \epsilon g'(x) + O(\epsilon^2)) \\ &= f(g(x)) + \epsilon g'(x) f'(g(x)) + O(\epsilon^2) \end{aligned}$$

# Chain Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- Let  $h(x) = f(g(x))$
- Then

$$\begin{aligned} h(x + \epsilon) &= f(g(x + \epsilon)) = f(g(x) + \epsilon g'(x) + O(\epsilon^2)) \\ &= f(g(x)) + \epsilon g'(x) f'(g(x)) + O(\epsilon^2) \end{aligned}$$

- Thus

$$h'(x) = \lim_{\epsilon \rightarrow 0} \frac{h(x + \epsilon) - h(x)}{\epsilon} = g'(x) f'(g(x))$$

# Chain Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- Let  $h(x) = f(g(x))$
- Then

$$\begin{aligned} h(x + \epsilon) &= f(g(x + \epsilon)) = f(g(x) + \epsilon g'(x) + O(\epsilon^2)) \\ &= f(g(x)) + \epsilon g'(x) f'(g(x)) + O(\epsilon^2) \end{aligned}$$

- Thus

$$h'(x) = \lim_{\epsilon \rightarrow 0} \frac{h(x + \epsilon) - h(x)}{\epsilon} = g'(x) f'(g(x))$$

- This is the famous **chain rule**

# Chain Rule

- Recall  $f(x + \epsilon) = f(x) + \epsilon f'(x) + O(\epsilon^2)$
- Let  $h(x) = f(g(x))$
- Then

$$\begin{aligned} h(x + \epsilon) &= f(g(x + \epsilon)) = f(g(x) + \epsilon g'(x) + O(\epsilon^2)) \\ &= f(g(x)) + \epsilon g'(x) f'(g(x)) + O(\epsilon^2) \end{aligned}$$

- Thus

$$h'(x) = \lim_{\epsilon \rightarrow 0} \frac{h(x + \epsilon) - h(x)}{\epsilon} = g'(x) f'(g(x))$$

- This is the famous **chain rule**. Together with the product rule it means you can differentiate almost everything

# More on chain rules

- We can also write the chain rule as

$$\frac{df(g(x))}{dx} = \frac{df(g)}{dg} \frac{dg(x)}{dx}$$

- Sometimes this is neater or easier to remember

$$\frac{de^{\cos(x^2)}}{dx} = \frac{de^{\cos(x^2)}}{d\cos(x^2)} \frac{d\cos(x^2)}{dx^2} \frac{dx^2}{dx}$$

# More on chain rules

- We can also write the chain rule as

$$\frac{df(g(x))}{dx} = \frac{df(g)}{dg} \frac{dg(x)}{dx}$$

- Sometimes this is neater or easier to remember

$$\frac{de^{\cos(x^2)}}{dx} = \frac{de^{\cos(x^2)}}{d\cos(x^2)} \frac{d\cos(x^2)}{dx^2} \frac{dx^2}{dx}$$

# More on chain rules

- We can also write the chain rule as

$$\frac{df(g(x))}{dx} = \frac{df(g)}{dg} \frac{dg(x)}{dx}$$

- Sometimes this is neater or easier to remember

$$\begin{aligned} \frac{de^{\cos(x^2)}}{dx} &= \frac{de^{\cos(x^2)}}{d\cos(x^2)} \frac{d\cos(x^2)}{dx^2} \frac{dx^2}{dx} \\ &= e^{\cos(x^2)} (-\sin(x^2)) 2x \end{aligned}$$



# More on chain rules

- We can also write the chain rule as

$$\frac{df(g(x))}{dx} = \frac{df(g)}{dg} \frac{dg(x)}{dx}$$

- Sometimes this is neater or easier to remember

$$\begin{aligned} \frac{de^{\cos(x^2)}}{dx} &= \frac{de^{\cos(x^2)}}{d\cos(x^2)} \frac{d\cos(x^2)}{dx^2} \frac{dx^2}{dx} \\ &= e^{\cos(x^2)} (-\sin(x^2)) 2x \\ &= -2x \sin(x^2) e^{\cos(x^2)} \end{aligned}$$

# Inverse functions

- Suppose  $g(y) = f^{-1}(y)$  is the inverse of  $f(x)$  in the sense that  $g(f(x)) = f^{-1}(f(x)) = x$
- Using the chain rule

$$\frac{dg(f(x))}{dx} = f'(x)g'(f(x))$$

- So  $g'(f(x)) = 1/f'(x)$
- Writing  $y = f(x)$  so that  $x = f^{-1}(y) = g(y)$  we find  $g'(y) = 1/f'(g(y))$  that is

$$\frac{dg(y)}{dy} = \frac{1}{f'(g(y))}$$

# Inverse functions

- Suppose  $g(y) = f^{-1}(y)$  is the inverse of  $f(x)$  in the sense that  $g(f(x)) = f^{-1}(f(x)) = x$
- Using the chain rule

$$\frac{dg(f(x))}{dx} = f'(x)g'(f(x))$$

- So  $g'(f(x)) = 1/f'(x)$
- Writing  $y = f(x)$  so that  $x = f^{-1}(y) = g(y)$  we find  $g'(y) = 1/f'(g(y))$  that is

$$\frac{dg(y)}{dy} = \frac{1}{f'(g(y))}$$

# Inverse functions

- Suppose  $g(y) = f^{-1}(y)$  is the inverse of  $f(x)$  in the sense that  $g(f(x)) = f^{-1}(f(x)) = x$
- Using the chain rule

$$\frac{dg(f(x))}{dx} = f'(x)g'(f(x)) = 1$$

since  $g(f(x)) = x$

- So  $g'(f(x)) = 1/f'(x)$
- Writing  $y = f(x)$  so that  $x = f^{-1}(y) = g(y)$  we find  $g'(y) = 1/f'(g(y))$  that is

$$\frac{dg(y)}{dy} = \frac{1}{f'(g(y))}$$

# Inverse functions

- Suppose  $g(y) = f^{-1}(y)$  is the inverse of  $f(x)$  in the sense that  $g(f(x)) = f^{-1}(f(x)) = x$
- Using the chain rule

$$\frac{dg(f(x))}{dx} = f'(x)g'(f(x)) = 1$$

since  $g(f(x)) = x$

- So  $g'(f(x)) = 1/f'(x)$
- Writing  $y = f(x)$  so that  $x = f^{-1}(y) = g(y)$  we find  $g'(y) = 1/f'(g(y))$  that is

$$\frac{dg(y)}{dy} = \frac{1}{f'(g(y))}$$

# Inverse functions

- Suppose  $g(y) = f^{-1}(y)$  is the inverse of  $f(x)$  in the sense that  $g(f(x)) = f^{-1}(f(x)) = x$
- Using the chain rule

$$\frac{dg(f(x))}{dx} = f'(x)g'(f(x)) = 1$$

since  $g(f(x)) = x$

- So  $g'(f(x)) = 1/f'(x)$
- Writing  $y = f(x)$  so that  $x = f^{-1}(y) = g(y)$  we find  $g'(y) = 1/f'(g(y))$  that is

$$\frac{dg(y)}{dy} = \frac{1}{f'(g(y))}$$

# Inverse functions

- Suppose  $g(y) = f^{-1}(y)$  is the inverse of  $f(x)$  in the sense that  $g(f(x)) = f^{-1}(f(x)) = x$
- Using the chain rule

$$\frac{dg(f(x))}{dx} = f'(x)g'(f(x)) = 1$$

since  $g(f(x)) = x$

- So  $g'(f(x)) = 1/f'(x)$
- Writing  $y = f(x)$  so that  $x = f^{-1}(y) = g(y)$  we find  $g'(y) = 1/f'(g(y))$  that is

$$\frac{dg(y)}{dy} = \frac{1}{f'(g(y))}$$

$$\frac{df^{-1}(y)}{dy} = \frac{1}{f'(f^{-1}(y))}$$

# Exponentials


- Note that  $a^{b+c} = a^b a^c$  (that is we multiply  $a$  together  $b + c$  times)



# Exponentials

- Note that  $a^{b+c} = a^b a^c$  (that is we multiply  $a$  together  $b + c$  times)


- Now  $e^\epsilon \approx (1 + \epsilon)$



# Exponentials

- Note that  $a^{b+c} = a^b a^c$  (that is we multiply  $a$  together  $b + c$  times)

- Now  $e^\epsilon \approx (1 + \epsilon)$




- But  $e^{x+\epsilon} = e^x e^\epsilon = e^x (1 + \epsilon + O(\epsilon^2))$

# Exponentials

- Note that  $a^{b+c} = a^b a^c$  (that is we multiply  $a$  together  $b + c$  times)

- Now  $e^\epsilon \approx (1 + \epsilon)$




- But  $e^{x+\epsilon} = e^x e^\epsilon = e^x (1 + \epsilon + O(\epsilon^2)) = e^x + \epsilon e^x + O(\epsilon^2)$

# Exponentials

- Note that  $a^{b+c} = a^b a^c$  (that is we multiply  $a$  together  $b + c$  times)

- Now  $e^\epsilon \approx (1 + \epsilon)$




- But  $e^{x+\epsilon} = e^x e^\epsilon = e^x (1 + \epsilon + O(\epsilon^2)) = e^x + \epsilon e^x + O(\epsilon^2)$

$$\frac{de^x}{dx} = \lim_{\epsilon \rightarrow 0} \frac{e^{x+\epsilon} - e^x}{\epsilon}$$

# Exponentials

- Note that  $a^{b+c} = a^b a^c$  (that is we multiply  $a$  together  $b + c$  times)

- Now  $e^\epsilon \approx (1 + \epsilon)$



- But  $e^{x+\epsilon} = e^x e^\epsilon = e^x (1 + \epsilon + O(\epsilon^2)) = e^x + \epsilon e^x + O(\epsilon^2)$

$$\frac{de^x}{dx} = \lim_{\epsilon \rightarrow 0} \frac{e^{x+\epsilon} - e^x}{\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{\epsilon e^x + O(\epsilon^2)}{\epsilon} = e^x$$

# Functions of Exponentials

- What about  $f(x) = e^{cx}$

$$\frac{de^{cx}}{dx} = \frac{de^{cx}}{dcx} \frac{dcx}{dx}$$

- More generally using the chain rule

$$\frac{de^{g(x)}}{dx} = g'(x)e^{g(x)}$$

# Functions of Exponentials

- What about  $f(x) = e^{cx}$

$$\frac{de^{cx}}{dx} = \frac{de^{cx}}{dcx} \frac{dcx}{dx} = ce^{cx}$$

- More generally using the chain rule

$$\frac{de^{g(x)}}{dx} = g'(x)e^{g(x)}$$

# Functions of Exponentials

- What about  $f(x) = e^{cx}$

$$\frac{de^{cx}}{dx} = \frac{de^{cx}}{dcx} \frac{dcx}{dx} = ce^{cx}$$

- More generally using the chain rule

$$\frac{de^{g(x)}}{dx} = g'(x)e^{g(x)}$$



# Functions of Exponentials

- What about  $f(x) = e^{cx}$

$$\frac{de^{cx}}{dx} = \frac{de^{cx}}{dcx} \frac{dcx}{dx} = ce^{cx}$$

- More generally using the chain rule

$$\frac{de^{g(x)}}{dx} = g'(x)e^{g(x)}$$

- Also  $a^{bc} = (a^b)^c$  (that is we multiply  $a$  together  $b \times c$  times)

# Functions of Exponentials

- What about  $f(x) = e^{cx}$

$$\frac{de^{cx}}{dx} = \frac{de^{cx}}{dcx} \frac{dcx}{dx} = ce^{cx}$$

- More generally using the chain rule

$$\frac{de^{g(x)}}{dx} = g'(x)e^{g(x)}$$

- Also  $a^{bc} = (a^b)^c$  (that is we multiply  $a$  together  $b \times c$  times)

$$\frac{da^x}{dx} = \frac{d(e^{\ln(a)})^x}{dx}$$

# Functions of Exponentials

- What about  $f(x) = e^{cx}$

$$\frac{de^{cx}}{dx} = \frac{de^{cx}}{dcx} \frac{dcx}{dx} = ce^{cx}$$

- More generally using the chain rule

$$\frac{de^{g(x)}}{dx} = g'(x)e^{g(x)}$$

- Also  $a^{bc} = (a^b)^c$  (that is we multiply  $a$  together  $b \times c$  times)

$$\frac{da^x}{dx} = \frac{d(e^{\ln(a)})^x}{dx} = \frac{de^{\ln(a)x}}{dx}$$

# Functions of Exponentials

- What about  $f(x) = e^{cx}$

$$\frac{de^{cx}}{dx} = \frac{de^{cx}}{dcx} \frac{dcx}{dx} = ce^{cx}$$

- More generally using the chain rule

$$\frac{de^{g(x)}}{dx} = g'(x)e^{g(x)}$$

- Also  $a^{bc} = (a^b)^c$  (that is we multiply  $a$  together  $b \times c$  times)

$$\frac{da^x}{dx} = \frac{d(e^{\ln(a)})^x}{dx} = \frac{de^{\ln(a)x}}{dx} = \ln(a)e^{\ln(a)x}$$

# Functions of Exponentials

- What about  $f(x) = e^{cx}$

$$\frac{de^{cx}}{dx} = \frac{de^{cx}}{dcx} \frac{dcx}{dx} = ce^{cx}$$

- More generally using the chain rule

$$\frac{de^{g(x)}}{dx} = g'(x)e^{g(x)}$$

- Also  $a^{bc} = (a^b)^c$  (that is we multiply  $a$  together  $b \times c$  times)

$$\frac{da^x}{dx} = \frac{d(e^{\ln(a)})^x}{dx} = \frac{de^{\ln(a)x}}{dx} = \ln(a)e^{\ln(a)x} = \ln(a)a^x$$

# Natural Logarithms

- The natural logarithm is defined as the inverse of  $e^x$

$$\ln(e^x) = x \qquad e^{\ln(y)} = y$$

- Recall that if  $g(y) = f^{-1}(y)$  then  $g'(y) = 1/f'(g(y))$
- Consider  $g(y) = \ln(y)$  and  $f(x) = e^x$  (with  $f'(x) = e^x$ )

$$\frac{d\ln(y)}{dy} = \frac{1}{e^{\ln(y)}}$$

# Natural Logarithms

- The natural logarithm is defined as the inverse of  $e^x$

$$\ln(e^x) = x \qquad e^{\ln(y)} = y$$

- Recall that if  $g(y) = f^{-1}(y)$  then  $g'(y) = 1/f'(g(y))$
- Consider  $g(y) = \ln(y)$  and  $f(x) = e^x$  (with  $f'(x) = e^x$ )

$$\frac{d\ln(y)}{dy} = \frac{1}{e^{\ln(y)}}$$

# Natural Logarithms

- The natural logarithm is defined as the inverse of  $e^x$

$$\ln(e^x) = x \qquad e^{\ln(y)} = y$$

- Recall that if  $g(y) = f^{-1}(y)$  then  $g'(y) = 1/f'(g(y))$
- Consider  $g(y) = \ln(y)$  and  $f(x) = e^x$  (with  $f'(x) = e^x$ )

$$\frac{d\ln(y)}{dy} = \frac{1}{e^{\ln(y)}}$$



# Natural Logarithms


- The natural logarithm is defined as the inverse of  $e^x$

$$\ln(e^x) = x \qquad e^{\ln(y)} = y$$


- Recall that if  $g(y) = f^{-1}(y)$  then  $g'(y) = 1/f'(g(y))$
- Consider  $g(y) = \ln(y)$  and  $f(x) = e^x$  (with  $f'(x) = e^x$ )

$$\frac{d\ln(y)}{dy} = \frac{1}{e^{\ln(y)}} = \frac{1}{y}$$


# Exponentials and Logarithms




# Exponentials and Logarithms




# Exponentials and Logarithms




# Exponentials and Logarithms




# Exponentials and Logarithms



# Exponentials and Logarithms




# Exponentials and Logarithms





# Outline

1. Why Calculus?
2. Differentiation
3. **Vector and Matrix Calculus**



# Derivatives in High Dimensions

- When working with functions  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  in many dimensions then there will typically be different derivative in different directions
- To compute the derivative in a direction  $\mathbf{u} \in \mathbb{R}^n$  (where  $\|\mathbf{u}\| = 1$ ) at a point  $\mathbf{x} \in \mathbb{R}^n$  we use

$$\partial_{\mathbf{u}} F(\mathbf{x}) = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{x} + \epsilon \mathbf{u}) - f(\mathbf{x})}{\epsilon}$$

- If  $\mathbf{u} = \boldsymbol{\delta}_i = (0, \dots, 0, 1, 0, \dots, 0)$  (i.e.  $u_i = 1$ ) then

$$\frac{\partial f(\mathbf{x})}{\partial x_i} = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{x} + \epsilon \boldsymbol{\delta}_i) - f(\mathbf{x})}{\epsilon}$$

# Derivatives in High Dimensions

- When working with functions  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  in many dimensions then there will typically be different derivative in different directions
- To compute the derivative in a direction  $\mathbf{u} \in \mathbb{R}^n$  (where  $\|\mathbf{u}\| = 1$ ) at a point  $\mathbf{x} \in \mathbb{R}^n$  we use

$$\partial_{\mathbf{u}} F(\mathbf{x}) = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{x} + \epsilon \mathbf{u}) - f(\mathbf{x})}{\epsilon}$$

- If  $\mathbf{u} = \delta_i = (0, \dots, 0, 1, 0, \dots, 0)$  (i.e.  $u_i = 1$ ) then

$$\frac{\partial f(\mathbf{x})}{\partial x_i} = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{x} + \epsilon \delta_i) - f(\mathbf{x})}{\epsilon}$$

# Derivatives in High Dimensions

- When working with functions  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  in many dimensions then there will typically be different derivative in different directions
- To compute the derivative in a direction  $\mathbf{u} \in \mathbb{R}^n$  (where  $\|\mathbf{u}\| = 1$ ) at a point  $\mathbf{x} \in \mathbb{R}^n$  we use

$$\partial_{\mathbf{u}} F(\mathbf{x}) = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{x} + \epsilon \mathbf{u}) - f(\mathbf{x})}{\epsilon}$$

- If  $\mathbf{u} = \delta_i = (0, \dots, 0, 1, 0, \dots, 0)$  (i.e.  $u_i = 1$ ) then

$$\frac{\partial f(\mathbf{x})}{\partial x_i} = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{x} + \epsilon \delta_i) - f(\mathbf{x})}{\epsilon}$$

# Taylor

- If we expand  $f(\mathbf{x} + \epsilon \mathbf{u})$  to first order in  $\epsilon$

$$f(\mathbf{x} + \epsilon \mathbf{u}) = f(\mathbf{x}) + \epsilon \mathbf{u}^\top \mathbf{g}(\mathbf{x}) + O(\epsilon^2)$$

then  $g_i(\mathbf{x}) = \frac{\partial f(\mathbf{x})}{\partial x_i}$

- Recall we defined the vector of first order derivatives of  $f(\mathbf{x})$  to be the gradient

$$\nabla f(\mathbf{x}) = \begin{pmatrix} \frac{\partial f(\mathbf{x})}{\partial x_1} \\ \frac{\partial f(\mathbf{x})}{\partial x_2} \\ \vdots \\ \frac{\partial f(\mathbf{x})}{\partial x_n} \end{pmatrix}$$

- Thus

$$f(\mathbf{x} + \epsilon \mathbf{u}) = f(\mathbf{x}) + \epsilon \mathbf{u}^\top \nabla f(\mathbf{x}) + O(\epsilon^2)$$

# Taylor

- If we expand  $f(\mathbf{x} + \epsilon \mathbf{u})$  to first order in  $\epsilon$

$$f(\mathbf{x} + \epsilon \mathbf{u}) = f(\mathbf{x}) + \epsilon \mathbf{u}^\top \mathbf{g}(\mathbf{x}) + O(\epsilon^2)$$

then  $g_i(\mathbf{x}) = \frac{\partial f(\mathbf{x})}{\partial x_i}$

- Recall we defined the vector of first order derivatives of  $f(\mathbf{x})$  to be the gradient

$$\nabla f(\mathbf{x}) = \begin{pmatrix} \frac{\partial f(\mathbf{x})}{\partial x_1} \\ \frac{\partial f(\mathbf{x})}{\partial x_2} \\ \vdots \\ \frac{\partial f(\mathbf{x})}{\partial x_n} \end{pmatrix}$$

- Thus

$$f(\mathbf{x} + \epsilon \mathbf{u}) = f(\mathbf{x}) + \epsilon \mathbf{u}^\top \nabla f(\mathbf{x}) + O(\epsilon^2)$$

# Taylor

- If we expand  $f(\mathbf{x} + \epsilon \mathbf{u})$  to first order in  $\epsilon$

$$f(\mathbf{x} + \epsilon \mathbf{u}) = f(\mathbf{x}) + \epsilon \mathbf{u}^\top \mathbf{g}(\mathbf{x}) + O(\epsilon^2)$$

then  $g_i(\mathbf{x}) = \frac{\partial f(\mathbf{x})}{\partial x_i}$

- Recall we defined the vector of first order derivatives of  $f(\mathbf{x})$  to be the gradient

$$\nabla f(\mathbf{x}) = \begin{pmatrix} \frac{\partial f(\mathbf{x})}{\partial x_1} \\ \frac{\partial f(\mathbf{x})}{\partial x_2} \\ \vdots \\ \frac{\partial f(\mathbf{x})}{\partial x_n} \end{pmatrix}$$

- Thus

$$f(\mathbf{x} + \epsilon \mathbf{u}) = f(\mathbf{x}) + \epsilon \mathbf{u}^\top \nabla f(\mathbf{x}) + O(\epsilon^2)$$

# Taylor

- If we expand  $f(\mathbf{x} + \epsilon \mathbf{u})$  to first order in  $\epsilon$

$$f(\mathbf{x} + \epsilon \mathbf{u}) = f(\mathbf{x}) + \epsilon \mathbf{u}^\top \mathbf{g}(\mathbf{x}) + O(\epsilon^2)$$

then  $g_i(\mathbf{x}) = \frac{\partial f(\mathbf{x})}{\partial x_i}$

- Recall we defined the vector of first order derivatives of  $f(\mathbf{x})$  to be the gradient

$$\nabla f(\mathbf{x}) = \begin{pmatrix} \frac{\partial f(\mathbf{x})}{\partial x_1} \\ \frac{\partial f(\mathbf{x})}{\partial x_2} \\ \vdots \\ \frac{\partial f(\mathbf{x})}{\partial x_n} \end{pmatrix}$$

- Thus

$$f(\mathbf{x} + \epsilon \mathbf{u}) = f(\mathbf{x}) + \epsilon \mathbf{u}^\top \nabla f(\mathbf{x}) + O(\epsilon^2)$$

This is the start of the high-dimensional Taylor expansion



# Computing Gradients 1

- We can compute the gradient by writing out  $f(x)$  componentwise and performing the partial derivative with respect to  $x_i$

$$\nabla w^\top M w$$

# Computing Gradients 1

- We can compute the gradient by writing out  $f(\mathbf{x})$  componentwise and performing the partial derivative with respect to  $x_i$

$$\nabla w^\top M w$$

# Computing Gradients 1

- We can compute the gradient by writing out  $f(\mathbf{x})$  componentwise and performing the partial derivative with respect to  $x_i$

$$\nabla \mathbf{w}^\top \mathbf{M} \mathbf{w} = \begin{pmatrix} \frac{\partial}{\partial w_1} \\ \frac{\partial}{\partial w_2} \\ \frac{\partial}{\partial w_3} \\ \vdots \end{pmatrix} \sum_{i,j} w_i M_{ij} w_j$$

# Computing Gradients 1

- We can compute the gradient by writing out  $f(\mathbf{x})$  componentwise and performing the partial derivative with respect to  $x_i$

$$\nabla \mathbf{w}^\top \mathbf{M} \mathbf{w} = \begin{pmatrix} \frac{\partial}{\partial w_1} \\ \frac{\partial}{\partial w_2} \\ \frac{\partial}{\partial w_3} \\ \vdots \end{pmatrix} \sum_{i,j} w_i M_{ij} w_j = \begin{pmatrix} \sum_j M_{1j} w_j + \sum_i w_i M_{i1} \\ \sum_j M_{2j} w_j + \sum_i w_i M_{i2} \\ \sum_j M_{3j} w_j + \sum_i w_i M_{i3} \\ \vdots \end{pmatrix}$$

# Computing Gradients 1

- We can compute the gradient by writing out  $f(\mathbf{x})$  componentwise and performing the partial derivative with respect to  $x_i$

$$\begin{aligned}\nabla \mathbf{w}^\top \mathbf{M} \mathbf{w} &= \begin{pmatrix} \frac{\partial}{\partial w_1} \\ \frac{\partial}{\partial w_2} \\ \frac{\partial}{\partial w_3} \\ \vdots \end{pmatrix} \sum_{i,j} w_i M_{ij} w_j = \begin{pmatrix} \sum_j M_{1j} w_j + \sum_i w_i M_{i1} \\ \sum_j M_{2j} w_j + \sum_i w_i M_{i2} \\ \sum_j M_{3j} w_j + \sum_i w_i M_{i3} \\ \vdots \end{pmatrix} \\ &= \mathbf{M} \mathbf{w} + \mathbf{M}^\top \mathbf{w}\end{aligned}$$

# Computing Gradients 1

- We can compute the gradient by writing out  $f(\mathbf{x})$  componentwise and performing the partial derivative with respect to  $x_i$

$$\begin{aligned}\nabla \mathbf{w}^\top \mathbf{M} \mathbf{w} &= \begin{pmatrix} \frac{\partial}{\partial w_1} \\ \frac{\partial}{\partial w_2} \\ \frac{\partial}{\partial w_3} \\ \vdots \end{pmatrix} \sum_{i,j} w_i M_{ij} w_j = \begin{pmatrix} \sum_j M_{1j} w_j + \sum_i w_i M_{i1} \\ \sum_j M_{2j} w_j + \sum_i w_i M_{i2} \\ \sum_j M_{3j} w_j + \sum_i w_i M_{i3} \\ \vdots \end{pmatrix} \\ &= \mathbf{M} \mathbf{w} + \mathbf{M}^\top \mathbf{w}\end{aligned}$$

- It is tedious to compute these things component-wise, but when you need to understand what is going on then go back to the basics

# Computing Gradients 2

- A slicker way is just to expand  $f(\mathbf{x} + \epsilon \mathbf{u})$
- Consider  $f(\mathbf{x}) = \mathbf{x}^\top \mathbf{M} \mathbf{x} + \mathbf{a}^\top \mathbf{x}$

$$f(\mathbf{x} + \epsilon \mathbf{u}) = (\mathbf{x} + \epsilon \mathbf{u})^\top \mathbf{M} (\mathbf{x} + \epsilon \mathbf{u}) + \mathbf{a}^\top (\mathbf{x} + \epsilon \mathbf{u})$$

# Computing Gradients 2

- A slicker way is just to expand  $f(\mathbf{x} + \epsilon \mathbf{u})$
- Consider  $f(\mathbf{x}) = \mathbf{x}^\top \mathbf{M} \mathbf{x} + \mathbf{a}^\top \mathbf{x}$

$$f(\mathbf{x} + \epsilon \mathbf{u}) = (\mathbf{x} + \epsilon \mathbf{u})^\top \mathbf{M} (\mathbf{x} + \epsilon \mathbf{u}) + \mathbf{a}^\top (\mathbf{x} + \epsilon \mathbf{u})$$



# Computing Gradients 2

- A slicker way is just to expand  $f(x + \epsilon u)$
- Consider  $f(x) = x^\top \mathbf{M}x + a^\top x$

$$\begin{aligned} f(x + \epsilon u) &= (x + \epsilon u)^\top \mathbf{M}(x + \epsilon u) + a^\top (x + \epsilon u) \\ &= f(x) + \epsilon (u^\top \mathbf{M}x + x^\top \mathbf{M}u + a^\top u) + O(\epsilon^2) \end{aligned}$$

# Computing Gradients 2

- A slicker way is just to expand  $f(x + \epsilon u)$
- Consider  $f(x) = x^\top \mathbf{M}x + a^\top x$

$$\begin{aligned} f(x + \epsilon u) &= (x + \epsilon u)^\top \mathbf{M}(x + \epsilon u) + a^\top (x + \epsilon u) \\ &= f(x) + \epsilon (u^\top \mathbf{M}x + x^\top \mathbf{M}u + a^\top u) + O(\epsilon^2) \\ &= f(x) + \epsilon u^\top (\mathbf{M}x + \mathbf{M}^\top x + a) + O(\epsilon^2) \end{aligned}$$

using  $x^\top \mathbf{M}u = u^\top \mathbf{M}^\top x$  and  $a^\top u = u^\top a$

# Computing Gradients 2

- A slicker way is just to expand  $f(\mathbf{x} + \epsilon \mathbf{u})$
- Consider  $f(\mathbf{x}) = \mathbf{x}^\top \mathbf{M} \mathbf{x} + \mathbf{a}^\top \mathbf{x}$

$$\begin{aligned} f(\mathbf{x} + \epsilon \mathbf{u}) &= (\mathbf{x} + \epsilon \mathbf{u})^\top \mathbf{M} (\mathbf{x} + \epsilon \mathbf{u}) + \mathbf{a}^\top (\mathbf{x} + \epsilon \mathbf{u}) \\ &= f(\mathbf{x}) + \epsilon (\mathbf{u}^\top \mathbf{M} \mathbf{x} + \mathbf{x}^\top \mathbf{M} \mathbf{u} + \mathbf{a}^\top \mathbf{u}) + O(\epsilon^2) \\ &= f(\mathbf{x}) + \epsilon \mathbf{u}^\top (\mathbf{M} \mathbf{x} + \mathbf{M}^\top \mathbf{x} + \mathbf{a}) + O(\epsilon^2) \end{aligned}$$

using  $\mathbf{x}^\top \mathbf{M} \mathbf{u} = \mathbf{u}^\top \mathbf{M}^\top \mathbf{x}$  and  $\mathbf{a}^\top \mathbf{u} = \mathbf{u}^\top \mathbf{a}$

- But  $f(\mathbf{x} + \epsilon \mathbf{u}) = f(\mathbf{x}) + \epsilon \mathbf{u}^\top \nabla f(\mathbf{x}) + O(\epsilon^2)$  so

$$\nabla f(\mathbf{x}) = \mathbf{M} \mathbf{x} + \mathbf{M}^\top \mathbf{x} + \mathbf{a}$$

# Differentiating Matrices

- Often we have loss functions with respect to a matrix  $\mathbf{W}$ , e.g.

$$L(\mathbf{W}) = (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c)^2$$

- We might want to find the minimum with respect to  $\mathbf{W}$
- This occurs at a point  $\mathbf{W}^*$  where  $L(\mathbf{W})$  does not increase as we change  $\mathbf{W}$  in any way
- That is, we seek a  $\mathbf{W}^*$  such that, for any matrices  $\mathbf{U}$

$$L(\mathbf{W}^* + \epsilon \mathbf{U}) - L(\mathbf{W}^*) = O(\epsilon^2)$$

# Differentiating Matrices

- Often we have loss functions with respect to a matrix  $\mathbf{W}$ , e.g.

$$L(\mathbf{W}) = (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c)^2$$

- We might want to find the minimum with respect to  $\mathbf{W}$
- This occurs at a point  $\mathbf{W}^*$  where  $L(\mathbf{W})$  does not increase as we change  $\mathbf{W}$  in any way
- That is, we seek a  $\mathbf{W}^*$  such that, for any matrices  $\mathbf{U}$

$$L(\mathbf{W}^* + \epsilon \mathbf{U}) - L(\mathbf{W}^*) = O(\epsilon^2)$$

# Differentiating Matrices

- Often we have loss functions with respect to a matrix  $\mathbf{W}$ , e.g.

$$L(\mathbf{W}) = (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c)^2$$

- We might want to find the minimum with respect to  $\mathbf{W}$
- This occurs at a point  $\mathbf{W}^*$  where  $L(\mathbf{W})$  does not increase as we change  $\mathbf{W}$  in any way
- That is, we seek a  $\mathbf{W}^*$  such that, for any matrices  $\mathbf{U}$

$$L(\mathbf{W}^* + \epsilon \mathbf{U}) - L(\mathbf{W}^*) = O(\epsilon^2)$$

# Differentiating Matrices

- Often we have loss functions with respect to a matrix  $\mathbf{W}$ , e.g.

$$L(\mathbf{W}) = (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c)^2$$

- We might want to find the minimum with respect to  $\mathbf{W}$
- This occurs at a point  $\mathbf{W}^*$  where  $L(\mathbf{W})$  does not increase as we change  $\mathbf{W}$  in any way
- That is, we seek a  $\mathbf{W}^*$  such that, for any matrices  $\mathbf{U}$

$$L(\mathbf{W}^* + \epsilon \mathbf{U}) - L(\mathbf{W}^*) = O(\epsilon^2)$$

# Generalised Gradient

- We can generalise the idea of gradient to matrices

$$\frac{\partial L(\mathbf{W})}{\partial \mathbf{W}} = \begin{pmatrix} \frac{\partial L(\mathbf{W})}{\partial W_{11}} & \frac{\partial L(\mathbf{W})}{\partial W_{12}} & \cdots & \frac{\partial L(\mathbf{W})}{\partial W_{1m}} \\ \frac{\partial L(\mathbf{W})}{\partial W_{21}} & \frac{\partial L(\mathbf{W})}{\partial W_{22}} & \cdots & \frac{\partial L(\mathbf{W})}{\partial W_{2m}} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial L(\mathbf{W})}{\partial W_{n1}} & \frac{\partial L(\mathbf{W})}{\partial W_{n2}} & \cdots & \frac{\partial L(\mathbf{W})}{\partial W_{nm}} \end{pmatrix}$$

- From an identical argument we used for vectors

$$L(\mathbf{W} + \epsilon \mathbf{U}) = L(\mathbf{W}) + \epsilon \text{tr} \mathbf{U}^\top \frac{\partial L(\mathbf{W})}{\partial \mathbf{W}} + O(\epsilon^2)$$

where

$$\text{tr} \mathbf{U}^\top \mathbf{G} = \sum_i [\mathbf{U}^\top \mathbf{G}]_{ii} = \sum_{ij} U_{ji} G_{ji} = \sum_{ij} U_{ij} G_{ij} = \langle \mathbf{U}, \mathbf{G} \rangle$$



# Generalised Gradient

- We can generalise the idea of gradient to matrices

$$\frac{\partial L(\mathbf{W})}{\partial \mathbf{W}} = \begin{pmatrix} \frac{\partial L(\mathbf{W})}{\partial W_{11}} & \frac{\partial L(\mathbf{W})}{\partial W_{12}} & \cdots & \frac{\partial L(\mathbf{W})}{\partial W_{1m}} \\ \frac{\partial L(\mathbf{W})}{\partial W_{21}} & \frac{\partial L(\mathbf{W})}{\partial W_{22}} & \cdots & \frac{\partial L(\mathbf{W})}{\partial W_{2m}} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial L(\mathbf{W})}{\partial W_{n1}} & \frac{\partial L(\mathbf{W})}{\partial W_{n2}} & \cdots & \frac{\partial L(\mathbf{W})}{\partial W_{nm}} \end{pmatrix}$$

- From an identical argument we used for vectors

$$L(\mathbf{W} + \epsilon \mathbf{U}) = L(\mathbf{W}) + \epsilon \text{tr} \mathbf{U}^\top \frac{\partial L(\mathbf{W})}{\partial \mathbf{W}} + O(\epsilon^2)$$

where

$$\text{tr} \mathbf{U}^\top \mathbf{G} = \sum_i [\mathbf{U}^\top \mathbf{G}]_{ii} = \sum_{ij} U_{ji} G_{ji} = \sum_{ij} U_{ij} G_{ij} = \langle \mathbf{U}, \mathbf{G} \rangle$$

# Generalised Gradient

- We can generalise the idea of gradient to matrices

$$\frac{\partial L(\mathbf{W})}{\partial \mathbf{W}} = \begin{pmatrix} \frac{\partial L(\mathbf{W})}{\partial W_{11}} & \frac{\partial L(\mathbf{W})}{\partial W_{12}} & \cdots & \frac{\partial L(\mathbf{W})}{\partial W_{1m}} \\ \frac{\partial L(\mathbf{W})}{\partial W_{21}} & \frac{\partial L(\mathbf{W})}{\partial W_{22}} & \cdots & \frac{\partial L(\mathbf{W})}{\partial W_{2m}} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial L(\mathbf{W})}{\partial W_{n1}} & \frac{\partial L(\mathbf{W})}{\partial W_{n2}} & \cdots & \frac{\partial L(\mathbf{W})}{\partial W_{nm}} \end{pmatrix}$$

- From an identical argument we used for vectors

$$L(\mathbf{W} + \epsilon \mathbf{U}) = L(\mathbf{W}) + \epsilon \text{tr} \mathbf{U}^\top \frac{\partial L(\mathbf{W})}{\partial \mathbf{W}} + O(\epsilon^2)$$

where

$$\text{tr} \mathbf{U}^\top \mathbf{G} = \sum_i [\mathbf{U}^\top \mathbf{G}]_{ii} = \sum_{ij} U_{ji} G_{ji} = \sum_{ij} U_{ij} G_{ij} = \langle \mathbf{U}, \mathbf{G} \rangle$$

# Example

- Suppose

$$L(\mathbf{W}) = (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c)^2$$

then

$$L(\mathbf{W} + \epsilon \mathbf{U}) = (\mathbf{a}^\top (\mathbf{W} + \epsilon \mathbf{U}) \mathbf{b} - c)^2$$

# Example

- Suppose

$$L(\mathbf{W}) = (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c)^2$$

then

$$L(\mathbf{W} + \epsilon \mathbf{U}) = (\mathbf{a}^\top (\mathbf{W} + \epsilon \mathbf{U}) \mathbf{b} - c)^2$$

# Example

- Suppose

$$L(\mathbf{W}) = (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c)^2$$

then

$$L(\mathbf{W} + \epsilon \mathbf{U}) = (\mathbf{a}^\top (\mathbf{W} + \epsilon \mathbf{U}) \mathbf{b} - c)^2 = (\mathbf{a}^\top \mathbf{W} \mathbf{b} + \epsilon \mathbf{a}^\top \mathbf{U} \mathbf{b} - c)^2$$

# Example

- Suppose

$$L(\mathbf{W}) = (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c)^2$$

then

$$\begin{aligned} L(\mathbf{W} + \epsilon \mathbf{U}) &= (\mathbf{a}^\top (\mathbf{W} + \epsilon \mathbf{U}) \mathbf{b} - c)^2 = (\mathbf{a}^\top \mathbf{W} \mathbf{b} + \epsilon \mathbf{a}^\top \mathbf{U} \mathbf{b} - c)^2 \\ &= L(\mathbf{W}) + 2\epsilon (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c) (\mathbf{a}^\top \mathbf{U} \mathbf{b}) + O(\epsilon^2) \end{aligned}$$

# Example

- Suppose

$$L(\mathbf{W}) = (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c)^2$$

then

$$\begin{aligned} L(\mathbf{W} + \epsilon \mathbf{U}) &= (\mathbf{a}^\top (\mathbf{W} + \epsilon \mathbf{U}) \mathbf{b} - c)^2 = (\mathbf{a}^\top \mathbf{W} \mathbf{b} + \epsilon \mathbf{a}^\top \mathbf{U} \mathbf{b} - c)^2 \\ &= L(\mathbf{W}) + 2\epsilon (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c) (\mathbf{a}^\top \mathbf{U} \mathbf{b}) + O(\epsilon^2) \end{aligned}$$

- Now

$$\mathbf{a}^\top \mathbf{U} \mathbf{b} = \sum_{ij} a_i U_{ij} b_j$$

# Example

- Suppose

$$L(\mathbf{W}) = (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c)^2$$

then

$$\begin{aligned} L(\mathbf{W} + \epsilon \mathbf{U}) &= (\mathbf{a}^\top (\mathbf{W} + \epsilon \mathbf{U}) \mathbf{b} - c)^2 = (\mathbf{a}^\top \mathbf{W} \mathbf{b} + \epsilon \mathbf{a}^\top \mathbf{U} \mathbf{b} - c)^2 \\ &= L(\mathbf{W}) + 2\epsilon (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c) (\mathbf{a}^\top \mathbf{U} \mathbf{b}) + O(\epsilon^2) \end{aligned}$$

- Now

$$\mathbf{a}^\top \mathbf{U} \mathbf{b} = \sum_{ij} a_i U_{ij} b_j = \sum_{ij} U_{ji} a_j b_i$$



# Example

- Suppose

$$L(\mathbf{W}) = (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c)^2$$

then

$$\begin{aligned} L(\mathbf{W} + \epsilon \mathbf{U}) &= (\mathbf{a}^\top (\mathbf{W} + \epsilon \mathbf{U}) \mathbf{b} - c)^2 = (\mathbf{a}^\top \mathbf{W} \mathbf{b} + \epsilon \mathbf{a}^\top \mathbf{U} \mathbf{b} - c)^2 \\ &= L(\mathbf{W}) + 2\epsilon (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c) (\mathbf{a}^\top \mathbf{U} \mathbf{b}) + O(\epsilon^2) \end{aligned}$$

- Now

$$\mathbf{a}^\top \mathbf{U} \mathbf{b} = \sum_{ij} a_i U_{ij} b_j = \sum_{ij} U_{ji} a_j b_i = \text{tr} \mathbf{U}^\top \mathbf{a} \mathbf{b}^\top$$

# Example

- Suppose

$$L(\mathbf{W}) = (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c)^2$$

then

$$\begin{aligned} L(\mathbf{W} + \epsilon \mathbf{U}) &= (\mathbf{a}^\top (\mathbf{W} + \epsilon \mathbf{U}) \mathbf{b} - c)^2 = (\mathbf{a}^\top \mathbf{W} \mathbf{b} + \epsilon \mathbf{a}^\top \mathbf{U} \mathbf{b} - c)^2 \\ &= L(\mathbf{W}) + 2\epsilon (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c) (\mathbf{a}^\top \mathbf{U} \mathbf{b}) + O(\epsilon^2) \end{aligned}$$

- Now

$$\mathbf{a}^\top \mathbf{U} \mathbf{b} = \sum_{ij} a_i U_{ij} b_j = \sum_{ij} U_{ji} a_j b_i = \text{tr} \mathbf{U}^\top \mathbf{a} \mathbf{b}^\top$$

$$\text{Thus } \frac{\partial L(\mathbf{W})}{\partial \mathbf{W}} = 2 (\mathbf{a}^\top \mathbf{W} \mathbf{b} - c) \mathbf{a} \mathbf{b}^\top$$

# Traces

- The trace of a matrix is the sum of its diagonal elements

$$\text{tr} \mathbf{A} = \text{tr} \mathbf{A}^T = \sum_i A_{ii}$$

- Clearly  $\text{tr} c\mathbf{A} = c\text{tr} \mathbf{A}$
- Also  $\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr} \mathbf{A} + \text{tr} \mathbf{B}$
- We note that

$$\text{tr} \mathbf{A} \mathbf{B} = \sum_{i,j} A_{ij} B_{ji} = \sum_{i,j} B_{ij} A_{ji} = \text{tr} \mathbf{B} \mathbf{A}$$

- It follows that

$$\text{tr} \mathbf{A} \mathbf{B} \mathbf{C} \mathbf{D} = \text{tr} \mathbf{D} \mathbf{A} \mathbf{B} \mathbf{C}$$

# Traces

- The trace of a matrix is the sum of its diagonal elements

$$\text{tr}\mathbf{A} = \text{tr}\mathbf{A}^\top = \sum_i A_{ii}$$

- Clearly  $\text{trc}\mathbf{A} = \text{ctr}\mathbf{A}$
- Also  $\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr}\mathbf{A} + \text{tr}\mathbf{B}$
- We note that

$$\text{tr}\mathbf{A}\mathbf{B} = \sum_{i,j} A_{ij}B_{ji} = \sum_{i,j} B_{ij}A_{ji} = \text{tr}\mathbf{B}\mathbf{A}$$

- It follows that

$$\text{tr}\mathbf{ABCD} = \text{tr}\mathbf{DABC}$$

# Traces

- The trace of a matrix is the sum of its diagonal elements

$$\text{tr}\mathbf{A} = \text{tr}\mathbf{A}^\top = \sum_i A_{ii}$$

- Clearly  $\text{tr}c\mathbf{A} = c\text{tr}\mathbf{A}$
- Also  $\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr}\mathbf{A} + \text{tr}\mathbf{B}$
- We note that

$$\text{tr}\mathbf{A}\mathbf{B} = \sum_{i,j} A_{ij}B_{ji} = \sum_{i,j} B_{ij}A_{ji} = \text{tr}\mathbf{B}\mathbf{A}$$

- It follows that

$$\text{tr}\mathbf{ABCD} = \text{tr}\mathbf{DABC}$$

# Traces

- The trace of a matrix is the sum of its diagonal elements

$$\text{tr}\mathbf{A} = \text{tr}\mathbf{A}^\top = \sum_i A_{ii}$$

- Clearly  $\text{tr}c\mathbf{A} = c\text{tr}\mathbf{A}$
- Also  $\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr}\mathbf{A} + \text{tr}\mathbf{B}$
- We note that

$$\text{tr}\mathbf{A}\mathbf{B} = \sum_{i,j} A_{ij}B_{ji} = \sum_{i,j} B_{ij}A_{ji} = \text{tr}\mathbf{B}\mathbf{A}$$

- It follows that

$$\text{tr}\mathbf{ABCD} = \text{tr}\mathbf{DABC}$$

# Traces

- The trace of a matrix is the sum of its diagonal elements

$$\text{tr}\mathbf{A} = \text{tr}\mathbf{A}^\top = \sum_i A_{ii}$$

- Clearly  $\text{tr}c\mathbf{A} = c\text{tr}\mathbf{A}$
- Also  $\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr}\mathbf{A} + \text{tr}\mathbf{B}$
- We note that

$$\text{tr}\mathbf{A}\mathbf{B} = \sum_{i,j} A_{ij}B_{ji} = \sum_{i,j} B_{ij}A_{ji} = \text{tr}\mathbf{B}\mathbf{A}$$

- It follows that

$$\text{tr}\mathbf{ABCD} = \text{tr}\mathbf{DABC}$$

# Traces

- The trace of a matrix is the sum of its diagonal elements

$$\text{tr}\mathbf{A} = \text{tr}\mathbf{A}^\top = \sum_i A_{ii}$$

- Clearly  $\text{tr}c\mathbf{A} = c\text{tr}\mathbf{A}$
- Also  $\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr}\mathbf{A} + \text{tr}\mathbf{B}$
- We note that

$$\text{tr}\mathbf{A}\mathbf{B} = \sum_{i,j} A_{ij}B_{ji} = \sum_{i,j} B_{ij}A_{ji} = \text{tr}\mathbf{B}\mathbf{A}$$

- It follows that

$$\text{tr}\mathbf{A}\mathbf{B}\mathbf{C}\mathbf{D} = \text{tr}\mathbf{D}\mathbf{A}\mathbf{B}\mathbf{C} = \text{tr}\mathbf{C}\mathbf{D}\mathbf{A}\mathbf{B}$$



# Traces

- The trace of a matrix is the sum of its diagonal elements

$$\text{tr}\mathbf{A} = \text{tr}\mathbf{A}^\top = \sum_i A_{ii}$$

- Clearly  $\text{tr}c\mathbf{A} = c\text{tr}\mathbf{A}$
- Also  $\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr}\mathbf{A} + \text{tr}\mathbf{B}$
- We note that

$$\text{tr}\mathbf{A}\mathbf{B} = \sum_{i,j} A_{ij}B_{ji} = \sum_{i,j} B_{ij}A_{ji} = \text{tr}\mathbf{B}\mathbf{A}$$

- It follows that

$$\text{tr}\mathbf{A}\mathbf{B}\mathbf{C}\mathbf{D} = \text{tr}\mathbf{D}\mathbf{A}\mathbf{B}\mathbf{C} = \text{tr}\mathbf{C}\mathbf{D}\mathbf{A}\mathbf{B} = \text{tr}\mathbf{B}\mathbf{C}\mathbf{D}\mathbf{A}$$

# Quick Matrix Differentiation

- Let

$$\partial_{\mathbf{U}} f(\mathbf{X}) = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{X} + \epsilon \mathbf{U}) - f(\mathbf{X})}{\epsilon}$$

- E.g.

$$\begin{aligned} \partial_{\mathbf{U}} \text{tr} \mathbf{A} \mathbf{X} \mathbf{B} &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \text{tr} \mathbf{A} (\mathbf{X} + \epsilon \mathbf{U}) \mathbf{B} - \text{tr} \mathbf{A} \mathbf{X} \mathbf{B} \\ &= \text{tr} \mathbf{A} \mathbf{U} \mathbf{B} \end{aligned}$$

# Quick Matrix Differentiation

- Let

$$\partial_{\mathbf{U}} f(\mathbf{X}) = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{X} + \epsilon \mathbf{U}) - f(\mathbf{X})}{\epsilon} = \text{tr} \mathbf{U}^{\top} \frac{\partial f(\mathbf{X})}{\partial \mathbf{X}}$$

- E.g.

$$\begin{aligned} \partial_{\mathbf{U}} \text{tr} \mathbf{A} \mathbf{X} \mathbf{B} &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \text{tr} \mathbf{A} (\mathbf{X} + \epsilon \mathbf{U}) \mathbf{B} - \text{tr} \mathbf{A} \mathbf{X} \mathbf{B} \\ &= \text{tr} \mathbf{A} \mathbf{U} \mathbf{B} \end{aligned}$$

# Quick Matrix Differentiation

- Let

$$\partial_{\mathbf{U}} f(\mathbf{X}) = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{X} + \epsilon \mathbf{U}) - f(\mathbf{X})}{\epsilon} = \text{tr} \mathbf{U}^T \frac{\partial f(\mathbf{X})}{\partial \mathbf{X}}$$

- E.g.

$$\begin{aligned} \partial_{\mathbf{U}} \text{tr} \mathbf{A} \mathbf{X} \mathbf{B} &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \text{tr} \mathbf{A} (\mathbf{X} + \epsilon \mathbf{U}) \mathbf{B} - \text{tr} \mathbf{A} \mathbf{X} \mathbf{B} \\ &= \text{tr} \mathbf{A} \mathbf{U} \mathbf{B} \end{aligned}$$

# Quick Matrix Differentiation

- Let

$$\partial_{\mathbf{U}} f(\mathbf{X}) = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{X} + \epsilon \mathbf{U}) - f(\mathbf{X})}{\epsilon} = \text{tr} \mathbf{U}^T \frac{\partial f(\mathbf{X})}{\partial \mathbf{X}}$$

- E.g.

$$\begin{aligned} \partial_{\mathbf{U}} \text{tr} \mathbf{A} \mathbf{X} \mathbf{B} &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \text{tr} \mathbf{A} (\mathbf{X} + \epsilon \mathbf{U}) \mathbf{B} - \text{tr} \mathbf{A} \mathbf{X} \mathbf{B} \\ &= \text{tr} \mathbf{A} \mathbf{U} \mathbf{B} = \text{tr} \mathbf{B}^T \mathbf{U}^T \mathbf{A}^T \end{aligned}$$

# Quick Matrix Differentiation

- Let

$$\partial_{\mathbf{U}} f(\mathbf{X}) = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{X} + \epsilon \mathbf{U}) - f(\mathbf{X})}{\epsilon} = \text{tr} \mathbf{U}^T \frac{\partial f(\mathbf{X})}{\partial \mathbf{X}}$$

- E.g.

$$\begin{aligned} \partial_{\mathbf{U}} \text{tr} \mathbf{A} \mathbf{X} \mathbf{B} &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \text{tr} \mathbf{A} (\mathbf{X} + \epsilon \mathbf{U}) \mathbf{B} - \text{tr} \mathbf{A} \mathbf{X} \mathbf{B} \\ &= \text{tr} \mathbf{A} \mathbf{U} \mathbf{B} = \text{tr} \mathbf{B}^T \mathbf{U}^T \mathbf{A}^T = \text{tr} \mathbf{U}^T \mathbf{A}^T \mathbf{B}^T \end{aligned}$$

# Quick Matrix Differentiation

- Let

$$\partial_{\mathbf{U}} f(\mathbf{X}) = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{X} + \epsilon \mathbf{U}) - f(\mathbf{X})}{\epsilon} = \text{tr} \mathbf{U}^T \frac{\partial f(\mathbf{X})}{\partial \mathbf{X}}$$

- E.g.

$$\begin{aligned} \partial_{\mathbf{U}} \text{tr} \mathbf{A} \mathbf{X} \mathbf{B} &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \text{tr} \mathbf{A} (\mathbf{X} + \epsilon \mathbf{U}) \mathbf{B} - \text{tr} \mathbf{A} \mathbf{X} \mathbf{B} \\ &= \text{tr} \mathbf{A} \mathbf{U} \mathbf{B} = \text{tr} \mathbf{B}^T \mathbf{U}^T \mathbf{A}^T = \text{tr} \mathbf{U}^T \mathbf{A}^T \mathbf{B}^T \end{aligned}$$

thus

$$\frac{\partial \text{tr} \mathbf{A} \mathbf{X} \mathbf{B}}{\partial \mathbf{X}} = \mathbf{A}^T \mathbf{B}^T$$

# Log Determinants

- We often come across logarithms of determinants of matrices,  $\log(|\mathbf{M}|)$
- For GP we want to choose  $\mathbf{K}$  to maximise the marginal likelihood,  $\log(|\mathbf{K} + \sigma^2 \mathbf{I}|)$
- To find the derivative of  $\log(|\mathbf{X}|)$  we consider

$$\log(|\mathbf{X} + \epsilon \mathbf{U}|) = \log(|\mathbf{X}(\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U})|)$$



# Log Determinants

- We often come across logarithms of determinants of matrices,  $\log(|\mathbf{M}|)$
- For GP we want to choose  $\mathbf{K}$  to maximise the marginal likelihood,  $\log(|\mathbf{K} + \sigma^2 \mathbf{I}|)$
- To find the derivative of  $\log(|\mathbf{X}|)$  we consider

$$\log(|\mathbf{X} + \epsilon \mathbf{U}|) = \log(|\mathbf{X}(\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U})|)$$

# Log Determinants

- We often come across logarithms of determinants of matrices,  $\log(|\mathbf{M}|)$
- For GP we want to choose  $\mathbf{K}$  to maximise the marginal likelihood,  $\log(|\mathbf{K} + \sigma^2 \mathbf{I}|)$
- To find the derivative of  $\log(|\mathbf{X}|)$  we consider

$$\log(|\mathbf{X} + \epsilon \mathbf{U}|) = \log(|\mathbf{X}(\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U})|)$$

# Log Determinants

- We often come across logarithms of determinants of matrices,  $\log(|\mathbf{M}|)$
- For GP we want to choose  $\mathbf{K}$  to maximise the marginal likelihood,  $\log(|\mathbf{K} + \sigma^2 \mathbf{I}|)$
- To find the derivative of  $\log(|\mathbf{X}|)$  we consider

$$\begin{aligned}\log(|\mathbf{X} + \epsilon \mathbf{U}|) &= \log(|\mathbf{X}(\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U})|) \\ &= \log(|\mathbf{X}| |\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U}|)\end{aligned}$$

★ Using  $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$

# Log Determinants

- We often come across logarithms of determinants of matrices,  $\log(|\mathbf{M}|)$
- For GP we want to choose  $\mathbf{K}$  to maximise the marginal likelihood,  $\log(|\mathbf{K} + \sigma^2 \mathbf{I}|)$
- To find the derivative of  $\log(|\mathbf{X}|)$  we consider

$$\begin{aligned}\log(|\mathbf{X} + \epsilon \mathbf{U}|) &= \log(|\mathbf{X}(\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U})|) \\ &= \log(|\mathbf{X}| |\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U}|) \\ &= \log(|\mathbf{X}|) + \log(|\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U}|)\end{aligned}$$

- ★ Using  $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$
- ★ Using  $\log(ab) = \log(a) + \log(b)$

# Determinants

$$|\mathbf{I} + \epsilon \mathbf{M}| = \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix}$$

# Determinants

$$|\mathbf{I} + \epsilon \mathbf{M}| = \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix}$$

# Determinants

$$|\mathbf{I} + \epsilon \mathbf{M}| = \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12}$$

# Determinants

$$\begin{aligned} |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\ &= 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2) \end{aligned}$$



# Determinants

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\
 &= 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2)
 \end{aligned}$$

$$|\mathbf{I} + \epsilon \mathbf{M}| = \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{21} & \epsilon M_{31} & \epsilon M_{41} & \epsilon M_{51} \\ \epsilon M_{12} & 1 + \epsilon M_{22} & \epsilon M_{32} & \epsilon M_{42} & \epsilon M_{52} \\ \epsilon M_{13} & \epsilon M_{23} & 1 + \epsilon M_{33} & \epsilon M_{43} & \epsilon M_{53} \\ \epsilon M_{14} & \epsilon M_{24} & \epsilon M_{34} & 1 + \epsilon M_{44} & \epsilon M_{54} \\ \epsilon M_{15} & \epsilon M_{25} & \epsilon M_{35} & \epsilon M_{45} & 1 + \epsilon M_{55} \end{vmatrix}$$

# Determinants

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\
 &= 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2)
 \end{aligned}$$

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} & \epsilon M_{13} & \epsilon M_{14} & \epsilon M_{15} \\ \epsilon M_{21} & 1 + \epsilon M_{22} & \epsilon M_{23} & \epsilon M_{24} & \epsilon M_{25} \\ \epsilon M_{31} & \epsilon M_{32} & 1 + \epsilon M_{33} & \epsilon M_{34} & \epsilon M_{35} \\ \epsilon M_{41} & \epsilon M_{42} & \epsilon M_{43} & 1 + \epsilon M_{44} & \epsilon M_{45} \\ \epsilon M_{51} & \epsilon M_{52} & \epsilon M_{53} & \epsilon M_{54} & 1 + \epsilon M_{55} \end{vmatrix} \\
 &= (1 + \epsilon M_{11}) C_{11}
 \end{aligned}$$

# Determinants

$$|\mathbf{I} + \epsilon \mathbf{M}| = \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\ = 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2)$$

$$|\mathbf{I} + \epsilon \mathbf{M}| = \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} & \epsilon M_{13} & \epsilon M_{14} & \epsilon M_{15} \\ \epsilon M_{21} & 1 + \epsilon M_{22} & \epsilon M_{23} & \epsilon M_{24} & \epsilon M_{25} \\ \epsilon M_{31} & \epsilon M_{32} & 1 + \epsilon M_{33} & \epsilon M_{34} & \epsilon M_{35} \\ \epsilon M_{41} & \epsilon M_{42} & \epsilon M_{43} & 1 + \epsilon M_{44} & \epsilon M_{45} \\ \epsilon M_{51} & \epsilon M_{52} & \epsilon M_{53} & \epsilon M_{54} & 1 + \epsilon M_{55} \end{vmatrix} \\ = (1 + \epsilon M_{11}) C_{11} - \epsilon M_{21} C_{21}$$

# Determinants

$$|\mathbf{I} + \epsilon \mathbf{M}| = \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\ = 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2)$$

$$|\mathbf{I} + \epsilon \mathbf{M}| = \begin{vmatrix} 1 & \epsilon M_{12} & \epsilon M_{13} & \epsilon M_{14} & \epsilon M_{15} \\ \epsilon M_{21} & 1 + \epsilon M_{22} & \epsilon M_{23} & \epsilon M_{24} & \epsilon M_{25} \\ \epsilon M_{31} & \epsilon M_{32} & 1 + \epsilon M_{33} & \epsilon M_{34} & \epsilon M_{35} \\ \epsilon M_{41} & \epsilon M_{42} & \epsilon M_{43} & 1 + \epsilon M_{44} & \epsilon M_{45} \\ \epsilon M_{51} & \epsilon M_{52} & \epsilon M_{53} & \epsilon M_{54} & 1 + \epsilon M_{55} \end{vmatrix} \\ = (1 + \epsilon M_{11}) C_{11} - \epsilon M_{21} C_{21} + \epsilon M_{31} C_{31}$$

# Determinants

$$|\mathbf{I} + \epsilon \mathbf{M}| = \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\ = 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2)$$

$$|\mathbf{I} + \epsilon \mathbf{M}| = \begin{vmatrix} 1 & \epsilon M_{12} & \epsilon M_{13} & \epsilon M_{14} & \epsilon M_{15} \\ \epsilon M_{21} & 1 + \epsilon M_{22} & \epsilon M_{23} & \epsilon M_{24} & \epsilon M_{25} \\ \epsilon M_{31} & \epsilon M_{32} & 1 + \epsilon M_{33} & \epsilon M_{34} & \epsilon M_{35} \\ \epsilon M_{41} & \epsilon M_{42} & \epsilon M_{43} & 1 + \epsilon M_{44} & \epsilon M_{45} \\ \epsilon M_{51} & \epsilon M_{52} & \epsilon M_{53} & \epsilon M_{54} & 1 + \epsilon M_{55} \end{vmatrix} \\ = (1 + \epsilon M_{11}) C_{11} - \epsilon M_{21} C_{21} + \epsilon M_{31} C_{31} - \epsilon M_{41} C_{41}$$

# Determinants

$$|\mathbf{I} + \epsilon \mathbf{M}| = \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\ = 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2)$$

$$|\mathbf{I} + \epsilon \mathbf{M}| = \begin{vmatrix} 1 & \epsilon M_{12} & \epsilon M_{13} & \epsilon M_{14} & \epsilon M_{15} \\ \epsilon M_{21} & 1 + \epsilon M_{22} & \epsilon M_{23} & \epsilon M_{24} & \epsilon M_{25} \\ \epsilon M_{31} & \epsilon M_{32} & 1 + \epsilon M_{33} & \epsilon M_{34} & \epsilon M_{35} \\ \epsilon M_{41} & \epsilon M_{42} & \epsilon M_{43} & 1 + \epsilon M_{44} & \epsilon M_{45} \\ \epsilon M_{51} & \epsilon M_{52} & \epsilon M_{53} & \epsilon M_{54} & 1 + \epsilon M_{55} \end{vmatrix} \\ = (1 + \epsilon M_{11}) C_{11} - \epsilon M_{21} C_{21} + \epsilon M_{31} C_{31} - \epsilon M_{41} C_{41} + \epsilon M_{51} C_{51}$$

# Determinants

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\
 &= 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2)
 \end{aligned}$$

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{21} & \epsilon M_{31} & \epsilon M_{41} & \epsilon M_{51} \\ \epsilon M_{12} & 1 + \epsilon M_{22} & \epsilon M_{32} & \epsilon M_{42} & \epsilon M_{52} \\ \epsilon M_{13} & \epsilon M_{23} & 1 + \epsilon M_{33} & \epsilon M_{43} & \epsilon M_{53} \\ \epsilon M_{14} & \epsilon M_{24} & \epsilon M_{34} & 1 + \epsilon M_{44} & \epsilon M_{54} \\ \epsilon M_{15} & \epsilon M_{25} & \epsilon M_{35} & \epsilon M_{45} & 1 + \epsilon M_{55} \end{vmatrix} \\
 &= (1 + \epsilon M_{11}) C_{11} + O(\epsilon^2)
 \end{aligned}$$

# Determinants

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\
 &= 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2)
 \end{aligned}$$

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} & \epsilon M_{13} & \epsilon M_{14} & \epsilon M_{15} \\ \epsilon M_{21} & 1 + \epsilon M_{22} & \epsilon M_{23} & \epsilon M_{24} & \epsilon M_{25} \\ \epsilon M_{31} & \epsilon M_{32} & 1 + \epsilon M_{33} & \epsilon M_{34} & \epsilon M_{35} \\ \epsilon M_{41} & \epsilon M_{42} & \epsilon M_{43} & 1 + \epsilon M_{44} & \epsilon M_{45} \\ \epsilon M_{51} & \epsilon M_{52} & \epsilon M_{53} & \epsilon M_{54} & 1 + \epsilon M_{55} \end{vmatrix} \\
 &= (1 + \epsilon M_{11}) C_{11} + O(\epsilon^2)
 \end{aligned}$$



# Determinants

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\
 &= 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2)
 \end{aligned}$$

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} & \epsilon M_{13} & \epsilon M_{14} & \epsilon M_{15} \\ \epsilon M_{21} & 1 + \epsilon M_{22} & \epsilon M_{23} & \epsilon M_{24} & \epsilon M_{25} \\ \epsilon M_{31} & \epsilon M_{32} & 1 + \epsilon M_{33} & \epsilon M_{34} & \epsilon M_{35} \\ \epsilon M_{41} & \epsilon M_{42} & \epsilon M_{43} & 1 + \epsilon M_{44} & \epsilon M_{45} \\ \epsilon M_{51} & \epsilon M_{52} & \epsilon M_{53} & \epsilon M_{54} & 1 + \epsilon M_{55} \end{vmatrix} \\
 &= (1 + \epsilon M_{11})(1 + \epsilon M_{22}) C'_{22} + O(\epsilon^2)
 \end{aligned}$$

# Determinants

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\
 &= 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2)
 \end{aligned}$$

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{21} & \epsilon M_{31} & \epsilon M_{41} & \epsilon M_{51} \\ \epsilon M_{12} & 1 + \epsilon M_{22} & \epsilon M_{32} & \epsilon M_{42} & \epsilon M_{52} \\ \epsilon M_{13} & \epsilon M_{23} & 1 + \epsilon M_{33} & \epsilon M_{43} & \epsilon M_{53} \\ \epsilon M_{14} & \epsilon M_{24} & \epsilon M_{34} & 1 + \epsilon M_{44} & \epsilon M_{54} \\ \epsilon M_{15} & \epsilon M_{25} & \epsilon M_{35} & \epsilon M_{45} & 1 + \epsilon M_{55} \end{vmatrix} \\
 &= \prod_i (1 + \epsilon M_{ii}) + O(\epsilon^2)
 \end{aligned}$$

# Determinants

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\
 &= 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2)
 \end{aligned}$$

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} & \epsilon M_{13} & \epsilon M_{14} & \epsilon M_{15} \\ \epsilon M_{21} & 1 + \epsilon M_{22} & \epsilon M_{23} & \epsilon M_{24} & \epsilon M_{25} \\ \epsilon M_{31} & \epsilon M_{32} & 1 + \epsilon M_{33} & \epsilon M_{34} & \epsilon M_{35} \\ \epsilon M_{41} & \epsilon M_{42} & \epsilon M_{43} & 1 + \epsilon M_{44} & \epsilon M_{45} \\ \epsilon M_{51} & \epsilon M_{52} & \epsilon M_{53} & \epsilon M_{54} & 1 + \epsilon M_{55} \end{vmatrix} \\
 &= (1 + \epsilon \sum_i M_{ii}) + O(\epsilon^2)
 \end{aligned}$$

# Determinants

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{12} \\ \epsilon M_{21} & 1 + \epsilon M_{22} \end{vmatrix} = (1 + \epsilon M_{11})(1 + \epsilon M_{22}) - \epsilon^2 M_{21} M_{12} \\
 &= 1 + \epsilon(M_{11} + M_{22}) + O(\epsilon^2)
 \end{aligned}$$

$$\begin{aligned}
 |\mathbf{I} + \epsilon \mathbf{M}| &= \begin{vmatrix} 1 + \epsilon M_{11} & \epsilon M_{21} & \epsilon M_{31} & \epsilon M_{41} & \epsilon M_{51} \\ \epsilon M_{12} & 1 + \epsilon M_{22} & \epsilon M_{32} & \epsilon M_{42} & \epsilon M_{52} \\ \epsilon M_{13} & \epsilon M_{23} & 1 + \epsilon M_{33} & \epsilon M_{43} & \epsilon M_{53} \\ \epsilon M_{14} & \epsilon M_{24} & \epsilon M_{34} & 1 + \epsilon M_{44} & \epsilon M_{54} \\ \epsilon M_{15} & \epsilon M_{25} & \epsilon M_{35} & \epsilon M_{45} & 1 + \epsilon M_{55} \end{vmatrix} \\
 &= (1 + \epsilon \operatorname{tr} \mathbf{M}) + O(\epsilon^2)
 \end{aligned}$$

# Putting it Together

- Recall

$$\log(|\mathbf{X} + \epsilon \mathbf{U}|) - \log(|\mathbf{X}|) = \log(|\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U}|)$$

# Putting it Together

- Recall

$$\begin{aligned}\log(|\mathbf{X} + \epsilon \mathbf{U}|) - \log(|\mathbf{X}|) &= \log(|\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U}|) \\ &= \log(1 + \epsilon \operatorname{tr} \mathbf{X}^{-1} \mathbf{U} + O(\epsilon)^2)\end{aligned}$$

# Putting it Together

- Recall

$$\begin{aligned}\log(|\mathbf{X} + \epsilon \mathbf{U}|) - \log(|\mathbf{X}|) &= \log(|\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U}|) \\ &= \log(1 + \epsilon \operatorname{tr} \mathbf{X}^{-1} \mathbf{U} + O(\epsilon)^2) \\ &= \epsilon \operatorname{tr} \mathbf{X}^{-1} \mathbf{U} + O(\epsilon)^2\end{aligned}$$

using  $\log(1 + x) = x + \frac{x^2}{2} + \dots$

# Putting it Together

- Recall

$$\begin{aligned}\log(|\mathbf{X} + \epsilon \mathbf{U}|) - \log(|\mathbf{X}|) &= \log(|\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U}|) \\ &= \log(1 + \epsilon \operatorname{tr} \mathbf{X}^{-1} \mathbf{U} + O(\epsilon)^2) \\ &= \epsilon \operatorname{tr} \mathbf{X}^{-1} \mathbf{U} + O(\epsilon)^2 \\ &= \epsilon \operatorname{tr} \mathbf{U}^{\top} (\mathbf{X}^{-1})^{\top} + O(\epsilon)\end{aligned}$$

$$\text{using } \log(1 + x) = x + \frac{x^2}{2} + \dots$$



# Putting it Together

- Recall

$$\begin{aligned}\log(|\mathbf{X} + \epsilon \mathbf{U}|) - \log(|\mathbf{X}|) &= \log(|\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U}|) \\ &= \log(1 + \epsilon \operatorname{tr} \mathbf{X}^{-1} \mathbf{U} + O(\epsilon)^2) \\ &= \epsilon \operatorname{tr} \mathbf{X}^{-1} \mathbf{U} + O(\epsilon)^2 \\ &= \epsilon \operatorname{tr} \mathbf{U}^T (\mathbf{X}^{-1})^T + O(\epsilon)\end{aligned}$$

using  $\log(1 + x) = x + \frac{x^2}{2} + \dots$

- Thus  $\partial_{\mathbf{U}} \log(|\mathbf{X}|) = \operatorname{tr} \mathbf{U}^T (\mathbf{X}^{-1})^T$

# Putting it Together

- Recall

$$\begin{aligned}\log(|\mathbf{X} + \epsilon \mathbf{U}|) - \log(|\mathbf{X}|) &= \log(|\mathbf{I} + \epsilon \mathbf{X}^{-1} \mathbf{U}|) \\ &= \log(1 + \epsilon \operatorname{tr} \mathbf{X}^{-1} \mathbf{U} + O(\epsilon)^2) \\ &= \epsilon \operatorname{tr} \mathbf{X}^{-1} \mathbf{U} + O(\epsilon)^2 \\ &= \epsilon \operatorname{tr} \mathbf{U}^T (\mathbf{X}^{-1})^T + O(\epsilon)\end{aligned}$$

using  $\log(1 + x) = x + \frac{x^2}{2} + \dots$

- Thus  $\partial_{\mathbf{U}} \log(|\mathbf{X}|) = \operatorname{tr} \mathbf{U}^T (\mathbf{X}^{-1})^T$

- Or

$$\frac{\partial \log(|\mathbf{X}|)}{\partial \mathbf{X}} = (\mathbf{X}^{-1})^T$$

# Summary

- With care you can differentiate most expressions
- The chain and product rule are incredibly powerful tools
- We can generalise differentiation to vectors and matrices
- There are a number of surprisingly useful results

# Summary

- With care you can differentiate most expressions
- The chain and product rule are incredibly powerful tools
- We can generalise differentiation to vectors and matrices
- There are a number of surprisingly useful results

# Summary

- With care you can differentiate most expressions
- The chain and product rule are incredibly powerful tools
- We can generalise differentiation to vectors and matrices
- There are a number of surprisingly useful results

# Summary

- With care you can differentiate most expressions
- The chain and product rule are incredibly powerful tools
- We can generalise differentiation to vectors and matrices
- There are a number of surprisingly useful results

# Summary

- With care you can differentiate most expressions
- The chain and product rule are incredibly powerful tools
- We can generalise differentiation to vectors and matrices
- There are a number of surprisingly useful results: see **The Matrix Cookbook**

# Summary

- With care you can differentiate most expressions
- The chain and product rule are incredibly powerful tools
- We can generalise differentiation to vectors and matrices
- There are a number of surprisingly useful results: see **The Matrix Cookbook**
- Next stop: integration