

(c) Now consider the case when $\mathbf{x} \in \mathbb{R}^n$. We assume that

$$g(\mathbf{x}) = \frac{1}{2}(\mathbf{x} - \mathbf{x}^*)^\top \mathbf{Q}(\mathbf{x} - \mathbf{x}^*)$$

where \mathbf{Q} is a symmetric, positive-definite matrix. The Hessian, \mathbf{H} , of $g(\mathbf{x})$ is a matrix with components

$$H_{ij} = \frac{\partial^2 g(\mathbf{x})}{\partial x_i \partial x_j}.$$

By writing out $g(\mathbf{x})$ as a double sum over the components compute the Hessian
[2 marks]

2

(d) Gradient descent in \mathbb{R}^n is given by

$$\mathbf{x}^{(t+1)} = \mathbf{x}^{(t)} - r \nabla g(\mathbf{x}).$$

Using the definition of $g(\mathbf{x})$ write down a recursion relation between $\mathbf{x}^{(t+1)}$ and $\mathbf{x}^{(t)}$.
[2 marks]

2

(e) Defining $\Delta^{(t)} = (\mathbf{x}^{(t)} - \mathbf{x}^*)$ obtain a recursion relation between $\Delta^{(t+1)}$ and $\Delta^{(t)}$.
(This is easy if you subtract \mathbf{x}^* from both sides of the recursion equation for $\mathbf{x}^{(t+1)}$.)
[2 marks]

2

- (f) Using the eigenvalue decomposition $\mathbf{Q} = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^T$ and defining $z^{(t)} = \mathbf{V}^T \Delta^t$ write out a recursion relation between $z^{(t+1)}$ and $z^{(t)}$. (This is helped by multiplying the recursion relation on the left by \mathbf{V}^T and using the fact that \mathbf{V} is an orthogonal matrix.) [3 marks]

3

- (g) Solve the recursion relation to obtain a formula for $x^{(t)}$ in terms of the initial state $x^{(0)}$. Express this formula for the i^{th} component of $x^{(t)}$ and hence find a condition on the learning rate r to ensure convergence. [4 marks]

4

End of question 1

(a) $\frac{1}{2}$ (b) $\frac{1}{5}$ (c) $\frac{1}{2}$ (d) $\frac{1}{2}$ (e) $\frac{1}{2}$ (f) $\frac{1}{3}$ (g) $\frac{1}{4}$ Total $\frac{1}{20}$
--

$$f(x) = \frac{c}{2}(x - x^*)^2 + \frac{d}{6}(x - x^*)^3$$
$$x^{(t+1)} = x^{(t)} - \frac{f'(x^{(t)})}{f''(x^{(t)})}.$$

(To do this we need to expand a term with the structure

$$\frac{r \vdash s \in}{u \vdash v \in}.$$

$$\frac{1}{u+v\epsilon} = \frac{1}{u} \frac{1}{1+\frac{v}{u}\epsilon} = \frac{1}{u} \left(1 - \frac{v}{u}\epsilon + \left(\frac{v}{u}\epsilon\right)^2 - \dots \right)$$

which is convergent provided $\left| \frac{v}{u} \right| < 1$.) [5 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There is no handwriting or other markings on the paper.

(b) Consider the function

$$h(x) = -x \log(x)$$

defined for $0 < x \leq 1$. By computing $h'(x)$ and setting $h'(x) = 0$ compute the value of x^* that maximises $h(x)$. [2 marks]

<hr/> <hr/> <hr/> <hr/> <hr/> <hr/>

$\frac{1}{2}$

(c) Compute $h''(x)$ and thus compute the Newton update function

$$n(x) = x - \frac{h'(x)}{h''(x)}$$

(the answer is rather surprising and in no way general). [3 marks]

<hr/> <hr/> <hr/> <hr/> <hr/> <hr/>


$\frac{1}{3}$

- (d) On the axes given below plot $x^{(t)}$ for $t = 0, 1, 2, \dots, 10$, starting from $x^{(0)} = 0.5$ where we use the gradient ascent updates

$$x^{(t+1)} = x^{(t)} + r h'(x^{(t)})$$

for $r = \{0.3, 0.4, 0.5\}$ (that is you should plot three curves).

Also plot $x^{(t)}$ where $x^{(t+1)} = n(x^{(t)})$ (that is using Newton's update formula) for $t = 0, 1, 2, 3$ and 4—note that to machine precision $x^{(5)} = x^*$. [10 marks]



10

End of question 2

(a) $\frac{5}{5}$	(b) $\frac{2}{2}$	(c) $\frac{3}{3}$	(d) $\frac{10}{10}$	Total $\frac{20}{20}$
-------------------	-------------------	-------------------	---------------------	-----------------------

3(a) Show that for $p_i > 0$ the function

$$h(\mathbf{p}) = -\sum_i p_i \log(p_i)$$

is strongly convex-down. Hint: show that the Hessian matrix is negative-definite.
[3 marks]

3

(b) Write down the Lagrangian, \mathcal{L} , for the problem of maximising $h(\mathbf{p})$ subject to the constraints

$$\sum_i p_i = 1 \qquad \sum_i p_i E_i = U.$$

Then explain why there is a unique solution to this constrained optimisation problem.

[2 marks]

2

- (c) By setting $\partial \mathcal{L} / \partial p_i = 0$ find the value of p_i that maximises \mathcal{L} in terms of E_i and the Lagrange multipliers. Use the constraint $\sum_i p_i = 1$ to eliminate the Lagrange multiplier that enforces this constraint. [5 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There is no handwriting or other markings on the paper.

15

End of question 3

(a) $\frac{1}{3}$ (b) $\frac{1}{2}$ (c) $\frac{1}{5}$ Total $\frac{1}{10}$