

# Advanced Machine Learning

## Vector Spaces

$$Mx=b$$

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 2 & 4 & 8 & 16 \\ 1 & 3 & 9 & 27 & 81 \\ 1 & 4 & 16 & 64 & 256 \\ 1 & 5 & 25 & 125 & 625 \end{pmatrix} \begin{pmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \\ a_4 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 2 \\ -3 \\ 1 \end{pmatrix}$$

$$b=M^{-1}x$$

$$Mv$$

$$v_i = \lambda_i v_i$$

$$\text{Tr}(X^{-1}A) = -X^{-1}AX^{-1}$$

Vectors, vector spaces, operators

# Outline

1. **Vector Spaces**
2. Operators

$$Mx=b$$

$$Mv_i = \lambda_i v_i$$

$$b=Mv$$

# Matrices, Vectors and All That

- The language of machine learning is mathematics
- Sometimes we draw pretty pictures to explain the mathematics
- Much of the mathematics we will use involves vectors, matrices and functions
- You need to master the language of mathematics, otherwise you won't understand the algorithms
- I'm going to spend this lecture and the next revising the mathematics you need to know

# Matrices, Vectors and All That

- The language of machine learning is mathematics
- Sometimes we draw pretty pictures to explain the mathematics
- Much of the mathematics we will use involves vectors, matrices and functions
- You need to master the language of mathematics, otherwise you won't understand the algorithms
- I'm going to spend this lecture and the next revising the mathematics you need to know

# Matrices, Vectors and All That

- The language of machine learning is mathematics
- Sometimes we draw pretty pictures to explain the mathematics
- Much of the mathematics we will use involves vectors, matrices and functions
- You need to master the language of mathematics, otherwise you won't understand the algorithms
- I'm going to spend this lecture and the next revising the mathematics you need to know

# Matrices, Vectors and All That

- The language of machine learning is mathematics
- Sometimes we draw pretty pictures to explain the mathematics
- Much of the mathematics we will use involves vectors, matrices and functions
- You need to master the language of mathematics, otherwise you won't understand the algorithms
- I'm going to spend this lecture and the next revising the mathematics you need to know

# Matrices, Vectors and All That

- The language of machine learning is mathematics
- Sometimes we draw pretty pictures to explain the mathematics
- Much of the mathematics we will use involves vectors, matrices and functions
- You need to master the language of mathematics, otherwise you won't understand the algorithms
- I'm going to spend this lecture and the next revising the mathematics you need to know

# Matrices, Vectors and All That

- The language of machine learning is mathematics
- Sometimes we draw pretty pictures to explain the mathematics
- Much of the mathematics we will use involves vectors, matrices and functions
- You need to master the language of mathematics, otherwise you won't understand the algorithms
- I'm going to spend this lecture and the next revising the mathematics you need to know (but I'm going use a slightly posher language than you are probably used to)



# Vectors

- We often work with objects with many components (features)
- To help handle this we will use vector notation
- We represent vectors by bold symbols
- All our vectors are column vectors by default
- We treat them as  $n \times 1$  matrix
- We write row vectors as transposes of column vectors

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

$$\mathbf{y}^T = (y_1, y_2, \dots, y_n)$$

# Vectors

- We often work with objects with many components (features)
- To help handle this we will use vector notation

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

- We represent vectors by bold symbols
  - All our vectors are column vectors by default
  - We treat them as  $n \times 1$  matrix
- We write row vectors as transposes of column vectors

$$\mathbf{y}^T = (y_1, y_2, \dots, y_n)$$

# Vectors

- We often work with objects with many components (features)
- To help handle this we will use vector notation
- We represent vectors by bold symbols
- All our vectors are column vectors by default
- We treat them as  $n \times 1$  matrix
- We write row vectors as transposes of column vectors

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

$$\mathbf{y}^T = (y_1, y_2, \dots, y_n)$$

# Vectors

- We often work with objects with many components (features)
- To help handle this we will use vector notation

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

- We represent vectors by bold symbols
  - All our vectors are column vectors by default
  - We treat them as  $n \times 1$  matrix
- We write row vectors as transposes of column vectors

$$\mathbf{y}^T = (y_1, y_2, \dots, y_n)$$

# Vectors

- We often work with objects with many components (features)
- To help handle this we will use vector notation
- We represent vectors by bold symbols
- All our vectors are column vectors by default
- We treat them as  $n \times 1$  matrix
- We write row vectors as transposes of column vectors

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

$$\mathbf{y}^T = (y_1, y_2, \dots, y_n)$$

# Vectors

- We often work with objects with many components (features)
- To help handle this we will use vector notation
- We represent vectors by bold symbols
- All our vectors are column vectors by default
- We treat them as  $n \times 1$  matrix
- We write row vectors as transposes of column vectors

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

$$\mathbf{y}^T = (y_1, y_2, \dots, y_n)$$

# Vectors


- We often work with objects with many components (features)
- To help handle this we will use vector notation
- We represent vectors by bold symbols
- All our vectors are column vectors by default
- We treat them as  $n \times 1$  matrix
- We write row vectors as transposes of column vectors

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

$$\mathbf{y}^T = (y_1, y_2, \dots, y_n)$$

# Basic Vector Operations

- The basic vector operations are adding




- multiplying by a scalar (a number)



# Basic Vector Operations


- The basic vector operations are adding



- multiplying by a scalar (a number)

# Basic Vector Operations


- The basic vector operations are adding



- multiplying by a scalar (a number)

# Basic Vector Operations


- The basic vector operations are adding




- multiplying by a scalar (a number)

# Basic Vector Operations

- The basic vector operations are adding




- multiplying by a scalar (a number)




# Basic Vector Operations

- The basic vector operations are adding



- multiplying by a scalar (a number)



# Vector Space

- A vector space,  $\mathcal{V}$ , is a set of vectors which satisfies

1. if  $\mathbf{v}, \mathbf{w} \in \mathcal{V}$  then  $a \mathbf{v} \in \mathcal{V}$  and  $\mathbf{v} + \mathbf{w} \in \mathcal{V}$  (closure)
2.  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}$  (commutativity of addition)
3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associativity of addition)
4.  $\mathbf{v} + \mathbf{0} = \mathbf{v}$  (existence of additive identity  $\mathbf{0}$ )
5.  $1 \mathbf{v} = \mathbf{v}$  (existence of multiplicative identity 1)
6.  $a (b \mathbf{v}) = (a b) \mathbf{v}$  (distributive properties)
7.  $a (\mathbf{v} + \mathbf{w}) = a \mathbf{v} + a \mathbf{w}$
8.  $(a + b) \mathbf{v} = a \mathbf{v} + b \mathbf{v}$

(You don't need to remember these)

- Just from these properties we can deduce other properties

# Vector Space

- A vector space,  $\mathcal{V}$ , is a set of vectors which satisfies
  1. if  $\mathbf{v}, \mathbf{w} \in \mathcal{V}$  then  $a \mathbf{v} \in \mathcal{V}$  and  $\mathbf{v} + \mathbf{w} \in \mathcal{V}$  (closure)
  2.  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}$  (commutativity of addition)
  3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associativity of addition)
  4.  $\mathbf{v} + \mathbf{0} = \mathbf{v}$  (existence of additive identity  $\mathbf{0}$ )
  5.  $1 \mathbf{v} = \mathbf{v}$  (existence of multiplicative identity 1)
  6.  $a (b \mathbf{v}) = (a b) \mathbf{v}$  (distributive properties)
  7.  $a (\mathbf{v} + \mathbf{w}) = a \mathbf{v} + a \mathbf{w}$
  8.  $(a + b) \mathbf{v} = a \mathbf{v} + b \mathbf{v}$

(You don't need to remember these)

- Just from these properties we can deduce other properties

# Vector Space

- A vector space,  $\mathcal{V}$ , is a set of vectors which satisfies
  1. if  $\mathbf{v}, \mathbf{w} \in \mathcal{V}$  then  $a \mathbf{v} \in \mathcal{V}$  and  $\mathbf{v} + \mathbf{w} \in \mathcal{V}$  (closure)
  2.  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}$  (commutativity of addition)
  3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associativity of addition)
  4.  $\mathbf{v} + \mathbf{0} = \mathbf{v}$  (existence of additive identity  $\mathbf{0}$ )
  5.  $1 \mathbf{v} = \mathbf{v}$  (existence of multiplicative identity 1)
  6.  $a (b \mathbf{v}) = (a b) \mathbf{v}$  (distributive properties)
  7.  $a (\mathbf{v} + \mathbf{w}) = a \mathbf{v} + a \mathbf{w}$
  8.  $(a + b) \mathbf{v} = a \mathbf{v} + b \mathbf{v}$

(You don't need to remember these)

- Just from these properties we can deduce other properties



# Vector Space

- A vector space,  $\mathcal{V}$ , is a set of vectors which satisfies
  1. if  $\mathbf{v}, \mathbf{w} \in \mathcal{V}$  then  $a \mathbf{v} \in \mathcal{V}$  and  $\mathbf{v} + \mathbf{w} \in \mathcal{V}$  (closure)
  2.  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}$  (commutativity of addition)
  3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associativity of addition)
  4.  $\mathbf{v} + \mathbf{0} = \mathbf{v}$  (existence of additive identity  $\mathbf{0}$ )
  5.  $1 \mathbf{v} = \mathbf{v}$  (existence of multiplicative identity 1)
  6.  $a (b \mathbf{v}) = (a b) \mathbf{v}$  (distributive properties)
  7.  $a (\mathbf{v} + \mathbf{w}) = a \mathbf{v} + a \mathbf{w}$
  8.  $(a + b) \mathbf{v} = a \mathbf{v} + b \mathbf{v}$

(You don't need to remember these)

- Just from these properties we can deduce other properties

# Vector Space

- A vector space,  $\mathcal{V}$ , is a set of vectors which satisfies
  1. if  $\mathbf{v}, \mathbf{w} \in \mathcal{V}$  then  $a \mathbf{v} \in \mathcal{V}$  and  $\mathbf{v} + \mathbf{w} \in \mathcal{V}$  (closure)
  2.  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}$  (commutativity of addition)
  3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associativity of addition)
  4.  $\mathbf{v} + \mathbf{0} = \mathbf{v}$  (existence of additive identity  $\mathbf{0}$ )
  5.  $1 \mathbf{v} = \mathbf{v}$  (existence of multiplicative identity 1)
  6.  $a (b \mathbf{v}) = (a b) \mathbf{v}$  (distributive properties)
  7.  $a (\mathbf{v} + \mathbf{w}) = a \mathbf{v} + a \mathbf{w}$
  8.  $(a + b) \mathbf{v} = a \mathbf{v} + b \mathbf{v}$

(You don't need to remember these)

- Just from these properties we can deduce other properties

# Vector Space

- A vector space,  $\mathcal{V}$ , is a set of vectors which satisfies
  1. if  $\mathbf{v}, \mathbf{w} \in \mathcal{V}$  then  $a \mathbf{v} \in \mathcal{V}$  and  $\mathbf{v} + \mathbf{w} \in \mathcal{V}$  (closure)
  2.  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}$  (commutativity of addition)
  3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associativity of addition)
  4.  $\mathbf{v} + \mathbf{0} = \mathbf{v}$  (existence of additive identity  $\mathbf{0}$ )
  5.  $1 \mathbf{v} = \mathbf{v}$  (existence of multiplicative identity 1)
  6.  $a (b \mathbf{v}) = (a b) \mathbf{v}$  (distributive properties)
  7.  $a (\mathbf{v} + \mathbf{w}) = a \mathbf{v} + a \mathbf{w}$
  8.  $(a + b) \mathbf{v} = a \mathbf{v} + b \mathbf{v}$

(You don't need to remember these)

- Just from these properties we can deduce other properties

# Vector Space

- A vector space,  $\mathcal{V}$ , is a set of vectors which satisfies
  1. if  $\mathbf{v}, \mathbf{w} \in \mathcal{V}$  then  $a \mathbf{v} \in \mathcal{V}$  and  $\mathbf{v} + \mathbf{w} \in \mathcal{V}$  (closure)
  2.  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}$  (commutativity of addition)
  3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associativity of addition)
  4.  $\mathbf{v} + \mathbf{0} = \mathbf{v}$  (existence of additive identity  $\mathbf{0}$ )
  5.  $1 \mathbf{v} = \mathbf{v}$  (existence of multiplicative identity 1)
  6.  $a (b \mathbf{v}) = (a b) \mathbf{v}$  (distributive properties)
  7.  $a (\mathbf{v} + \mathbf{w}) = a \mathbf{v} + a \mathbf{w}$
  8.  $(a + b) \mathbf{v} = a \mathbf{v} + b \mathbf{v}$

(You don't need to remember these)

- Just from these properties we can deduce other properties

# Vector Space


- A vector space,  $\mathcal{V}$ , is a set of vectors which satisfies
  1. if  $\mathbf{v}, \mathbf{w} \in \mathcal{V}$  then  $a \mathbf{v} \in \mathcal{V}$  and  $\mathbf{v} + \mathbf{w} \in \mathcal{V}$  (closure)
  2.  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}$  (commutativity of addition)
  3.  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associativity of addition)
  4.  $\mathbf{v} + \mathbf{0} = \mathbf{v}$  (existence of additive identity  $\mathbf{0}$ )
  5.  $1 \mathbf{v} = \mathbf{v}$  (existence of multiplicative identity 1)
  6.  $a (b \mathbf{v}) = (a b) \mathbf{v}$  (distributive properties)
  7.  $a (\mathbf{v} + \mathbf{w}) = a \mathbf{v} + a \mathbf{w}$
  8.  $(a + b) \mathbf{v} = a \mathbf{v} + b \mathbf{v}$

(You don't need to remember these)

- Just from these properties we can deduce other properties


$$\mathbb{R}^n$$

- When we first learn about vectors we think of them as arrows in 3-D space
- If we centre them all at the origin then there is a one-to-one correspondence between vectors and points in space
- We call this vector space  $\mathbb{R}^3$
- Any set of quantities  $\mathbf{x} = (x_1, x_2, \dots, x_n)^\top$  which satisfy the axioms above form a vector space  $\mathbb{R}^n$
- Of course, we can't so easily draw pictures of high-dimensional vectors




$$\mathbb{R}^n$$

- When we first learn about vectors we think of them as arrows in 3-D space
- If we centre them all at the origin then there is a one-to-one correspondence between vectors and points in space
- We call this vector space  $\mathbb{R}^3$
- Any set of quantities  $\mathbf{x} = (x_1, x_2, \dots, x_n)^\top$  which satisfy the axioms above form a vector space  $\mathbb{R}^n$
- Of course, we can't so easily draw pictures of high-dimensional vectors



$$\mathbb{R}^n$$


- When we first learn about vectors we think of them as arrows in 3-D space
- If we centre them all at the origin then there is a one-to-one correspondence between vectors and points in space
- We call this vector space  $\mathbb{R}^3$
- Any set of quantities  $\mathbf{x} = (x_1, x_2, \dots, x_n)^\top$  which satisfy the axioms above form a vector space  $\mathbb{R}^n$
- Of course, we can't so easily draw pictures of high-dimensional vectors






$$\mathbb{R}^n$$

- When we first learn about vectors we think of them as arrows in 3-D space
- If we centre them all at the origin then there is a one-to-one correspondence between vectors and points in space
- We call this vector space  $\mathbb{R}^3$
- Any set of quantities  $\mathbf{x} = (x_1, x_2, \dots, x_n)^T$  which satisfy the axioms above form a vector space  $\mathbb{R}^n$
- Of course, we can't so easily draw pictures of high-dimensional vectors



$$\mathbb{R}^n$$

- When we first learn about vectors we think of them as arrows in 3-D space
- If we centre them all at the origin then there is a one-to-one correspondence between vectors and points in space
- We call this vector space  $\mathbb{R}^3$
- Any set of quantities  $\mathbf{x} = (x_1, x_2, \dots, x_n)^\top$  which satisfy the axioms above form a vector space  $\mathbb{R}^n$
- Of course, we can't so easily draw pictures of high-dimensional vectors



# Other Vector Spaces

- Vectors are not the only object that form vector spaces
- Matrices satisfy all the conditions of a vector space
- Infinite sequences form a vector space
- Functions form a vector space
  - ★ Let  $C(a, b)$  be the set of functions defined on the interval  $[a, b]$
  - ★ Note that if  $f(x), g(x) \in C(a, b)$  then  $af(x) \in C(a, b)$  and  $f(x) + g(x) \in C(a, b)$
- Bounded vectors **don't** form a vector space

# Other Vector Spaces

- Vectors are not the only object that form vector spaces
- Matrices satisfy all the conditions of a vector space
- Infinite sequences form a vector space
- Functions form a vector space
  - ★ Let  $C(a, b)$  be the set of functions defined on the interval  $[a, b]$
  - ★ Note that if  $f(x), g(x) \in C(a, b)$  then  $af(x) \in C(a, b)$  and  $f(x) + g(x) \in C(a, b)$
- Bounded vectors **don't** form a vector space

# Other Vector Spaces

- Vectors are not the only object that form vector spaces
- Matrices satisfy all the conditions of a vector space
- Infinite sequences form a vector space
- Functions form a vector space
  - ★ Let  $C(a, b)$  be the set of functions defined on the interval  $[a, b]$
  - ★ Note that if  $f(x), g(x) \in C(a, b)$  then  $af(x) \in C(a, b)$  and  $f(x) + g(x) \in C(a, b)$
- Bounded vectors **don't** form a vector space

# Other Vector Spaces

- Vectors are not the only object that form vector spaces
- Matrices satisfy all the conditions of a vector space
- Infinite sequences form a vector space
- Functions form a vector space
  - ★ Let  $C(a, b)$  be the set of functions defined on the interval  $[a, b]$
  - ★ Note that if  $f(x), g(x) \in C(a, b)$  then  $af(x) \in C(a, b)$  and  $f(x) + g(x) \in C(a, b)$
- Bounded vectors **don't** form a vector space

# Other Vector Spaces

- Vectors are not the only object that form vector spaces
- Matrices satisfy all the conditions of a vector space
- Infinite sequences form a vector space
- Functions form a vector space
  - ★ Let  $C(a, b)$  be the set of functions defined on the interval  $[a, b]$
  - ★ Note that if  $f(x), g(x) \in C(a, b)$  then  $af(x) \in C(a, b)$  and  $f(x) + g(x) \in C(a, b)$
- Bounded vectors **don't** form a vector space

# Other Vector Spaces

- Vectors are not the only object that form vector spaces
- Matrices satisfy all the conditions of a vector space
- Infinite sequences form a vector space
- Functions form a vector space
  - ★ Let  $C(a, b)$  be the set of functions defined on the interval  $[a, b]$
  - ★ Note that if  $f(x), g(x) \in C(a, b)$  then  $af(x) \in C(a, b)$  and  $f(x) + g(x) \in C(a, b)$
- Bounded vectors **don't** form a vector space



# Metrics

- Vector spaces become more interesting if we have a notion of distance
- We say  $d(\mathbf{x}, \mathbf{y})$  is a proper distance or **metric** if
  1.  $d(\mathbf{x}, \mathbf{y}) \geq 0$  (non-negativity)
  2.  $d(\mathbf{x}, \mathbf{y}) = 0$  iff  $\mathbf{x} = \mathbf{y}$  (identity of indiscernibles)
  3.  $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$  (symmetry)
  4.  $d(\mathbf{x}, \mathbf{y}) \leq d(\mathbf{x}, \mathbf{z}) + d(\mathbf{z}, \mathbf{y})$  (triangular inequality)
- There are typically many possible distances (e.g. Euclidean distance, Manhattan distance, etc.)
- Often one or more condition isn't satisfied then we have a **pseudo-metric**

# Metrics

- Vector spaces become more interesting if we have a notion of distance
- We say  $d(\mathbf{x}, \mathbf{y})$  is a proper distance or **metric** if
  1.  $d(\mathbf{x}, \mathbf{y}) \geq 0$  (non-negativity)
  2.  $d(\mathbf{x}, \mathbf{y}) = 0$  iff  $\mathbf{x} = \mathbf{y}$  (identity of indiscernibles)
  3.  $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$  (symmetry)
  4.  $d(\mathbf{x}, \mathbf{y}) \leq d(\mathbf{x}, \mathbf{z}) + d(\mathbf{z}, \mathbf{y})$  (triangular inequality)
- There are typically many possible distances (e.g. Euclidean distance, Manhattan distance, etc.)
- Often one or more condition isn't satisfied then we have a **pseudo-metric**

# Metrics

- Vector spaces become more interesting if we have a notion of distance
- We say  $d(\mathbf{x}, \mathbf{y})$  is a proper distance or **metric** if
  1.  $d(\mathbf{x}, \mathbf{y}) \geq 0$  (non-negativity)
  2.  $d(\mathbf{x}, \mathbf{y}) = 0$  iff  $\mathbf{x} = \mathbf{y}$  (identity of indiscernibles)
  3.  $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$  (symmetry)
  4.  $d(\mathbf{x}, \mathbf{y}) \leq d(\mathbf{x}, \mathbf{z}) + d(\mathbf{z}, \mathbf{y})$  (triangular inequality)
- There are typically many possible distances (e.g. Euclidean distance, Manhattan distance, etc.)
- Often one or more condition isn't satisfied then we have a **pseudo-metric**

# Metrics

- Vector spaces become more interesting if we have a notion of distance
- We say  $d(\mathbf{x}, \mathbf{y})$  is a proper distance or **metric** if
  1.  $d(\mathbf{x}, \mathbf{y}) \geq 0$  (non-negativity)
  2.  $d(\mathbf{x}, \mathbf{y}) = 0$  iff  $\mathbf{x} = \mathbf{y}$  (identity of indiscernibles)
  3.  $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$  (symmetry)
  4.  $d(\mathbf{x}, \mathbf{y}) \leq d(\mathbf{x}, \mathbf{z}) + d(\mathbf{z}, \mathbf{y})$  (triangular inequality)
- There are typically many possible distances (e.g. Euclidean distance, Manhattan distance, etc.)
- Often one or more condition isn't satisfied then we have a **pseudo-metric**

# Metrics

- Vector spaces become more interesting if we have a notion of distance
- We say  $d(\mathbf{x}, \mathbf{y})$  is a proper distance or **metric** if
  1.  $d(\mathbf{x}, \mathbf{y}) \geq 0$  (non-negativity)
  2.  $d(\mathbf{x}, \mathbf{y}) = 0$  iff  $\mathbf{x} = \mathbf{y}$  (identity of indiscernibles)
  3.  $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$  (symmetry)
  4.  $d(\mathbf{x}, \mathbf{y}) \leq d(\mathbf{x}, \mathbf{z}) + d(\mathbf{z}, \mathbf{y})$  (triangular inequality)
- There are typically many possible distances (e.g. Euclidean distance, Manhattan distance, etc.)
- Often one or more condition isn't satisfied then we have a **pseudo-metric**

# Metrics

- Vector spaces become more interesting if we have a notion of distance
- We say  $d(\mathbf{x}, \mathbf{y})$  is a proper distance or **metric** if
  1.  $d(\mathbf{x}, \mathbf{y}) \geq 0$  (non-negativity)
  2.  $d(\mathbf{x}, \mathbf{y}) = 0$  iff  $\mathbf{x} = \mathbf{y}$  (identity of indiscernibles)
  3.  $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$  (symmetry)
  4.  $d(\mathbf{x}, \mathbf{y}) \leq d(\mathbf{x}, \mathbf{z}) + d(\mathbf{z}, \mathbf{y})$  (triangular inequality)
- There are typically many possible distances (e.g. Euclidean distance, Manhattan distance, etc.)
- Often one or more condition isn't satisfied then we have a **pseudo-metric**

# Metrics

- Vector spaces become more interesting if we have a notion of distance
- We say  $d(\mathbf{x}, \mathbf{y})$  is a proper distance or **metric** if
  1.  $d(\mathbf{x}, \mathbf{y}) \geq 0$  (non-negativity)
  2.  $d(\mathbf{x}, \mathbf{y}) = 0$  iff  $\mathbf{x} = \mathbf{y}$  (identity of indiscernibles)
  3.  $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$  (symmetry)
  4.  $d(\mathbf{x}, \mathbf{y}) \leq d(\mathbf{x}, \mathbf{z}) + d(\mathbf{z}, \mathbf{y})$  (triangular inequality)
- There are typically many possible distances (e.g. Euclidean distance, Manhattan distance, etc.)
- Often one or more condition isn't satisfied then we have a **pseudo-metric**

# Norms

- Vector spaces are even more interesting with a notion of length
- **Norms** provide some measure of the size of a vector
- To formalise this we define the **norm** of an object  $v$  as  $\|v\|$  satisfying
  1.  $\|v\| > 0$  if  $v \neq \mathbf{0}$
  2.  $\|a v\| = |a| \|v\|$
  3.  $\|u + v\| \leq \|u\| + \|v\|$  (triangular inequality)
- When some criteria aren't satisfied we have a **pseudo-norms**
- Norms provide a metric  $d(x, y) = \|x - y\|$  (they are metric spaces)



# Norms

- Vector spaces are even more interesting with a notion of length
- **Norms** provide some measure of the size of a vector
- To formalise this we define the **norm** of an object  $v$  as  $\|v\|$  satisfying
  1.  $\|v\| > 0$  if  $v \neq \mathbf{0}$
  2.  $\|a v\| = |a| \|v\|$
  3.  $\|u + v\| \leq \|u\| + \|v\|$  (triangular inequality)
- When some criteria aren't satisfied we have a **pseudo-norms**
- Norms provide a metric  $d(x, y) = \|x - y\|$  (they are metric spaces)

# Norms

- Vector spaces are even more interesting with a notion of length
- **Norms** provide some measure of the size of a vector
- To formalise this we define the **norm** of an object  $\mathbf{v}$  as  $\|\mathbf{v}\|$  satisfying
  1.  $\|\mathbf{v}\| > 0$  if  $\mathbf{v} \neq \mathbf{0}$
  2.  $\|a \mathbf{v}\| = |a| \|\mathbf{v}\|$
  3.  $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$  (triangular inequality)
- When some criteria aren't satisfied we have a **pseudo-norms**
- Norms provide a metric  $d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|$  (they are metric spaces)

# Norms

- Vector spaces are even more interesting with a notion of length
- **Norms** provide some measure of the size of a vector
- To formalise this we define the **norm** of an object  $v$  as  $\|v\|$  satisfying
  1.  $\|v\| > 0$  if  $v \neq 0$
  2.  $\|a v\| = |a| \|v\|$
  3.  $\|u + v\| \leq \|u\| + \|v\|$  (triangular inequality)
- When some criteria aren't satisfied we have a **pseudo-norms**
- Norms provide a metric  $d(x, y) = \|x - y\|$  (they are metric spaces)

# Norms

- Vector spaces are even more interesting with a notion of length
- **Norms** provide some measure of the size of a vector
- To formalise this we define the **norm** of an object  $v$  as  $\|v\|$  satisfying
  1.  $\|v\| > 0$  if  $v \neq \mathbf{0}$
  2.  $\|a v\| = |a| \|v\|$
  3.  $\|u + v\| \leq \|u\| + \|v\|$  (triangular inequality)
- When some criteria aren't satisfied we have a **pseudo-norms**
- Norms provide a metric  $d(x, y) = \|x - y\|$  (they are metric spaces)

# Norms

- Vector spaces are even more interesting with a notion of length
- **Norms** provide some measure of the size of a vector
- To formalise this we define the **norm** of an object  $v$  as  $\|v\|$  satisfying
  1.  $\|v\| > 0$  if  $v \neq \mathbf{0}$
  2.  $\|a v\| = |a| \|v\|$
  3.  $\|u + v\| \leq \|u\| + \|v\|$  (triangular inequality)
- When some criteria aren't satisfied we have a **pseudo-norms**
- Norms provide a metric  $d(x, y) = \|x - y\|$  (they are metric spaces)

# Norms

- Vector spaces are even more interesting with a notion of length
- **Norms** provide some measure of the size of a vector
- To formalise this we define the **norm** of an object  $\mathbf{v}$  as  $\|\mathbf{v}\|$  satisfying
  1.  $\|\mathbf{v}\| > 0$  if  $\mathbf{v} \neq \mathbf{0}$
  2.  $\|a \mathbf{v}\| = |a| \|\mathbf{v}\|$
  3.  $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$  (triangular inequality)
- When some criteria aren't satisfied we have a **pseudo-norms**
- Norms provide a metric  $d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|$  (they are metric spaces)

# Vector Norms

- The familiar vector norm is the (Euclidean) two norm

$$\|\mathbf{v}\|_2 = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$$

- Other norms exist, such as the  $p$ -norm

$$\|\mathbf{v}\|_p = \left( \sum_{i=1}^n |v_i|^p \right)^{1/p}$$

- Other special cases include the 1-norm and the infinite norm

$$\|\mathbf{v}\|_1 = \sum_{i=1}^n |v_i| \qquad \|\mathbf{v}\|_\infty = \max_i |v_i|$$

- The 0-norm is a pseudo-norm as it does not satisfy condition 2

$$\|\mathbf{v}\|_0 = \text{number of non-zero components}$$

# Vector Norms

- The familiar vector norm is the (Euclidean) two norm

$$\|\mathbf{v}\|_2 = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$$

- Other norms exist, such as the  $p$ -norm

$$\|\mathbf{v}\|_p = \left( \sum_{i=1}^n |v_i|^p \right)^{1/p}$$

- Other special cases include the 1-norm and the infinite norm

$$\|\mathbf{v}\|_1 = \sum_{i=1}^n |v_i| \qquad \|\mathbf{v}\|_\infty = \max_i |v_i|$$

- The 0-norm is a pseudo-norm as it does not satisfy condition 2

$$\|\mathbf{v}\|_0 = \text{number of non-zero components}$$



# Vector Norms

- The familiar vector norm is the (Euclidean) two norm

$$\|\mathbf{v}\|_2 = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$$

- Other norms exist, such as the  $p$ -norm

$$\|\mathbf{v}\|_p = \left( \sum_{i=1}^n |v_i|^p \right)^{1/p}$$

- Other special cases include the 1-norm and the infinite norm

$$\|\mathbf{v}\|_1 = \sum_{i=1}^n |v_i| \qquad \|\mathbf{v}\|_\infty = \max_i |v_i|$$

- The 0-norm is a pseudo-norm as it does not satisfy condition 2

$$\|\mathbf{v}\|_0 = \text{number of non-zero components}$$

# Vector Norms

- The familiar vector norm is the (Euclidean) two norm

$$\|\mathbf{v}\|_2 = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$$

- Other norms exist, such as the  $p$ -norm

$$\|\mathbf{v}\|_p = \left( \sum_{i=1}^n |v_i|^p \right)^{1/p}$$

- Other special cases include the 1-norm and the infinite norm

$$\|\mathbf{v}\|_1 = \sum_{i=1}^n |v_i| \qquad \|\mathbf{v}\|_\infty = \max_i |v_i|$$

- The 0-norm is a pseudo-norm as it does not satisfy condition 2

$$\|\mathbf{v}\|_0 = \text{number of non-zero components}$$

# Vector Norms

- The familiar vector norm is the (Euclidean) two norm

$$\|\mathbf{v}\|_2 = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$$

- Other norms exist, such as the  $p$ -norm

$$\|\mathbf{v}\|_p = \left( \sum_{i=1}^n |v_i|^p \right)^{1/p}$$

- Other special cases include the 1-norm and the infinite norm

$$\|\mathbf{v}\|_1 = \sum_{i=1}^n |v_i| \qquad \|\mathbf{v}\|_\infty = \max_i |v_i|$$

- The 0-norm is a pseudo-norm as it does not satisfy condition 2

$$\|\mathbf{v}\|_0 = \text{number of non-zero components}$$

# Matrix Norms

- We can define norms for other objects
- The norm of a matrix encodes how large the mapping is
- The Frobenius norm is defined by

$$\|\mathbf{A}\|_F = \sqrt{\sum_{i=1}^m \sum_{j=1}^n |A_{ij}|^2}$$

- Many other norms exist including 1-norm, max-norm, etc.
- For square matrices, some, but not all, norms satisfy the inequality

$$\|\mathbf{A} \mathbf{B}\| \leq \|\mathbf{A}\| \times \|\mathbf{B}\|$$

# Matrix Norms

- We can define norms for other objects
- The norm of a matrix encodes how large the mapping is
- The Frobenius norm is defined by

$$\|\mathbf{A}\|_F = \sqrt{\sum_{i=1}^m \sum_{j=1}^n |A_{ij}|^2}$$

- Many other norms exist including 1-norm, max-norm, etc.
- For square matrices, some, but not all, norms satisfy the inequality

$$\|\mathbf{A} \mathbf{B}\| \leq \|\mathbf{A}\| \times \|\mathbf{B}\|$$

# Matrix Norms

- We can define norms for other objects
- The norm of a matrix encodes how large the mapping is
- The Frobenius norm is defined by

$$\|\mathbf{A}\|_F = \sqrt{\sum_{i=1}^m \sum_{j=1}^n |A_{ij}|^2}$$

- Many other norms exist including 1-norm, max-norm, etc.
- For square matrices, some, but not all, norms satisfy the inequality

$$\|\mathbf{A} \mathbf{B}\| \leq \|\mathbf{A}\| \times \|\mathbf{B}\|$$

# Matrix Norms

- We can define norms for other objects
- The norm of a matrix encodes how large the mapping is
- The Frobenius norm is defined by

$$\|\mathbf{A}\|_F = \sqrt{\sum_{i=1}^m \sum_{j=1}^n |A_{ij}|^2}$$

- Many other norms exist including 1-norm, max-norm, etc.
- For square matrices, some, but not all, norms satisfy the inequality

$$\|\mathbf{A} \mathbf{B}\| \leq \|\mathbf{A}\| \times \|\mathbf{B}\|$$

# Matrix Norms

- We can define norms for other objects
- The norm of a matrix encodes how large the mapping is
- The Frobenius norm is defined by

$$\|\mathbf{A}\|_F = \sqrt{\sum_{i=1}^m \sum_{j=1}^n |A_{ij}|^2}$$

- Many other norms exist including 1-norm, max-norm, etc.
- For square matrices, some, but not all, norms satisfy the inequality

$$\|\mathbf{A} \mathbf{B}\| \leq \|\mathbf{A}\| \times \|\mathbf{B}\|$$



# Compatible Norms

- A vector and matrix norm are said to be compatible if

$$\|\mathbf{M}\mathbf{v}\|_b \leq \|\mathbf{M}\|_a \times \|\mathbf{v}\|_b$$

(Frobenius and Euclidean norms are compatible)

- Norms provide quick ways to bound the maximum growth of a vector under a mapping induced by the matrix
- We will see that a measure of the sensitivity of a mapping is in terms of the ratio of its maximum effect to its minimum effect on a vector
- This is known as the **conditioning**, given by  $\|\mathbf{M}\| \times \|\mathbf{M}^{-1}\|$

# Compatible Norms

- A vector and matrix norm are said to be compatible if

$$\|\mathbf{M}\mathbf{v}\|_b \leq \|\mathbf{M}\|_a \times \|\mathbf{v}\|_b$$

(Frobenius and Euclidean norms are compatible)

- Norms provide quick ways to bound the maximum growth of a vector under a mapping induced by the matrix
- We will see that a measure of the sensitivity of a mapping is in terms of the ratio of its maximum effect to its minimum effect on a vector
- This is known as the **conditioning**, given by  $\|\mathbf{M}\| \times \|\mathbf{M}^{-1}\|$

# Compatible Norms

- A vector and matrix norm are said to be compatible if

$$\|\mathbf{M}\mathbf{v}\|_b \leq \|\mathbf{M}\|_a \times \|\mathbf{v}\|_b$$

(Frobenius and Euclidean norms are compatible)

- Norms provide quick ways to bound the maximum growth of a vector under a mapping induced by the matrix
- We will see that a measure of the sensitivity of a mapping is in terms of the ratio of its maximum effect to its minimum effect on a vector
- This is known as the **conditioning**, given by  $\|\mathbf{M}\| \times \|\mathbf{M}^{-1}\|$

# Compatible Norms

- A vector and matrix norm are said to be compatible if

$$\|\mathbf{M}\mathbf{v}\|_b \leq \|\mathbf{M}\|_a \times \|\mathbf{v}\|_b$$

(Frobenius and Euclidean norms are compatible)

- Norms provide quick ways to bound the maximum growth of a vector under a mapping induced by the matrix
- We will see that a measure of the sensitivity of a mapping is in terms of the ratio of its maximum effect to its minimum effect on a vector
- This is known as the **conditioning**, given by  $\|\mathbf{M}\| \times \|\mathbf{M}^{-1}\|$

# Function Norms

- Functions can also have norms, for example, if  $f(x)$  is defined in some interval  $\mathcal{I}$

$$\|f\|_{L_2} = \sqrt{\int_{x \in \mathcal{I}} f^2(x) \, dx}$$

- The  $L_2$  vector space is the set of function where  $\|f\|_{L_2} < \infty$
- The  $L_1$ -norm is given by  $\|f\|_{L_1} = \int_{x \in \mathcal{I}} |f(x)| \, dx$
- The infinite-norm is given by  $\|f\|_{\infty} = \max_{x \in \mathcal{I}} |f(x)|$

# Function Norms

- Functions can also have norms, for example, if  $f(x)$  is defined in some interval  $\mathcal{I}$

$$\|f\|_{L_2} = \sqrt{\int_{x \in \mathcal{I}} f^2(x) \, dx}$$

- The  $L_2$  vector space is the set of function where  $\|f\|_{L_2} < \infty$
- The  $L_1$ -norm is given by  $\|f\|_{L_1} = \int_{x \in \mathcal{I}} |f(x)| \, dx$
- The infinite-norm is given by  $\|f\|_{\infty} = \max_{x \in \mathcal{I}} |f(x)|$

# Function Norms

- Functions can also have norms, for example, if  $f(x)$  is defined in some interval  $\mathcal{I}$

$$\|f\|_{L_2} = \sqrt{\int_{x \in \mathcal{I}} f^2(x) \, dx}$$

- The  $L_2$  vector space is the set of function where  $\|f\|_{L_2} < \infty$
- The  $L_1$ -norm is given by  $\|f\|_{L_1} = \int_{x \in \mathcal{I}} |f(x)| \, dx$
- The infinite-norm is given by  $\|f\|_{\infty} = \max_{x \in \mathcal{I}} |f(x)|$

# Function Norms

- Functions can also have norms, for example, if  $f(x)$  is defined in some interval  $\mathcal{I}$

$$\|f\|_{L_2} = \sqrt{\int_{x \in \mathcal{I}} f^2(x) \, dx}$$

- The  $L_2$  vector space is the set of function where  $\|f\|_{L_2} < \infty$
- The  $L_1$ -norm is given by  $\|f\|_{L_1} = \int_{x \in \mathcal{I}} |f(x)| \, dx$
- The infinite-norm is given by  $\|f\|_{\infty} = \max_{x \in \mathcal{I}} |f(x)|$



# Normed Vector Spaces

- Vector spaces with a distance (metric spaces) and vector spaces with a norm (normed vector spaces) are interesting objects
- They allow you to define a topology (open/closed sets, etc.)
- You can build up ideas about connectedness, continuity, contractive maps, fixed-point theorems, . . .
- For the most part we are going to consider an even more powerful vector space that has an inner-product defined

# Normed Vector Spaces

- Vector spaces with a distance (metric spaces) and vector spaces with a norm (normed vector spaces) are interesting objects
- They allow you to define a topology (open/closed sets, etc.)
- You can build up ideas about connectedness, continuity, contractive maps, fixed-point theorems, . . .
- For the most part we are going to consider an even more powerful vector space that has an inner-product defined

# Normed Vector Spaces

- Vector spaces with a distance (metric spaces) and vector spaces with a norm (normed vector spaces) are interesting objects
- They allow you to define a topology (open/closed sets, etc.)
- You can build up ideas about connectedness, continuity, contractive maps, fixed-point theorems, . . .
- For the most part we are going to consider an even more powerful vector space that has an inner-product defined

# Normed Vector Spaces

- Vector spaces with a distance (metric spaces) and vector spaces with a norm (normed vector spaces) are interesting objects
- They allow you to define a topology (open/closed sets, etc.)
- You can build up ideas about connectedness, continuity, contractive maps, fixed-point theorems, . . .
- For the most part we are going to consider an even more powerful vector space that has an inner-product defined

# Inner Products

- We will often consider objects with an *inner product*
- For vectors in  $\mathbb{R}^n$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^\top \mathbf{y} = \sum_{i=1}^n x_i y_i$$

- For functions

$$\langle f, g \rangle = \int_{x \in \mathcal{I}} f(x) g(x) \, dx$$

- With matrices we don't usually think about inner products

# Inner Products

- We will often consider objects with an *inner product*
- For vectors in  $\mathbb{R}^n$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^\top \mathbf{y} = \sum_{i=1}^n x_i y_i$$

- For functions

$$\langle f, g \rangle = \int_{x \in \mathcal{I}} f(x) g(x) \, dx$$

- With matrices we don't usually think about inner products

# Inner Products

- We will often consider objects with an *inner product*
- For vectors in  $\mathbb{R}^n$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^\top \mathbf{y} = \sum_{i=1}^n x_i y_i$$

- For functions

$$\langle f, g \rangle = \int_{x \in \mathcal{I}} f(x) g(x) \, dx$$

- With matrices we don't usually think about inner products

# Inner Products

- We will often consider objects with an *inner product*
- For vectors in  $\mathbb{R}^n$

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^\top \mathbf{y} = \sum_{i=1}^n x_i y_i$$

- For functions


$$\langle f, g \rangle = \int_{x \in \mathcal{I}} f(x) g(x) \, dx$$

- With matrices we don't usually think about inner products



# Angles Between Vectors

- A natural interpretation of the inner product is in providing a measure of the angle between vectors



$$\langle x, y \rangle = x^T y = \|x\| \|y\| \cos(\theta)$$


- Vectors are orthogonal if  $\langle x, y \rangle = 0$
- We can extend this idea to functions

$$\langle f(x), g(x) \rangle = \int_{x \in \mathcal{I}} f(x) g(x) dx = \|f(x)\| \|g(x)\| \cos(\theta)$$

- Note that  $\sin(x)$  and  $\cos(x)$  are orthogonal in the interval  $[0, 2\pi]$

# Angles Between Vectors

- A natural interpretation of the inner product is in providing a measure of the angle between vectors



$$\langle x, y \rangle = x^T y = \|x\| \|y\| \cos(\theta)$$


- Vectors are orthogonal if  $\langle x, y \rangle = 0$
- We can extend this idea to functions

$$\langle f(x), g(x) \rangle = \int_{x \in \mathcal{I}} f(x) g(x) dx = \|f(x)\| \|g(x)\| \cos(\theta)$$

- Note that  $\sin(x)$  and  $\cos(x)$  are orthogonal in the interval  $[0, 2\pi]$

# Angles Between Vectors

- A natural interpretation of the inner product is in providing a measure of the angle between vectors



$$\langle x, y \rangle = x^T y = \|x\| \|y\| \cos(\theta)$$


- Vectors are orthogonal if  $\langle x, y \rangle = 0$
- We can extend this idea to functions

$$\langle f(x), g(x) \rangle = \int_{x \in \mathcal{I}} f(x) g(x) dx = \|f(x)\| \|g(x)\| \cos(\theta)$$

- Note that  $\sin(x)$  and  $\cos(x)$  are orthogonal in the interval  $[0, 2\pi]$

# Angles Between Vectors

- A natural interpretation of the inner product is in providing a measure of the angle between vectors



$$\langle x, y \rangle = x^T y = \|x\| \|y\| \cos(\theta)$$

- Vectors are orthogonal if  $\langle x, y \rangle = 0$
- We can extend this idea to functions

$$\langle f(x), g(x) \rangle = \int_{x \in \mathcal{I}} f(x) g(x) dx = \|f(x)\| \|g(x)\| \cos(\theta)$$

- Note that  $\sin(x)$  and  $\cos(x)$  are orthogonal in the interval  $[0, 2\pi]$

# Basis Functions

- Any set of vectors  $\{\mathbf{b}_i | i = 1, \dots\}$  that span the space can be used as a basis or coordinate system
- The simplest and most useful case is when the vectors are orthogonal and normalised (i.e.  $\|\mathbf{b}_i\| = 1$ )
- In  $\mathbb{R}^3$  we could use  $\mathbf{b}_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ ,  $\mathbf{b}_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ ,  $\mathbf{b}_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$
- This is not unique as we can rotate our basis vectors
- For an orthogonal basis we can write any vector as  $\mathbf{x} = \begin{pmatrix} \mathbf{x}^\top \mathbf{b}_1 \\ \mathbf{x}^\top \mathbf{b}_2 \\ \mathbf{x}^\top \mathbf{b}_3 \end{pmatrix}$

# Basis Functions

- Any set of vectors  $\{\mathbf{b}_i | i = 1, \dots\}$  that span the space can be used as a basis or coordinate system
- The simplest and most useful case is when the vectors are orthogonal and normalised (i.e.  $\|\mathbf{b}_i\| = 1$ )

- In  $\mathbb{R}^3$  we could use  $\mathbf{b}_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ ,  $\mathbf{b}_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ ,  $\mathbf{b}_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$

- This is not unique as we can rotate our basis vectors

- For an orthogonal basis we can write any vector as  $\mathbf{x} = \begin{pmatrix} \mathbf{x}^\top \mathbf{b}_1 \\ \mathbf{x}^\top \mathbf{b}_2 \\ \mathbf{x}^\top \mathbf{b}_3 \end{pmatrix}$

# Basis Functions

- Any set of vectors  $\{\mathbf{b}_i | i = 1, \dots\}$  that span the space can be used as a basis or coordinate system
- The simplest and most useful case is when the vectors are orthogonal and normalised (i.e.  $\|\mathbf{b}_i\| = 1$ )

- In  $\mathbb{R}^3$  we could use  $\mathbf{b}_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ ,  $\mathbf{b}_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ ,  $\mathbf{b}_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$

- This is not unique as we can rotate our basis vectors

- For an orthogonal basis we can write any vector as  $\mathbf{x} = \begin{pmatrix} \mathbf{x}^\top \mathbf{b}_1 \\ \mathbf{x}^\top \mathbf{b}_2 \\ \mathbf{x}^\top \mathbf{b}_3 \end{pmatrix}$

# Basis Functions

- Any set of vectors  $\{\mathbf{b}_i | i = 1, \dots\}$  that span the space can be used as a basis or coordinate system
- The simplest and most useful case is when the vectors are orthogonal and normalised (i.e.  $\|\mathbf{b}_i\| = 1$ )

- In  $\mathbb{R}^3$  we could use  $\mathbf{b}_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ ,  $\mathbf{b}_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ ,  $\mathbf{b}_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$

- This is not unique as we can rotate our basis vectors

- For an orthogonal basis we can write any vector as  $\mathbf{x} = \begin{pmatrix} \mathbf{x}^\top \mathbf{b}_1 \\ \mathbf{x}^\top \mathbf{b}_2 \\ \mathbf{x}^\top \mathbf{b}_3 \end{pmatrix}$



# Basis Functions

- Any set of vectors  $\{\mathbf{b}_i | i = 1, \dots\}$  that span the space can be used as a basis or coordinate system
- The simplest and most useful case is when the vectors are orthogonal and normalised (i.e.  $\|\mathbf{b}_i\| = 1$ )



- In  $\mathbb{R}^3$  we could use  $\mathbf{b}_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ ,  $\mathbf{b}_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ ,  $\mathbf{b}_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$

- This is not unique as we can rotate our basis vectors

- For an orthogonal basis we can write any vector as  $\mathbf{x} = \begin{pmatrix} \mathbf{x}^\top \mathbf{b}_1 \\ \mathbf{x}^\top \mathbf{b}_2 \\ \mathbf{x}^\top \mathbf{b}_3 \end{pmatrix}$



# Orthogonal Functions

- For functions we can use any ortho-normal set of functions as a basis
- The most familiar are the Fourier functions  $\sin(n\theta)$  and  $\cos(n\theta)$
- Any function in  $C(0, 2\pi)$  can be represented by a point  $\mathbf{f} = \begin{pmatrix} \langle f(x), b_1(x) \rangle \\ \langle f(x), b_2(x) \rangle \\ \vdots \end{pmatrix}$
- There might be an infinite number of components
- This is analogous to points in  $\mathbb{R}^n$  (for large  $n$ )





# Orthogonal Functions

- For functions we can use any ortho-normal set of functions as a basis
- The most familiar are the Fourier functions  $\sin(n\theta)$  and  $\cos(n\theta)$
- Any function in  $C(0, 2\pi)$  can be represented by a point  $\mathbf{f} = \begin{pmatrix} \langle f(x), b_1(x) \rangle \\ \langle f(x), b_2(x) \rangle \\ \vdots \end{pmatrix}$
- There might be an infinite number of components
- This is analogous to points in  $\mathbb{R}^n$  (for large  $n$ )





# Orthogonal Functions

- For functions we can use any ortho-normal set of functions as a basis
- The most familiar are the Fourier functions  $\sin(n\theta)$  and  $\cos(n\theta)$
- Any function in  $C(0, 2\pi)$  can be represented by a point  $\mathbf{f} = \begin{pmatrix} \langle f(x), b_1(x) \rangle \\ \langle f(x), b_2(x) \rangle \\ \vdots \end{pmatrix}$
- There might be an infinite number of components
- This is analogous to points in  $\mathbb{R}^n$  (for large  $n$ )





# Orthogonal Functions

- For functions we can use any ortho-normal set of functions as a basis
- The most familiar are the Fourier functions  $\sin(n\theta)$  and  $\cos(n\theta)$
- Any function in  $C(0, 2\pi)$  can be represented by a point  $\mathbf{f} = \begin{pmatrix} \langle f(x), b_1(x) \rangle \\ \langle f(x), b_2(x) \rangle \\ \vdots \end{pmatrix}$
- There might be an infinite number of components
- This is analogous to points in  $\mathbb{R}^n$  (for large  $n$ )



# Orthogonal Functions

- For functions we can use any ortho-normal set of functions as a basis
- The most familiar are the Fourier functions  $\sin(n\theta)$  and  $\cos(n\theta)$
- Any function in  $C(0, 2\pi)$  can be represented by a point  $\mathbf{f} = \begin{pmatrix} \langle f(x), b_1(x) \rangle \\ \langle f(x), b_2(x) \rangle \\ \vdots \end{pmatrix}$
- There might be an infinite number of components
- This is analogous to points in  $\mathbb{R}^n$  (for large  $n$ )



# Algebraic Structure

- We have gone to these lengths as we want to show that many properties of vectors are shared by other objects (matrices, functions, etc.)
- Mathematicians study *algebraic structures* such as vector spaces, metric spaces, Hilbert spaces (infinite dimensional spaces with a norm and an inner product)
- Many of the concepts (distance/metric, norms, inner products) will re-occur many times within the course

# Algebraic Structure

- We have gone to these lengths as we want to show that many properties of vectors are shared by other objects (matrices, functions, etc.)
- Mathematicians study *algebraic structures* such as vector spaces, metric spaces, Hilbert spaces (infinite dimensional spaces with a norm and an inner product)
- Many of the concepts (distance/metric, norms, inner products) will re-occur many times within the course



# Algebraic Structure

- We have gone to these lengths as we want to show that many properties of vectors are shared by other objects (matrices, functions, etc.)
- Mathematicians study *algebraic structures* such as vector spaces, metric spaces, Hilbert spaces (infinite dimensional spaces with a norm and an inner product)
- Many of the concepts (distance/metric, norms, inner products) will re-occur many times within the course

# Outline

1. Vector Spaces
2. Operators

$$Mx=b$$

$$Mv_i = \lambda_i v_i$$

$$b=Mv$$

# Operators

- In machine learning we are interested in transforming our input vectors into some output predictions
- To accomplish this we will apply some mapping or operators on the vector  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}'$
- This says that  $\mathcal{T}$  maps some object  $x \in \mathcal{V}$  to an object  $y = \mathcal{T}[x]$  in a new vector space  $\mathcal{V}'$
- This new vector space may or may not be the same as the original vector space
- Our objects may be any object in a vector space such as a function

# Operators

- In machine learning we are interested in transforming our input vectors into some output predictions
- To accomplish this we will apply some mapping or operators on the vector  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}'$
- This says that  $\mathcal{T}$  maps some object  $x \in \mathcal{V}$  to an object  $y = \mathcal{T}[x]$  in a new vector space  $\mathcal{V}'$
- This new vector space may or may not be the same as the original vector space
- Our objects may be any object in a vector space such as a function

# Operators

- In machine learning we are interested in transforming our input vectors into some output predictions
- To accomplish this we will apply some mapping or operators on the vector  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}'$
- This says that  $\mathcal{T}$  maps some object  $x \in \mathcal{V}$  to an object  $y = \mathcal{T}[x]$  in a new vector space  $\mathcal{V}'$
- This new vector space may or may not be the same as the original vector space
- Our objects may be any object in a vector space such as a function

# Operators

- In machine learning we are interested in transforming our input vectors into some output predictions
- To accomplish this we will apply some mapping or operators on the vector  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}'$
- This says that  $\mathcal{T}$  maps some object  $x \in \mathcal{V}$  to an object  $y = \mathcal{T}[x]$  in a new vector space  $\mathcal{V}'$
- This new vector space may or may not be the same as the original vector space
- Our objects may be any object in a vector space such as a function

# Operators

- In machine learning we are interested in transforming our input vectors into some output predictions
- To accomplish this we will apply some mapping or operators on the vector  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}'$
- This says that  $\mathcal{T}$  maps some object  $x \in \mathcal{V}$  to an object  $y = \mathcal{T}[x]$  in a new vector space  $\mathcal{V}'$
- This new vector space may or may not be the same as the original vector space
- Our objects may be any object in a vector space such as a function

# Linear Operators

- Operators are in general very complicated, but a particular nice set of operators are linear operators
- $\mathcal{T}$  is a linear operator if
  1.  $\mathcal{T}[a \mathbf{x}] = a \mathcal{T}[\mathbf{x}]$
  2.  $\mathcal{T}[\mathbf{x} + \mathbf{y}] = \mathcal{T}[\mathbf{x}] + \mathcal{T}[\mathbf{y}]$
- For normal vectors the most general linear operation is

$$\mathcal{T}[\mathbf{x}] = \mathbf{M} \mathbf{x}$$

where  $\mathbf{M}$  is a matrix



# Linear Operators

- Operators are in general very complicated, but a particular nice set of operators are linear operators
- $\mathcal{T}$  is a linear operator if
  1.  $\mathcal{T}[a \mathbf{x}] = a \mathcal{T}[\mathbf{x}]$
  2.  $\mathcal{T}[\mathbf{x} + \mathbf{y}] = \mathcal{T}[\mathbf{x}] + \mathcal{T}[\mathbf{y}]$
- For normal vectors the most general linear operation is

$$\mathcal{T}[\mathbf{x}] = \mathbf{M} \mathbf{x}$$

where  $\mathbf{M}$  is a matrix

# Linear Operators

- Operators are in general very complicated, but a particular nice set of operators are linear operators
- $\mathcal{T}$  is a linear operator if
  1.  $\mathcal{T}[a \mathbf{x}] = a \mathcal{T}[\mathbf{x}]$
  2.  $\mathcal{T}[\mathbf{x} + \mathbf{y}] = \mathcal{T}[\mathbf{x}] + \mathcal{T}[\mathbf{y}]$
- For normal vectors the most general linear operation is

$$\mathcal{T}[\mathbf{x}] = \mathbf{M} \mathbf{x}$$

where  $\mathbf{M}$  is a matrix

# Matrix multiplication

- For an  $\ell \times m$  matrix  $\mathbf{A}$  and an  $m \times n$  matrix  $\mathbf{B}$  we can compute the  $(\ell \times n)$  product,  $\mathbf{C} = \mathbf{A} \mathbf{B}$ , such that

$$C_{ij} = \sum_{k=1}^m A_{ik} B_{kj} \quad \left( \begin{array}{|c|} \hline \text{---} \\ \text{---} \\ \text{---} \\ \hline \end{array} \right) \left( \begin{array}{|c|c|c|} \hline | & | & | \\ \hline \end{array} \right) = \left( \begin{array}{|c|c|c|} \hline \blacksquare & \blacksquare & \blacksquare \\ \blacksquare & \blacksquare & \blacksquare \\ \blacksquare & \blacksquare & \blacksquare \\ \hline \end{array} \right)$$

- Treating the vector  $\mathbf{x}$  as a  $n \times 1$  matrix then

$$\mathbf{y} = \mathbf{A} \mathbf{x} \quad \Rightarrow \quad y_i = \sum_j M_{ij} x_j \quad \left( \begin{array}{|c|} \hline \text{---} \\ \text{---} \\ \text{---} \\ \hline \end{array} \right) \left( \begin{array}{|c|} \hline | \\ \hline \end{array} \right) = \left( \begin{array}{|c|} \hline \blacksquare \\ \blacksquare \\ \blacksquare \\ \hline \end{array} \right)$$

- Using the same matrix notation we can define the inner product as

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^\top \mathbf{y} = \sum_{i=1}^n x_i y_i \quad \left( \text{---} \right) \left( \begin{array}{|c|} \hline | \\ \hline \end{array} \right) = \left( \blacksquare \right)$$

# Matrix multiplication

- For an  $\ell \times m$  matrix  $\mathbf{A}$  and an  $m \times n$  matrix  $\mathbf{B}$  we can compute the  $(\ell \times n)$  product,  $\mathbf{C} = \mathbf{A} \mathbf{B}$ , such that

$$C_{ij} = \sum_{k=1}^m A_{ik} B_{kj} \quad \left( \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right) \left( \begin{array}{c} | \\ | \\ | \\ | \end{array} \right) = \left( \begin{array}{c} \text{■} \text{■} \text{■} \text{■} \\ \text{■} \text{■} \text{■} \text{■} \\ \text{■} \text{■} \text{■} \text{■} \\ \text{■} \text{■} \text{■} \text{■} \end{array} \right)$$

- Treating the vector  $\mathbf{x}$  as a  $n \times 1$  matrix then

$$\mathbf{y} = \mathbf{A} \mathbf{x} \quad \Rightarrow \quad y_i = \sum_j M_{ij} x_j \quad \left( \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right) \left( \begin{array}{c} | \\ | \\ | \\ | \end{array} \right) = \left( \begin{array}{c} \text{■} \\ \text{■} \\ \text{■} \\ \text{■} \end{array} \right)$$

- Using the same matrix notation we can define the inner product as

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^\top \mathbf{y} = \sum_{i=1}^n x_i y_i \quad \left( \text{---} \right) \left( \begin{array}{c} | \\ | \\ | \\ | \end{array} \right) = \left( \text{■} \right)$$

# Matrix multiplication

- For an  $\ell \times m$  matrix  $\mathbf{A}$  and an  $m \times n$  matrix  $\mathbf{B}$  we can compute the  $(\ell \times n)$  product,  $\mathbf{C} = \mathbf{A} \mathbf{B}$ , such that

$$C_{ij} = \sum_{k=1}^m A_{ik} B_{kj} \quad \left( \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right) \left( \begin{array}{c} | \\ | \\ | \\ | \end{array} \right) = \left( \begin{array}{c} \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \end{array} \right)$$

- Treating the vector  $\mathbf{x}$  as a  $n \times 1$  matrix then

$$\mathbf{y} = \mathbf{A} \mathbf{x} \quad \Rightarrow \quad y_i = \sum_j M_{ij} x_j \quad \left( \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right) \left( \begin{array}{c} | \\ | \\ | \\ | \end{array} \right) = \left( \begin{array}{c} \blacksquare \\ \blacksquare \\ \blacksquare \\ \blacksquare \end{array} \right)$$

- Using the same matrix notation we can define the inner product as

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^T \mathbf{y} = \sum_{i=1}^n x_i y_i \quad \left( \text{---} \right) \left( \begin{array}{c} | \\ | \\ | \\ | \end{array} \right) = \left( \blacksquare \right)$$

# Kernels

- The equivalent of a matrix for functions (i.e. a linear operator) is known as a kernel  $K(x, y)$

$$g(x) = \mathcal{T}[f] = \int_{y \in \mathcal{I}} K(x, y) f(y) \, dy$$

- Our domain does not need to be one dimensional, e.g.

$$g(\boldsymbol{x}) = \mathcal{T}[f] = \int_{\boldsymbol{y} \in \mathcal{I}} K(\boldsymbol{x}, \boldsymbol{y}) f(\boldsymbol{y}) \, d\boldsymbol{y}$$

- We shall soon see examples of high-dimensional kernels

# Kernels

- The equivalent of a matrix for functions (i.e. a linear operator) is known as a kernel  $K(x, y)$

$$g(x) = \mathcal{T}[f] = \int_{y \in \mathcal{I}} K(x, y) f(y) \, dy$$

- Our domain does not need to be one dimensional, e.g.

$$g(\boldsymbol{x}) = \mathcal{T}[f] = \int_{\boldsymbol{y} \in \mathcal{I}} K(\boldsymbol{x}, \boldsymbol{y}) f(\boldsymbol{y}) \, d\boldsymbol{y}$$

- We shall soon see examples of high-dimensional kernels

# Kernels

- The equivalent of a matrix for functions (i.e. a linear operator) is known as a kernel  $K(x, y)$

$$g(x) = \mathcal{T}[f] = \int_{y \in \mathcal{I}} K(x, y) f(y) \, dy$$

- Our domain does not need to be one dimensional, e.g.

$$g(\boldsymbol{x}) = \mathcal{T}[f] = \int_{\boldsymbol{y} \in \mathcal{I}} K(\boldsymbol{x}, \boldsymbol{y}) f(\boldsymbol{y}) \, d\boldsymbol{y}$$

- We shall soon see examples of high-dimensional kernels



# Kernels in Machine Learning

- Kernels are used heavily in machine learning
- In kernel methods such as SVM, SVR, Kernel-PCA
- They are also used in Gaussian Processes
- In all these cases we consider symmetric, positive semi-definite kernels
- Sometimes they can be interpreted as covariance between random functions

$$K(\mathbf{x}, \mathbf{y}) = \mathbb{E}_{f \sim \mathcal{P}} [(f(\mathbf{x}) - \mu(\mathbf{x})) (f(\mathbf{y}) - \mu(\mathbf{y}))]$$

# Kernels in Machine Learning

- Kernels are used heavily in machine learning
- In kernel methods such as SVM, SVR, Kernel-PCA
- They are also used in Gaussian Processes
- In all these cases we consider symmetric, positive semi-definite kernels
- Sometimes they can be interpreted as covariance between random functions

$$K(\mathbf{x}, \mathbf{y}) = \mathbb{E}_{f \sim \mathcal{P}} [(f(\mathbf{x}) - \mu(\mathbf{x})) (f(\mathbf{y}) - \mu(\mathbf{y}))]$$

# Kernels in Machine Learning

- Kernels are used heavily in machine learning
- In kernel methods such as SVM, SVR, Kernel-PCA
- They are also used in Gaussian Processes
- In all these cases we consider symmetric, positive semi-definite kernels
- Sometimes they can be interpreted as covariance between random functions

$$K(\mathbf{x}, \mathbf{y}) = \mathbb{E}_{f \sim \mathcal{P}} [(f(\mathbf{x}) - \mu(\mathbf{x})) (f(\mathbf{y}) - \mu(\mathbf{y}))]$$

# Kernels in Machine Learning

- Kernels are used heavily in machine learning
- In kernel methods such as SVM, SVR, Kernel-PCA
- They are also used in Gaussian Processes
- In all these cases we consider symmetric, positive semi-definite kernels
- Sometimes they can be interpreted as covariance between random functions

$$K(\mathbf{x}, \mathbf{y}) = \mathbb{E}_{f \sim \mathcal{P}} [(f(\mathbf{x}) - \mu(\mathbf{x})) (f(\mathbf{y}) - \mu(\mathbf{y}))]$$

# Kernels in Machine Learning

- Kernels are used heavily in machine learning
- In kernel methods such as SVM, SVR, Kernel-PCA
- They are also used in Gaussian Processes
- In all these cases we consider symmetric, positive semi-definite kernels
- Sometimes they can be interpreted as covariance between random functions


$$K(\mathbf{x}, \mathbf{y}) = \mathbb{E}_{f \sim \mathcal{P}} [(f(\mathbf{x}) - \mu(\mathbf{x})) (f(\mathbf{y}) - \mu(\mathbf{y}))]$$

# General Linear Mappings

- In general a linear operator will map vectors between different vector spaces
- E.g.  $\mathbb{R}^3 \rightarrow \mathbb{R}^2$


# General Linear Mappings

- In general a linear operator will map vectors between different vector spaces
- E.g.  $\mathbb{R}^3 \rightarrow \mathbb{R}^2$



# General Linear Mappings


- In general a linear operator will map vectors between different vector spaces
- E.g.  $\mathbb{R}^3 \rightarrow \mathbb{R}^2$






# General Linear Mappings

- In general a linear operator will map vectors between different vector spaces
- E.g.  $\mathbb{R}^3 \rightarrow \mathbb{R}^2$




# General Linear Mappings

- In general a linear operator will map vectors between different vector spaces
- E.g.  $\mathbb{R}^3 \rightarrow \mathbb{R}^2$




# General Linear Mappings

- In general a linear operator will map vectors between different vector spaces
- E.g.  $\mathbb{R}^3 \rightarrow \mathbb{R}^2$




# General Linear Mappings

- In general a linear operator will map vectors between different vector spaces
- E.g.  $\mathbb{R}^3 \rightarrow \mathbb{R}^2$




# General Linear Mappings

- In general a linear operator will map vectors between different vector spaces
- E.g.  $\mathbb{R}^3 \rightarrow \mathbb{R}^2$




# General Linear Mappings

- In general a linear operator will map vectors between different vector spaces
- E.g.  $\mathbb{R}^3 \rightarrow \mathbb{R}^2$




# General Linear Mappings

- In general a linear operator will map vectors between different vector spaces
- E.g.  $\mathbb{R}^3 \rightarrow \mathbb{R}^2$



# General Linear Mappings

- In general a linear operator will map vectors between different vector spaces
- E.g.  $\mathbb{R}^3 \rightarrow \mathbb{R}^2$





# Square Matrices

- We will spend a lot of time on operators that map from a vector space onto itself  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}$
- For vectors in  $\mathbb{R}^n$  such linear operators are represented by square matrices
- When there is a one-to-one mapping then we have a unique inverse
- We will study such mappings in detail in the next lecture

# Square Matrices

- We will spend a lot of time on operators that map from a vector space onto itself  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}$
- For vectors in  $\mathbb{R}^n$  such linear operators are represented by square matrices
- When there is a one-to-one mapping then we have a unique inverse
- We will study such mappings in detail in the next lecture

# Square Matrices

- We will spend a lot of time on operators that map from a vector space onto itself  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}$
- For vectors in  $\mathbb{R}^n$  such linear operators are represented by square matrices
- When there is a one-to-one mapping then we have a unique inverse
- We will study such mappings in detail in the next lecture

# Square Matrices

- We will spend a lot of time on operators that map from a vector space onto itself  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}$
- For vectors in  $\mathbb{R}^n$  such linear operators are represented by square matrices
- When there is a one-to-one mapping then we have a unique inverse
- We will study such mappings in detail in the next lecture

# Summary

- We haven't covered much machine learning as such
- But mathematics is the language of machine learning and you have to get used to it
- Mathematics is like programming, if you don't understand the syntax and you can't write it down then its meaningless
- We've taken a high level view of vector spaces, this will pay us back later as we look at kernel methods

# Summary

- We haven't covered much machine learning as such—sorry
- But mathematics is the language of machine learning and you have to get used to it
- Mathematics is like programming, if you don't understand the syntax and you can't write it down then it's meaningless
- We've taken a high level view of vector spaces, this will pay us back later as we look at kernel methods

# Summary

- We haven't covered much machine learning as such—sorry
- But mathematics is the language of machine learning and you have to get used to it
- Mathematics is like programming, if you don't understand the syntax and you can't write it down then it's meaningless
- We've taken a high level view of vector spaces, this will pay us back later as we look at kernel methods

# Summary

- We haven't covered much machine learning as such—sorry
- But mathematics is the language of machine learning and you have to get used to it
- Mathematics is like programming, if you don't understand the syntax and you can't write it down then it's meaningless
- We've taken a high level view of vector spaces, this will pay us back later as we look at kernel methods



# Summary

- We haven't covered much machine learning as such—sorry
- But mathematics is the language of machine learning and you have to get used to it
- Mathematics is like programming, if you don't understand the syntax and you can't write it down then it's meaningless
- We've taken a high level view of vector spaces, this will pay us back later as we look at kernel methods