



Hartree-Fock Stability

Applications to the Homogeneous Electron Gas

Evan Curtin

November 7, 2016

University of Illinois at Urbana-Champaign

Table of contents

1. Background Information
2. Hartree-Fock Stability
3. Homogeneous Electron Gas
4. Notes on Iterative Subspace Eigenvalue Methods
5. Results
6. Concluding Remarks


Background Information

Levels of Hartree-Fock Theory

| Method | Spinorbital | DoF | Eigenfunction of |
|--------------|--|-----|------------------------|
| Restricted | $\chi_j^\alpha(\vec{r}, \sigma) = \sum_{i=1}^N c_{ij} \phi_i(\vec{r}) \alpha(\sigma)$ $\chi_j^\beta(\vec{r}, \sigma) = \sum_{i=1}^N c_{ij} \phi_i(\vec{r}) \beta(\sigma)$ | N/2 | \hat{S}^2, \hat{S}_z |
| Unrestricted | $\chi_j^\alpha(\vec{r}, \sigma) = \sum_{i=1}^N c_{ij}^\alpha \phi_i(\vec{r}) \alpha(\sigma)$ $\chi_j^\beta(\vec{r}, \sigma) = \sum_{i=1}^N c_{ij}^\beta \phi_i(\vec{r}) \beta(\sigma)$ | N | \hat{S}_z |
| General | $\chi_j(\vec{r}, \sigma) = \sum_{i=1}^N [c_{ij}^\alpha \phi_i(\vec{r}) \alpha(\sigma) + c_{ij}^\beta \phi_i(\vec{r}) \beta(\sigma)]$ | 2N | Neither |

Restricted Minimization

- Hartee-Fock SCF guarantees only stationary energy w.r.t. change in orbitals
- The solution may be a maximum, minimum or saddle point



Restricted Minimization


- Hartee-Fock SCF guarantees only stationary energy w.r.t. change in orbitals
- The solution may be a maximum, minimum or saddle point

Within the Constrained Space




Restricted Minimization

- Restricted minima may correspond to minima in another dimension




Restricted Minimization

- Restricted minima may correspond to minima in another dimension
- Restricted minima may correspond to maxima in another dimension



Restricted Minimization

- Restricted minima may correspond to minima in another dimension
- Restricted minima may correspond to maxima in another dimension
- Restricted minima may be nonstationary



Hartree-Fock Stability

Hartree-Fock Stability

- Solving the HF equations guarantees only that the energy is **stationary**. The first order variation is therefore 0.

Hartree-Fock Stability

- Solving the HF equations guarantees only that the energy is **stationary**. The first order variation is therefore 0.
- We need to know if this is indeed a minimum.

Hartree-Fock Stability

- Solving the HF equations guarantees only that the energy is **stationary**. The first order variation is therefore 0.
- We need to know if this is indeed a minimum.
- We can determine this if we inspect the second order variation in the energy.

Hartree-Fock Stability

- Solving the HF equations guarantees only that the energy is **stationary**. The first order variation is therefore 0.
- We need to know if this is indeed a minimum.
- We can determine this if we inspect the second order variation in the energy.
- Thouless¹ showed a physically motivated derivation using Time-Dependent Hartree-Fock theory (TDHF).

Hartree-Fock Stability Conditions

The TDHF solution is given by the Liouville Von-Neumann Equation of Motion,

$$i\hbar \frac{\partial \rho}{\partial t} = [T + U, \rho], \quad (1)$$

Hartree-Fock Stability Conditions

The TDHF solution is given by the Liouville Von-Neumann Equation of Motion,

$$i\hbar \frac{\partial \rho}{\partial t} = [T + U, \rho], \quad (1)$$

Where U is the self-consistent potential,

$$U_{pq} = \sum_{r,s} (\rho_{pq,rs} - \langle pr|sq \rangle) \rho_{sr}. \quad (2)$$

Hartree-Fock Stability Conditions

The TDHF solution is given by the Liouville Von-Neumann Equation of Motion,

$$i\hbar \frac{\partial \rho}{\partial t} = [T + U, \rho], \quad (1)$$

Where U is the self-consistent potential,

$$U_{pq} = \sum_{r,s} (\rho_{pq,rs} - \langle pr|sq \rangle) \rho_{sr}. \quad (2)$$

Under the restrictions,

$$\rho^2 = \rho \quad \text{and} \quad \rho^\dagger = \rho, \quad (3)$$

Hartree-Fock Stability Conditions

The TDHF solution is given by the Liouville Von-Neumann Equation of Motion,

$$i\hbar \frac{\partial \rho}{\partial t} = [T + U, \rho], \quad (1)$$

Where U is the self-consistent potential,

$$U_{pq} = \sum_{r,s} (\rho_{pq,rs} - \langle pr|sq \rangle) \rho_{sr}. \quad (2)$$

Under the restrictions,

$$\rho^2 = \rho \quad \text{and} \quad \rho^\dagger = \rho, \quad (3)$$

and to first order in ρ_{ai} , the elements of the density matrix are ,

$$\rho_{ai} = C_{ai}, \quad \rho_{ia} = C_{ai}^*. \quad (4)$$

Hartree-Fock Stability Conditions

The TDHF EoM becomes,

$$i\hbar \frac{dC_{ai}(t)}{dt} = (\epsilon_a - \epsilon_i) + \sum_i^{occ} \sum_b^{vir} [\langle aj || ib \rangle C_{bj}(t) + \langle ab || ij \rangle C_{bj}^*(t)] . \quad (5)$$

Hartree-Fock Stability Conditions

The TDHF EoM becomes,

$$i\hbar \frac{dC_{ai}(t)}{dt} = (\epsilon_a - \epsilon_i) + \sum_i^{occ} \sum_b^{vir} [\langle aj || ib \rangle C_{bj}(t) + \langle ab || ij \rangle C_{bj}^*(t)] . \quad (5)$$

The solutions are complex exponentials,

$$C_{ai}(t) = \alpha X_{ai} e^{-i\omega t} + \alpha^* Y_{ai}^* e^{i\omega^* t}, \quad (6)$$

Hartree-Fock Stability Conditions

The TDHF EoM becomes,

$$i\hbar \frac{dC_{ai}(t)}{dt} = (\epsilon_a - \epsilon_i) + \sum_i^{occ} \sum_b^{vir} [\langle aj||ib \rangle C_{bj}(t) + \langle ab||ij \rangle C_{bj}^*(t)]. \quad (5)$$

The solutions are complex exponentials,

$$C_{ai}(t) = \alpha X_{ai} e^{-i\omega t} + \alpha^* Y_{ai}^* e^{i\omega^* t}, \quad (6)$$

whose amplitudes, X_{ai} and Y_{ai} satisfy,

$$\begin{aligned} (\epsilon_a - \epsilon_i) X_{ai} + \sum_i^{occ} \sum_b^{vir} [\langle aj||ib \rangle X_{bj} + \langle ab||ij \rangle Y_{bj}] &= \hbar\omega X_{ai} \\ (\epsilon_a - \epsilon_i) Y_{ai} + \sum_i^{occ} \sum_b^{vir} [\langle aj||ib \rangle X_{bj} + \langle ab||ij \rangle Y_{bj}] &= -\hbar\omega Y_{ai}. \end{aligned} \quad (7)$$

Hartree-Fock Stability Conditions

The TDHF EoM becomes,

$$i\hbar \frac{dC_{ai}(t)}{dt} = (\epsilon_a - \epsilon_i) + \sum_i^{occ} \sum_b^{vir} [\langle aj||ib \rangle C_{bj}(t) + \langle ab||ij \rangle C_{bj}^*(t)]. \quad (5)$$

The solutions are complex exponentials,

$$C_{ai}(t) = \alpha X_{ai} e^{-i\omega t} + \alpha^* Y_{ai}^* e^{i\omega^* t}, \quad (6)$$

whose amplitudes, X_{ai} and Y_{ai} satisfy,

$$\begin{aligned} (\epsilon_a - \epsilon_i) X_{ai} + \sum_i^{occ} \sum_b^{vir} [\langle aj||ib \rangle X_{bj} + \langle ab||ij \rangle Y_{bj}] &= \hbar\omega X_{ai} \\ (\epsilon_a - \epsilon_i) Y_{ai} + \sum_i^{occ} \sum_b^{vir} [\langle aj||ib \rangle X_{bj} + \langle ab||ij \rangle Y_{bj}] &= -\hbar\omega Y_{ai}. \end{aligned} \quad (7)$$

These are the equations of the Random Phase Approximation (RPA)

Hartree-Fock Stability Conditions

- The RPA frequencies represent electronic oscillations about the ground state

Hartree-Fock Stability Conditions

- The RPA frequencies represent electronic oscillations about the ground state
- Real frequencies are stable, imaginary frequencies are unstable

Hartree-Fock Stability Conditions

- The RPA frequencies represent electronic oscillations about the ground state
- Real frequencies are stable, imaginary frequencies are unstable
- The RPA frequencies are imaginary when the **Orbital Hessian** (aka stability matrix, electronic Hessian) eigenvalues are negative

$$\begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{B}^* & \mathbf{A}^* \end{bmatrix} \begin{bmatrix} \mathbf{X} \\ \mathbf{Y} \end{bmatrix} = \omega \begin{bmatrix} \mathbf{X} \\ \mathbf{Y} \end{bmatrix} \quad (8)$$

Hartree-Fock Stability Conditions

- The RPA frequencies represent electronic oscillations about the ground state
- Real frequencies are stable, imaginary frequencies are unstable
- The RPA frequencies are imaginary when the **Orbital Hessian** (aka stability matrix, electronic Hessian) eigenvalues are negative

$$\begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{B}^* & \mathbf{A}^* \end{bmatrix} \begin{bmatrix} \mathbf{X} \\ \mathbf{Y} \end{bmatrix} = \omega \begin{bmatrix} \mathbf{X} \\ \mathbf{Y} \end{bmatrix} \quad (8)$$

- Where

$$\begin{aligned} A_{ia,jb} &= \langle i^a | H - \langle 0 | H | 0 \rangle | j^b \rangle = (\epsilon_a - \epsilon_i) \delta_{ij} \delta_{ab} + \langle aj || ib \rangle \\ B_{ia,jb} &= \langle ij^a | H - \langle 0 | H | 0 \rangle | 0 \rangle = \langle ab || ij \rangle . \end{aligned} \quad (9)$$

Hartree-Fock Stability Conditions

The eigenvalue equation can be factorized depending on symmetry.

TABLE I. Matrices required to have positive semidefinite character as a necessary condition for stability of self-consistent solution of type X in the space of type Y .

| X \ Y | | | | | | |
|-----------------------|-------------------------------------|-------------------------------------|-------------------------------------|-----------------------------|-------------------------------|---------------------------|
| | Real RHF ^a | Complex RHF | Real UHF | Complex UHF | Real GHF | Complex GHF |
| Real RHF ^a | ${}^1\mathbf{A}' + {}^1\mathbf{B}'$ | ${}^1\mathbf{A}' - {}^1\mathbf{B}'$ | ${}^3\mathbf{A}' + {}^3\mathbf{B}'$ | | | |
| Complex RHF | | ${}^1\mathbf{H}'$ | | ${}^3\mathbf{H}'$ | | |
| Real UHF | | | $\mathbf{A}' + \mathbf{B}'$ | $\mathbf{A}' - \mathbf{B}'$ | $\mathbf{A}'' + \mathbf{B}''$ | |
| Complex UHF | | | | \mathbf{H}' | | \mathbf{H}'' |
| Real GHF | | | | | $\mathbf{A} + \mathbf{B}$ | $\mathbf{A} - \mathbf{B}$ |
| Complex GHF | | | | | | \mathbf{H} |

^aSinglet closed shell case ($n_\alpha = n_\beta$) only.

Homogeneous Electron Gas

Brief Overview

- Homogeneous Electron Gas (HEG) model, also known as Uniform Electron Gas or Jellium Model.
- Electrons in a box with "smeared" nuclei \rightarrow uniform positive background charge
- The total charge is constrained to be neutral,

$$V_{bg}(\mathbf{r}) = \sum_i \frac{-Ze^2}{|\mathbf{r} - \mathbf{R}_i|} \rightarrow -e^2 \int \frac{d\mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|}, \quad (10)$$

- and the background and coulomb terms cancel exactly,

$$V_{ee} = e^2 \int \frac{d\mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|}. \quad (11)$$

Brief Overview

- The discretized solutions are given by,

$$\epsilon_{\vec{k}} = \frac{\hbar^2 k^2}{2m} - \sum_{\vec{k}'}^{|\vec{k}| < k_f} \langle \vec{k}, \vec{k}' | \vec{k}', \vec{k} \rangle \quad (12)$$

- Where the two electron integral is given by

$$\langle \vec{k}, \vec{k}' | \vec{k}'', \vec{k}''' \rangle \stackrel{2D, 3D}{=} \begin{cases} \frac{\pi}{V} \frac{2^{D-1}}{|\vec{k} - \vec{k}''|^{D-1}} & \vec{k}''' = \vec{k} + \vec{k}' - \vec{k}'' \\ 0 & \text{else} \end{cases}$$
$$\langle k, k' | k'', k''' \rangle \stackrel{1D}{=} \begin{cases} e^{|k-k''|^2 a^2} \text{Ei}(-|k-k''|^2 a^2) ; & k''' = k + k' - k'' \\ 0 ; & \text{else} \end{cases} \quad (13)$$

Project Goal

- Can we use the HF-Stability analysis to determine where the HEG is unstable?

Project Goal

- Can we use the HF-Stability analysis to determine where the HEG is unstable?
- Will this predict the known tendency of crystallization at low density?

Project Goal

- Can we use the HF-Stability analysis to determine where the HEG is unstable?
- Will this predict the known tendency of crystallization at low density?
- It is known that the RHF-GHF instability persists at all densities for the HEG¹.

Project Goal

- Can we use the HF-Stability analysis to determine where the HEG is unstable?
- Will this predict the known tendency of crystallization at low density?
- It is known that the RHF-GHF instability persists at all densities for the HEG¹.
- Can we show this numerically, and discriminate between RHF-UHF and RHF-GHF instability?

Project Goal

- Can we use the HF-Stability analysis to determine where the HEG is unstable?
- Will this predict the known tendency of crystallization at low density?
- It is known that the RHF-GHF instability persists at all densities for the HEG¹.
- Can we show this numerically, and discriminate between RHF-UHF and RHF-GHF instability?
- (Future) Can we combine this with to improve the efficacy of correlation theories (CC, MBPT)?

Notes on Iterative Subspace Eigenvalue Methods

Subspace Methods in 1 Slide

- Sometimes the eigenvalue problem is too large to solve completely, and only a few eigenpairs are needed
- Subspace methods allow for finding some eigenpairs without full diagonalization.
- They need only a matrix-vector product (maybe diagonal elements too).
- A small subspace is constructed, then expanded slowly until the eigenpair residual norm,

$$||\mathbf{Ax} - \lambda\mathbf{x}||, \quad (14)$$

is below some threshold.

- The **Lanczos** and **Arnoldi** methods are the most commonly used¹, while the **Davidson**² method is likely more useful in a majority of chemical applications.

1. Saad, Y. Numerical Methods for Large Eigenvalue Problems; SIAM, 2011.

2. Davidson, E. R. J. Comput. Phys. 1975, 17 (1), 8794.

Convergence Properties

- The convergence of these subspace algorithms depends on:

Convergence Properties

- The convergence of these subspace algorithms depends on:
 - Number of eigenvalues requested

Convergence Properties

- The convergence of these subspace algorithms depends on:
 - Number of eigenvalues requested
 - The block size of the algorithm

Convergence Properties

- The convergence of these subspace algorithms depends on:
 - Number of eigenvalues requested
 - The block size of the algorithm
 - The density of the eigenvalue spectrum

Convergence Properties

- The convergence of these subspace algorithms depends on:
 - Number of eigenvalues requested
 - The block size of the algorithm
 - The density of the eigenvalue spectrum
 - The initial guess eigenvectors

Convergence Properties

- The convergence of these subspace algorithms depends on:
 - Number of eigenvalues requested
 - The block size of the algorithm
 - The density of the eigenvalue spectrum
 - The initial guess eigenvectors
 - The condition number of the matrix

Convergence Properties

- The convergence of these subspace algorithms depends on:
 - Number of eigenvalues requested
 - The block size of the algorithm
 - The density of the eigenvalue spectrum
 - The initial guess eigenvectors
 - The condition number of the matrix
 - The preconditioner (Davidson only)

Convergence Properties

- The convergence of these subspace algorithms depends on:
 - Number of eigenvalues requested
 - The block size of the algorithm
 - The density of the eigenvalue spectrum
 - The initial guess eigenvectors
 - The condition number of the matrix
 - The preconditioner (Davidson only)
- They can converge to an incorrect answer if:

Convergence Properties

- The convergence of these subspace algorithms depends on:
 - Number of eigenvalues requested
 - The block size of the algorithm
 - The density of the eigenvalue spectrum
 - The initial guess eigenvectors
 - The condition number of the matrix
 - The preconditioner (Davidson only)
- They can converge to an incorrect answer if:
 - The block size is too small compared to degeneracy

Convergence Properties

- The convergence of these subspace algorithms depends on:
 - Number of eigenvalues requested
 - The block size of the algorithm
 - The density of the eigenvalue spectrum
 - The initial guess eigenvectors
 - The condition number of the matrix
 - The preconditioner (Davidson only)
- They can converge to an incorrect answer if:
 - The block size is too small compared to degeneracy
 - The approximate eigenvectors become non-orthogonal

Convergence Properties

- The convergence of these subspace algorithms depends on:
 - Number of eigenvalues requested
 - The block size of the algorithm
 - The density of the eigenvalue spectrum
 - The initial guess eigenvectors
 - The condition number of the matrix
 - The preconditioner (Davidson only)
- They can converge to an incorrect answer if:
 - The block size is too small compared to degeneracy
 - The approximate eigenvectors become non-orthogonal
- My recommendation for guess eigenvectors is

$$v_j^{(i)} = \text{normalize} \left(\frac{1}{|A_{ii} - A_{jj}| + 1} \right). \quad (15)$$

Orthogonalization

- The condition number, κ , is bound from below by

$$\kappa \geq \frac{\text{Max}(\lambda_i)}{\text{Min}(\lambda_j)} \quad (16)$$

- The Gram-Schmidt procedure has numerical issues,

$$\|\mathbf{I} - \mathbf{Q}^T \mathbf{Q}\| \leq \frac{\alpha \kappa^2}{1 - \beta \kappa^2}. \quad (17)$$

- Modified Gram-Schmidt is better, but not perfect,


$$\|\mathbf{I} - \mathbf{Q}^T \mathbf{Q}\| \leq \frac{\gamma \kappa}{1 - \eta \kappa} \quad (18)$$

- May need multiple orthogonalization steps

Results


Matrix Size Scaling

- Scales as $2N_{exc}$
- N_{occ} and N_{vir} both scale with $N_{kpoints}^D$




Matrix-Vector Scaling

- Scales as $N_{exc} \times N_{occ}$
due to momentum
conservation




Davidson Scaling

- Davidson is Asymptotically quadratic.
- Full diagonalization is almost cubic.




Algorithm Wall time

- Roughly quadratic wrt matrix dimension.
- Outliers correspond to high condition number




Efficacy of Davidson's Algorithm

- Scales as $N_{exc} \times N_{occ}$ due to momentum conservation




Dependence on r_s

Crossover from stable to unstable agrees with previous results.



My Result



Published Result. Red = Square Lattice, Blue = Triangular Lattice

Concluding Remarks

Concluding Remarks

- Finding all the low energy conformers for a molecule is hard

Concluding Remarks

- Finding all the low energy conformers for a molecule is hard
- In order to get accurate energies/structures, computationally expensive methods should be employed

Concluding Remarks

- Finding all the low energy conformers for a molecule is hard
- In order to get accurate energies/structures, computationally expensive methods should be employed
- These take time, so we want to minimize how many of these we do

Concluding Remarks

- Finding all the low energy conformers for a molecule is hard
- In order to get accurate energies/structures, computationally expensive methods should be employed
- These take time, so we want to minimize how many of these we do
- The Genetic Algorithm provides a framework for a refined global search

Concluding Remarks

- Finding all the low energy conformers for a molecule is hard
- In order to get accurate energies/structures, computationally expensive methods should be employed
- These take time, so we want to minimize how many of these we do
- The Genetic Algorithm provides a framework for a refined global search
- It shines when asked to find a host of low energy solutions

Concluding Remarks

- Finding all the low energy conformers for a molecule is hard
- In order to get accurate energies/structures, computationally expensive methods should be employed
- These take time, so we want to minimize how many of these we do
- The Genetic Algorithm provides a framework for a refined global search
- It shines when asked to find a host of low energy solutions
- GA wrapper can be interfaced with a variety of electronic structure packages(NWChem, ORCA) and is available under the GNU Lesser General Public License at <https://github.com/adrianasupady/fafoom>

Questions?

Empty Backup