



# Hartree-Fock Stability

## Applications to the Homogeneous Electron Gas

---

Evan Curtin

April 10, 2017

University of Illinois at Urbana-Champaign

# Table of contents

1. Background Information
2. Hartree-Fock Stability
3. Homogeneous Electron Gas
4. Results

## Background Information


---

# Levels of Hartree-Fock Theory

Method	Spinorbital	DoF	Eigenfunction of
Restricted	$\chi_j^\alpha(\vec{r}, \sigma) = \sum_{i=1}^N c_{ij} \phi_i(\vec{r}) \alpha(\sigma)$ $\chi_j^\beta(\vec{r}, \sigma) = \sum_{i=1}^N c_{ij} \phi_i(\vec{r}) \beta(\sigma)$	N/2	$\hat{S}^2, \hat{S}_z$
Unrestricted	$\chi_j^\alpha(\vec{r}, \sigma) = \sum_{i=1}^N c_{ij}^\alpha \phi_i(\vec{r}) \alpha(\sigma)$ $\chi_j^\beta(\vec{r}, \sigma) = \sum_{i=1}^N c_{ij}^\beta \phi_i(\vec{r}) \beta(\sigma)$	N	$\hat{S}_z$
General	$\chi_j(\vec{r}, \sigma) = \sum_{i=1}^N [c_{ij}^\alpha \phi_i(\vec{r}) \alpha(\sigma) + c_{ij}^\beta \phi_i(\vec{r}) \beta(\sigma)]$	2N	Neither

# Restricted Minimization


- Hartree-Fock SCF guarantees only stationary energy w.r.t. change in orbitals
- The solution may be a maximum, minimum or saddle point



# Restricted Minimization


- Hartree-Fock SCF guarantees only stationary energy w.r.t. change in orbitals
- The solution may be a maximum, minimum or saddle point

## Within the Constrained Space




# Restricted Minimization

- Restricted minima may correspond to minima in another dimension



# Restricted Minimization


- Restricted minima may correspond to minima in another dimension
- Restricted minima may correspond to maxima in another dimension





# Restricted Minimization

- Restricted minima may correspond to minima in another dimension
- Restricted minima may correspond to maxima in another dimension
- Restricted minima may be nonstationary



# Hartree-Fock Stability

---

# Hartree-Fock Stability

- Solving the HF equations guarantees only that the energy is **stationary**. The first order variation is therefore 0.

# Hartree-Fock Stability

- Solving the HF equations guarantees only that the energy is **stationary**. The first order variation is therefore 0.
- We need to know if this is indeed a minimum.

# Hartree-Fock Stability

- Solving the HF equations guarantees only that the energy is **stationary**. The first order variation is therefore 0.
- We need to know if this is indeed a minimum.
- We can determine this if we inspect the **second order variation** in the energy.

# Hartree-Fock Stability

- Solving the HF equations guarantees only that the energy is **stationary**. The first order variation is therefore 0.
- We need to know if this is indeed a minimum.
- We can determine this if we inspect the **second order variation** in the energy.
- Thouless<sup>1</sup> showed a physically motivated derivation using Time-Dependent Hartree-Fock theory (TDHF).

# Hartree-Fock Stability Conditions

- The RPA frequencies represent electronic oscillations about the ground state

# Hartree-Fock Stability Conditions

- The RPA frequencies represent electronic oscillations about the ground state
- Real frequencies are stable, imaginary frequencies are unstable



# Hartree-Fock Stability Conditions

- The RPA frequencies represent electronic oscillations about the ground state
- Real frequencies are stable, imaginary frequencies are unstable
- The RPA frequencies are imaginary when the **Orbital Hessian** (aka stability matrix, electronic Hessian) eigenvalues are **negative**

$$\begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{B}^* & \mathbf{A}^* \end{bmatrix} \begin{bmatrix} \mathbf{X} \\ \mathbf{Y} \end{bmatrix} = \omega \begin{bmatrix} \mathbf{X} \\ \mathbf{Y} \end{bmatrix} \quad (1)$$

# Hartree-Fock Stability Conditions

- The RPA frequencies represent electronic oscillations about the ground state
- Real frequencies are stable, imaginary frequencies are unstable
- The RPA frequencies are imaginary when the **Orbital Hessian** (aka stability matrix, electronic Hessian) eigenvalues are **negative**

$$\begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{B}^* & \mathbf{A}^* \end{bmatrix} \begin{bmatrix} \mathbf{X} \\ \mathbf{Y} \end{bmatrix} = \omega \begin{bmatrix} \mathbf{X} \\ \mathbf{Y} \end{bmatrix} \quad (1)$$

- Where

$$\begin{aligned} A_{ia,jb} &= \langle i^a | H - E_0 | j^b \rangle = (\epsilon_a - \epsilon_i) \delta_{ij} \delta_{ab} + \langle aj || ib \rangle \\ B_{ia,jb} &= \langle ij^a | H - E_0 | 0 \rangle = \langle ab || ij \rangle. \end{aligned} \quad (2)$$

# Hartree-Fock Stability Conditions

The eigenvalue equation can be factorized depending on symmetry.

Solution Type	Space Type					
	Real RHF	Complex RHF	Real UHF	Complex UHF	Real GHF	Complex GHF
Real RHF	${}^1\mathbf{A}' + {}^1\mathbf{B}'$	${}^1\mathbf{A}' - {}^1\mathbf{B}'$	${}^3\mathbf{A}' + {}^3\mathbf{B}'$	-	-	-
Complex RHF	-	${}^1\mathbf{H}'$	-	${}^3\mathbf{H}'$	-	-
Real UHF	-	-	$\mathbf{A}' + \mathbf{B}'$	$\mathbf{A}' - \mathbf{B}'$	$\mathbf{A}'' + \mathbf{B}''$	-
Complex UHF	-	-	-	$\mathbf{H}'$	-	$\mathbf{H}'$
Real GHF	-	-	-	-	$\mathbf{A} - \mathbf{B}$	$\mathbf{A} - \mathbf{B}$
Complex GHF	-	-	-	-	-	$\mathbf{H}$

Table reproduced from Seeger & Pople<sup>1</sup>

# Homogeneous Electron Gas

---

# Brief Overview

- Homogeneous Electron Gas (HEG) model, also known as Uniform Electron Gas or Jellium Model.
- Electrons in a box with "smeared" nuclei  $\rightarrow$  uniform positive background charge
- The total charge is constrained to be neutral,

$$V_{bg}(\mathbf{r}) = \sum_i \frac{-Ze^2}{|\mathbf{r} - \mathbf{R}_i|} \rightarrow -e^2 \int \frac{d\mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|}, \quad (3)$$

- and the background and coulomb terms cancel exactly,

$$V_{ee} = e^2 \int \frac{d\mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|}. \quad (4)$$

# Brief Overview

- The discretized solutions are given by,

$$\epsilon_{\vec{k}} = \frac{\hbar^2 k^2}{2m} - \sum_{\vec{k}'}^{|\vec{k}'| < k_f} \langle \vec{k}, \vec{k}' | \vec{k}', \vec{k} \rangle \quad (5)$$

- Where the two electron integral is given by

$$\langle \vec{k}, \vec{k}' | \vec{k}'', \vec{k}''' \rangle \stackrel{2D, 3D}{=} \begin{cases} \frac{\pi}{V} \frac{2^{D-1}}{|\vec{k} - \vec{k}''|^{D-1}} & \vec{k}''' = \vec{k} + \vec{k}' - \vec{k}'' \\ 0 & \text{else} \end{cases}$$
$$\langle k, k' | k'', k''' \rangle \stackrel{1D}{=} \begin{cases} e^{|k-k''|^2 a^2} \text{Ei}(-|k-k''|^2 a^2) ; & k''' = k + k' - k'' \\ 0 ; & \text{else} \end{cases} \quad (6)$$

# Project Goal

- Can we use the HF-Stability analysis to determine where the HEG is unstable?

# Project Goal

- Can we use the HF-Stability analysis to determine where the HEG is unstable?
- Will this predict the known tendency of crystallization at low density?



# Project Goal

- Can we use the HF-Stability analysis to determine where the HEG is unstable?
- Will this predict the known tendency of crystallization at low density?
- It is known that the RHF-GHF instability persists at all densities for the HEG<sup>1</sup>.

# Project Goal

- Can we use the HF-Stability analysis to determine where the HEG is unstable?
- Will this predict the known tendency of crystallization at low density?
- It is known that the RHF-GHF instability persists at all densities for the HEG<sup>1</sup>.
- Can we show this numerically, and discriminate between RHF-UHF and RHF-GHF instability?

# Project Goal


- Can we use the HF-Stability analysis to determine where the HEG is unstable?
- Will this predict the known tendency of crystallization at low density?
- It is known that the RHF-GHF instability persists at all densities for the HEG<sup>1</sup>.
- Can we show this numerically, and discriminate between RHF-UHF and RHF-GHF instability?
- (Future) Can we combine this with to improve the efficacy of correlation theories (CC, MBPT)?

## Results

---


# Orbital Energies

- $Nk = 57$  reproduces the orbital energies reasonably well
- Worst towards  $\Gamma$ , better for higher  $|\vec{k}|$




# Matrix Diagonals

- Spectrum is Dense




# Matrix Size Scaling

- Scales as  $2N_{exc}$
- $N_{occ}$  and  $N_{vir}$  both scale with  $N_{kpoints}^D$
- Use an iterative subspace method
- Jacobi-Davidson(-Liu)



# Matrix-Vector Scaling


- Scales as  $N_{exc} \times N_{occ}$   
due to momentum  
conservation





# Davidson Scaling


- Davidson is Asymptotically quadratic.
- Full diagonalization is almost cubic.
- Matrix multiplication is order<sup>1</sup>  $\log_2(7) \approx 2.807$



1) <http://mathworld.wolfram.com/StrassenFormulas.html>



# Efficacy of Davidson's Algorithm

- Reproduces Exact result to machine precision in all test cases.
- Odd spikes are due to approximating circle by squares





# Dependence on $r_s$

Crossover from stable to unstable agrees with previous results.



# Dependence on $r_s$

3D data is currently inconclusive.



**Questions?**