

Hartree-Fock Stability Theory and its Relation to Correlation Theories

Evan Curtin

I. BACKGROUND INFORMATION AND ACCOMPLISHMENTS

A. Hartree-Fock Stability

Hartree-Fock (HF) Theory has been the foundation for *ab initio* electronic structure theory throughout its history. An often overlooked aspect of the theory is that there are many solutions to the HF equations for a given system, but being a solution to the Hartree-Fock equation ensures only that the solution is stationary with respect to the determined orbitals. If the solution is indeed a minimum, it is called "stable", while if there is any displacement in the electronic structure which reduces the energy, the solution is "unstable". A method for determining the stability of a Hartree-Fock solution was proposed by Thouless in 1960[1]. The condition was rederived into the expression familiar to quantum chemists by Cizek and Paldus in 1967[2]. Furthermore, the stability equations factorize depending on the symmetry of the Hartree-Fock eigenfunctions. To this end, Seeger and Pople outlined a hierarchical approach to systematically evaluate the stability of HF states in the restricted, unrestricted and generalized Hartree-Fock procedures[3]. Recently, the method has been used to aid the in search for the lowest energy Unrestricted Hartree-Fock (UHF) solutions in molecules, as well as the General Hartree-Fock (GHF) solutions in geometrically frustrated hydrogen rings which cannot conform even to the UHF scheme [4][5].

The usual method of determining stability of a Hartree-Fock solution is to look for a negative eigenvalue the Electronic Hessian,

$$\mathbf{H} = \begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{B}^* & \mathbf{A}^* \end{bmatrix}, \quad (1)$$

where the matrices \mathbf{A} and \mathbf{B} have dimension $N_{occupied} \times N_{virtual}$ with elements given by

$$A_{i \rightarrow a, j \rightarrow b} = (\epsilon_a - \epsilon_i) \delta_{ij} \delta_{ab} + \langle aj || ib \rangle, \quad (2)$$

$$B_{i \rightarrow a, j \rightarrow b} = \langle ab || ij \rangle. \quad (3)$$

If \mathbf{H} has a negative eigenvalue, this indicates an instability. For a paramagnetic HF solution, the equations factorize into the matrices known to chemists as the singlet and triplet instability matrices [6][3],

$${}^1A'_{i \rightarrow a, j \rightarrow b} = (\epsilon_a - \epsilon_i) \delta_{ij} \delta_{ab} + 2 \langle aj|ib \rangle - \langle aj|bi \rangle \quad (4a)$$

$${}^3A'_{i \rightarrow a, j \rightarrow b} = (\epsilon_a - \epsilon_i) \delta_{ij} \delta_{ab} - \langle aj|bi \rangle \quad (4b)$$

$${}^1B'_{i \rightarrow a, j \rightarrow b} = 2 \langle ab|ij \rangle - \langle ab|ji \rangle \quad (4c)$$

$${}^3B'_{i \rightarrow a, j \rightarrow b} = - \langle ab|ji \rangle \quad (4d)$$

where the 1, 3 denote singlet and triplet instabilities, respectively. The lowest eigenvalue of both of these matrices will reveal the stability of the RHF solution. If they are both positive, the solution is stable with respect to symmetry breaking, while negative eigenvalues in either indicate instability with respect to either triplet or singlet perturbations (or both). An eigenvalue of 0 does not indicate instability, and this case is discussed in detail by Cui et al [7].

B. Hartree-Fock Stability of the Homogeneous Electron Gas

The Homogeneous Electron Gas (HEG) model of metallic systems is a convenient starting point for this study. The Energies of various phases thereof have been computed to great accuracy [8], while plane wave solutions to the Hartree-Fock equations can be demonstrated analytically [9]. Furthermore, within the Hartree-Fock approximation, it has been known for half a century that broken symmetry solutions are always lower in energy compared to their paramagnetic counterparts [10]. More recently, phase diagrams have been determined for the HEG in 2 and 3 dimensions[11][12][13]. In all cases the phase diagrams are made by computing the energies of the polarized and unpolarized states, and comparing them. This approach necessitates that the form of the solutions is known ahead of time.

The present work focuses on directly calculating the Hartree-Fock stability of the paramagnetic homogeneous electron gas as a function of electron density. In this way it is possible to map out where the symmetry broken Hartree-Fock reference is needed. To date, an optimal form of the Hartree-Fock solution to the HEG remains unknown, even though exact energy (including correlation) is essentially known. Therefore determining a lower energy HF solution can yield insight into the effects of correlation of this fundamental physical model.

C. Method

Starting with the Jellium Hamiltonian in atomic units,

$$\hat{H} = \sum_i \frac{\hat{p}_i^2}{2} + \frac{1}{2} \sum_{i < j} \frac{1}{|\vec{r}_i - \vec{r}_j|}, \quad (5)$$

one solution to the Hartree-Fock equations in D dimensions are plane waves of the form

$$\phi_{\vec{k}} = \frac{1}{\sqrt{L^D}} e^{i\vec{k}\vec{x}}, \quad (6)$$

where L is the length of one side of the direct lattice with cubic symmetry. Using this as the basis for the rest of the analysis, the two electron repulsion integrals are analytic and well-behaved in two and three dimensions [11][14]

$$\langle \vec{k}, \vec{k}' | \vec{k}'', \vec{k}''' \rangle \stackrel{2D, 3D}{=} \begin{cases} \frac{\pi}{\Omega} \frac{2^{D-1}}{|\vec{k} - \vec{k}''|^{D-1}} & \vec{k}''' = \vec{k} + \vec{k}' - \vec{k}'' + n\vec{G} \text{ and } |\vec{k} - \vec{k}''| \neq 0 \\ 0 & \text{else,} \end{cases} \quad (7)$$

where Ω is the direct lattice volume and $n\vec{G}$ is any integer times a reciprocal lattice vector of the system. The orbital energies are

$$\epsilon_{\vec{k}, \sigma} = \frac{\hbar^2 \vec{k}^2}{2m} - \sum_{\vec{k}}^{|\vec{k}| < k_f} n_{\vec{k}\sigma} \langle \vec{k}, \vec{k}' | \vec{k}', \vec{k} \rangle, \quad (8)$$

where $n_{\vec{k}\sigma}$ is the occupation number of the state with momentum \vec{k} and spin σ [14].

D. Results

Using the equations for energies and two electron integrals, the stability analysis was performed on the paramagnetic HEG model as a function of electron density (or equivalently, r_s). The naive approach of explicitly constructing and diagonalizing the orbital hessian is intractable both in terms of memory and calculation time requirements. Since the instability condition is that the any eigenvalue is negative, it suffices to calculate the sign of only the lowest eigenvalue. This is a task well suited for an iterative subspace solver. To this end a parallel version of the Jacobi-Davidson algorithm was implemented by representing the matrix as a matrix-vector product (made possible by the SLEPc library[15]). This allowed for two necessary optimizations. First, the algorithmic change from full diagonalization to a subspace algorithm reduces the memory requirements from $O(N^2)$ to $O(N)$ and the computational scaling from $O(N^3)$ to $O(N^2)$ (See Fig 1). Secondly, the parallelization of the algorithm allows the program to be run on the Blue Waters supercomputer with about 80% strong parallel scaling efficiency (Fig 2).




FIG. 1: The asymptotic scaling of both algorithms behaves as expected.




FIG. 2: The speedup of the entire calculation as the number of computer cores is sufficiently linear.

These optimizations allow the calculation to be performed with enough grid points to converge the resulting "transition r_s ", the Wigner-Seitz radius at which the HF solution transitions from stable to unstable.




FIG. 3: The Lowest eigenvalues of the triplet and singlet instability matrices in 1, 2 and 3 dimensions.

The transition r_s'' occurs at $r_s = 1$ in 2D and $r_s = 3$ in 3D. These results are consistent with those seen by other methods[16][12]. On the other hand, the transition r_s for one dimension tends towards zero and stays quite close as the number of gridpoints increases. Indeed, numerical issues appear at very high density (low r_s) due to divergence of the two electron integrals in the asymptotic limit. With this in mind, it is clear at least that the instability in the 1D case persists even at very high densities (potentially all finite densities), which is to be expected from Overhauser's proof of instability at all values of r_s in 1D for the delta function potential [10].




FIG. 4: The Lowest eigenvalues of the triplet and singlet instability matrices in 1, 2 and 3 dimensions.

II. GOALS AND RELEVANCE OF PROPOSED RESEARCH

The goal of electronic structure theory is to solve the electronic Schrödinger equation for a molecule or solid state system. The prevailing approach in *ab initio* quantum chemistry is to first compute a self-consistent single-determinant Hartree-Fock (HF) solution which will serve as a reference for many-body perturbative (MP) or coupled-cluster (CC) approaches. Implicit in these methods is the assumption that the reference wave function will be sufficiently close to the exact solution of the Schrödinger equation to converge to the correct answer in a computationally feasible manner. Both the MP and CC approaches have the attractive quality that they are systematically improvable, i.e. more computational power can be used on a problem to increase the accuracy of the resulting solution. These so-called post-Hartree-Fock methods are able to recover more of the total electronic energy by accounting for the *correlation energy* in the system. Traditionally, the correlation energy is defined by the difference between the exact energy and the Hartree-Fock energy [17]

$$E_{corr} = E_{exact} - E_{HF} \quad (9)$$

In many cases the correlation energy is relatively small and the Hartree-Fock reference captures a significant portion of the exact total energy. In such cases, the prevailing approach is usually successful and has only implementation issues for larger systems. On the other hand there are systems which are not well treated by this approach, which are often called "strongly correlated". Examples of such systems are those with degenerate or near-degenerate HOMO-LUMO gaps, metallic solids or molecules close to breaking bonds. Oftentimes for such systems the traditional Hartree-Fock solutions are not even qualitatively correct, implying inadequacies in either the single-reference or mean-field approach.

It is often neglected, however, that the usual way for computing a Hartree-Fock solution in practice is to use a restricted form of the single determinantal wavefunction. In the Restricted Hartree-Fock (RHF), the wave function is restricted to be an eigenfunction of the \hat{S}^2 and \hat{S}_z operators, while in the Unrestricted variant (UHF), the wave function need only be an eigenfunction of the \hat{S}_z . There is yet another lesser-known variation of the theory, known as General Hartree-Fock (GHF), which assumes no spin-symmetries in the wave function. This allows for greater variational flexibility at the cost of "broken symmetry". Together these three levels of Hartree-Fock theory form a hierarchy of increasing parameter space within which to minimize the total energy. Thus the term "Hartree-Fock Energy" is inherently ambiguous; this fact calling into question commonly held ideas about correlation and Hartree-Fock theory. Take for example the dissociation curve of H_2 in the RHF and UHF approximations, compared with the exact solution obtained by full configuration-interaction in the complete basis set limit.




FIG. 5: Dissociation curve in the RHF and UHF approximations, compared to the exact limit. The difference in correlation energies as determined by each reference illustrates the ambiguity of that quantity. In the large dissociation limit, the correlation energy vanishes.

The correlation energy for the RHF reference case is dramatically larger than that for the UHF reference. In other words, *the correlation energy is, by definition, inherently coupled to the Hartree-Fock reference*. The given example is well understood, but it brings to mind a more general idea that there may be more cases where apparently large correlation energies are due to simply using an inadequate level of Hartree-Fock theory. Scuseria and coworkers have leveraged this idea to create new coupled cluster theories while preserving the symmetry of the reference [18].

While the exact wave function will have the same symmetry as the exact Hamiltonian (to be exact, the wave function will be an eigenfunction of all symmetry operators that commute with the Hamiltonian), it is important to remember that the Hartree-Fock Hamiltonian is not exact, and we may have something to gain by taking the variational principle at face value; lower energy is always better. For predicting certain observables this is not the case, but for energies alone we are free to maintain this mindset. Therefore the overarching theme of the proposed work is that **many examples of strong correlation may be misleading, and in order to find out we have to find lower energy Hartree-Fock solutions, even at the cost of broken symmetry**.

For some time an approach to find lower energy HF solutions has been known to follow the instability eigenvectors down in energy to find lower energy solutions [3]. This approach will, by construction, always lead to a lower energy HF solution than the starting point. However, it is important to remember that the instability condition is a local phenomena, and the lower energy solution is not itself guaranteed to be the *lowest* energy solution. The standard

approach still suffers from the global optimization problem: whether a solution is a local or global minimum can only be known by comparing it to the other minima of the surface. In fact, the problem of finding the lowest energy HF solution has a direct correspondence to the problem of finding minimum energy points on a potential energy surface of atomic coordinates. It is therefore proposed to adopt approaches from that field to apply to the current problem. An attractive approach is the Global Reaction Route Mapping (GRRM) method with its Anharmonic Downward Distortion (ADD) algorithm [19][20]. In essence, the algorithm leverages the idea that the presence of a local minimum influences the surrounding energy landscape (in a smooth surface). The minimum will lower the energy pathways leading towards it, and softens the ascent of energy around a transition state. This softening of the PES can be seen by increased anharmonicity in the PES's Taylor expansion. Thus a waypoint from one minimum to another can be determined by finding the direction with the highest anharmonicity, and these can be followed to find new solutions. By adopting this approach to use molecular orbital expansion coefficients in place of normal mode coordinates, many minima of the Hartree-Fock electronic energy surface can be found.

III. REFERENCES

-
- [1] D. Thouless, Nuclear Physics **21**, 225 (1960), ISSN 00295582, URL <http://linkinghub.elsevier.com/retrieve/pii/0029558260900481>.
 - [2] J. Cizek and J. Paldus, The Journal of Chemical Physics **47**, 3976 (1967), ISSN 00219606, URL <http://scitation.aip.org/content/aip/journal/jcp/47/10/10.1063/1.1701562>.
 - [3] R. Seeger and J. A. Pople, The Journal of Chemical Physics **66**, 3045 (1977), ISSN 00219606, URL <http://scitation.aip.org/content/aip/journal/jcp/66/7/10.1063/1.434318>.
 - [4] P. Pulay and Z. Tóth, Preprint **164102**, 1 (2016), ISSN 0021-9606, URL <http://dx.doi.org/10.1063/1.4964903>.
 - [5] J. J. Goings, F. Ding, M. J. Frisch, and X. Li, The Journal of Chemical Physics **142**, 154109 (2015), ISSN 0021-9606, URL <http://dx.doi.org/10.1063/1.4918561><http://scitation.aip.org/content/aip/journal/jcp/142/15/10.1063/1.4918561>.
 - [6] T. H. Dunning, The Journal of Chemical Physics **47**, 1735 (1967), ISSN 00219606, URL <http://scitation.aip.org/content/aip/journal/jcp/47/5/10.1063/1.1712158>.
 - [7] Y. Cui, I. W. Bulik, C. A. Jimenez-Hoyos, T. M. Henderson, and G. E. Scuseria, The Journal of Chemical Physics **139**, 154107 (2013), ISSN 00219606, arXiv:1310.0942v1, URL <http://scitation.aip.org/content/aip/journal/jcp/139/15/154107>.

10.1063/1.4824905.

- [8] D. M. Ceperley and B. J. Alder, *Physical Review Letters* **45**, 566 (1980), ISSN 0031-9007, URL <http://link.aps.org/doi/10.1103/PhysRevLett.45.566>.
- [9] P. Phillips, *Advanced Solid State Physics* (Cambridge University Press, Cambridge, 2012), 2nd ed., ISBN 978-0-521-19490-7.
- [10] A. W. Overhauser, *Physical Review* **128**, 1437 (1962), ISSN 0031-899X, URL <http://link.aps.org/doi/10.1103/PhysRev.128.1437>.
- [11] F. Delyon, M. Duneau, B. Bernu, and M. Holzmann, pp. 1–12 (2008), 0807.0770, URL <http://arxiv.org/abs/0807.0770>.
- [12] B. Bernu, F. Delyon, M. Holzmann, and L. Baguet, *Physical Review B* **84**, 115115 (2011), ISSN 1098-0121, URL <http://link.aps.org/doi/10.1103/PhysRevB.84.115115>.
- [13] L. Baguet, F. Delyon, B. Bernu, and M. Holzmann, *Physical Review Letters* **111**, 166402 (2013), ISSN 0031-9007, URL <http://link.aps.org/doi/10.1103/PhysRevLett.111.166402>.
- [14] G. Guillian and G. Vignale, *Quantum Theory of the Electron Liquid* (Cambridge University Press, Cambridge, 2005), ISBN 978-0-521-82112-6.
- [15] V. Hernandez, J. E. Roman, and V. Vidal, *ACM Transactions on Mathematical Software* **31**, 351 (2005), ISSN 00983500, URL <http://portal.acm.org/citation.cfm?doid=1089014.1089019>.
- [16] L. Baguet, F. Delyon, B. Bernu, and M. Holzmann, *Physical Review B* **90**, 165131 (2014), ISSN 1098-0121, arXiv:1404.7652v1, URL <http://link.aps.org/doi/10.1103/PhysRevB.90.165131>.
- [17] I. Shavitt and R. J. Bartlett, *Many-Body Methods in Chemistry and Physics* (Cambridge University Press, 2009), ISBN 978-0-521-81832-2.
- [18] J. A. Gomez, T. M. Henderson, and G. E. Scuseria, *The Journal of Chemical Physics* **144**, 244117 (2016), ISSN 0021-9606, URL <http://scitation.aip.org/content/aip/journal/jcp/144/24/10.1063/1.4954891>.
- [19] K. Ohno and S. Maeda, *Journal of Physical Chemistry A* **110**, 8933 (2006), ISSN 10895639.
- [20] K. Ohno, *Chemical Record* **16**, 2198 (2016), ISSN 15280691.