


# TABLE OF CONTENTS





# HIGHER ORDER DERIVATIVES

Evaluate  $\frac{d}{dx} \left[ \frac{d}{dx} [x^5 - 2x^2 + 3] \right]$

# HIGHER ORDER DERIVATIVES

Evaluate  $\frac{d}{dx} \left[ \frac{d}{dx} [x^5 - 2x^2 + 3] \right]$

$$\frac{d}{dx} [x^5 - 2x^2 + 3] = 5x^4 - 4x$$

# HIGHER ORDER DERIVATIVES

Evaluate  $\frac{d}{dx} \left[ \frac{d}{dx} [x^5 - 2x^2 + 3] \right]$

$$\frac{d}{dx} [x^5 - 2x^2 + 3] = 5x^4 - 4x$$

$$\begin{aligned} \frac{d}{dx} \left[ \frac{d}{dx} [x^5 - 2x^2 + 3] \right] &= \frac{d}{dx} [5x^4 - 4x] \\ &= 20x^3 - 4 \end{aligned}$$

# HIGHER ORDER DERIVATIVES

Evaluate  $\frac{d}{dx} \left[ \frac{d}{dx} [x^5 - 2x^2 + 3] \right]$

$$\begin{aligned} \frac{d}{dx} [x^5 - 2x^2 + 3] &= 5x^4 - 4x \\ \frac{d}{dx} \left[ \frac{d}{dx} [x^5 - 2x^2 + 3] \right] &= \frac{d}{dx} [5x^4 - 4x] \\ &= 20x^3 - 4 \end{aligned}$$

## Notation 2.14.1

The derivative of a derivative is called the **second derivative**, written

$$f''(x) \quad \text{or} \quad \frac{d^2y}{dx^2}(x)$$

# HIGHER ORDER DERIVATIVES

Evaluate  $\frac{d}{dx} \left[ \frac{d}{dx} [x^5 - 2x^2 + 3] \right]$

$$\begin{aligned} \frac{d}{dx} [x^5 - 2x^2 + 3] &= 5x^4 - 4x \\ \frac{d}{dx} \left[ \frac{d}{dx} [x^5 - 2x^2 + 3] \right] &= \frac{d}{dx} [5x^4 - 4x] \\ &= 20x^3 - 4 \end{aligned}$$

## Notation 2.14.1

The derivative of a derivative is called the **second derivative**, written

$$f''(x) \quad \text{or} \quad \frac{d^2y}{dx^2}(x)$$

Similarly, the derivative of a second derivative is a third derivative, etc.

## Notation 2.14.1

- ▶  $f''(x)$  and  $f^{(2)}(x)$  and  $\frac{d^2f}{dx^2}(x)$  all mean  $\frac{d}{dx} \left( \frac{d}{dx} f(x) \right)$
- ▶  $f'''(x)$  and  $f^{(3)}(x)$  and  $\frac{d^3f}{dx^3}(x)$  all mean  $\frac{d}{dx} \left( \frac{d}{dx} \left( \frac{d}{dx} f(x) \right) \right)$
- ▶  $f^{(4)}(x)$  and  $\frac{d^4f}{dx^4}(x)$  both mean  $\frac{d}{dx} \left( \frac{d}{dx} \left( \frac{d}{dx} \left( \frac{d}{dx} f(x) \right) \right) \right)$
- ▶ and so on.



## TYPICAL EXAMPLE: ACCELERATION

- ▶ Velocity: rate of change of position

# TYPICAL EXAMPLE: ACCELERATION

- ▶ Velocity: rate of change of position
- ▶ Acceleration: rate of change of velocity.

## TYPICAL EXAMPLE: ACCELERATION

- ▶ Velocity: rate of change of position
- ▶ Acceleration: rate of change of velocity.

The position of an object at time  $t$  is given by  $s(t) = t(5 - t)$ . *Time is measured in seconds, and position is measured in metres.*


1. Sketch the graph giving the position of the object.
2. What is the velocity of the object when  $t = 1$ ? Include units.
3. What is the acceleration of the object when  $t = 1$ ? Include units.

# TYPICAL EXAMPLE: ACCELERATION

- ▶ Velocity: rate of change of position
- ▶ Acceleration: rate of change of velocity.

The position of an object at time  $t$  is given by  $s(t) = t(5 - t)$ . *Time is measured in seconds, and position is measured in metres.*

1. Sketch the graph giving the position of the object.
2. What is the velocity of the object when  $t = 1$ ? Include units.
3. What is the acceleration of the object when  $t = 1$ ? Include units.



# TYPICAL EXAMPLE: ACCELERATION

- ▶ Velocity: rate of change of position
- ▶ Acceleration: rate of change of velocity.

The position of an object at time  $t$  is given by  $s(t) = t(5 - t)$ . *Time is measured in seconds, and position is measured in metres.*

1. Sketch the graph giving the position of the object.
2. What is the velocity of the object when  $t = 1$ ? Include units.
3. What is the acceleration of the object when  $t = 1$ ? Include units.

$$s(t) = t(5 - t) = 5t - t^2$$

$$s'(t) = 5 - 2t$$

$$s'(1) = 5 - 2(1) = 3 = \text{vel}$$

$$\text{Units of velocity: } \frac{\Delta s}{\Delta t} = \frac{m}{s}$$

$$\text{Units of acceleration: } \frac{\Delta s'}{\Delta t} = \frac{m/s}{s} = \frac{m}{s^2}$$

## TYPICAL EXAMPLE: ACCELERATION

- ▶ Velocity: rate of change of position
- ▶ Acceleration: rate of change of velocity.

The position of an object at time  $t$  is given by  $s(t) = t(5 - t)$ . *Time is measured in seconds, and position is measured in metres.*

1. Sketch the graph giving the position of the object.
2. What is the velocity of the object when  $t = 1$ ? Include units.
3. What is the acceleration of the object when  $t = 1$ ? Include units.

$$s(t) = t(5 - t) = 5t - t^2$$

$$s'(t) = 5 - 2t$$

$$s'(1) = 5 - 2(1) = 3 = \text{vel}$$

$$s''(t) = -2$$

$$-2 = \text{acc}$$

$$\text{Units of velocity: } \frac{\Delta s}{\Delta t} = \frac{m}{s}$$

$$\text{Units of acceleration: } \frac{\Delta s'}{\Delta t} = \frac{m/s}{s} = \frac{m}{s^2}$$

## CONCEPT CHECK

**True or False:** If  $f'(1) = 18$ , then  $f''(1) = 0$ ,  
since the  $\frac{d}{dx}\{18\} = 0$ .

## CONCEPT CHECK

**True or False:** If  $f'(1) = 18$ , then  $f''(1) = 0$ ,  
since the  $\frac{d}{dx}\{18\} = 0$ .

False: for example,  $f(x) = 9x^2$ .



## CONCEPT CHECK

**True or False:** If  $f'(1) = 18$ , then  $f''(1) = 0$ , since the  $\frac{d}{dx}\{18\} = 0$ .

**False:** for example,  $f(x) = 9x^2$ .

**Which of the following is always true of a QUADRATIC polynomial  $f(x)$ ?**

- A.  $f(0) = 0$
- B.  $f'(0) = 0$
- C.  $f''(0) = 0$
- D.  $f'''(0) = 0$
- E.  $f^{(4)}(0) = 0$

## CONCEPT CHECK

**True or False:** If  $f'(1) = 18$ , then  $f''(1) = 0$ , since the  $\frac{d}{dx}\{18\} = 0$ .

**False:** for example,  $f(x) = 9x^2$ .

**Which of the following is always true of a QUADRATIC polynomial  $f(x)$ ?**

- A.  $f(0) = 0$      $f(x) = ax^2 + bx + c$
- B.  $f'(0) = 0$      $f'(x) = 2ax + b$
- C.  $f''(0) = 0$      $f''(x) = 2a$
- D.  $f'''(0) = 0$      $f'''(x) = 0$  ✓
- E.  $f^{(4)}(0) = 0$      $f^{(4)}(x) = 0$  ✓

## CONCEPT CHECK

**True or False:** If  $f'(1) = 18$ , then  $f''(1) = 0$ ,  
since the  $\frac{d}{dx}\{18\} = 0$ .

**False:** for example,  $f(x) = 9x^2$ .

**Which of the following is  
always true of a QUADRATIC  
polynomial  $f(x)$ ?**

- A.  $f(0) = 0$      $f(x) = ax^2 + bx + c$
- B.  $f'(0) = 0$      $f'(x) = 2ax + b$
- C.  $f''(0) = 0$      $f''(x) = 2a$
- D.  $f'''(0) = 0$      $f'''(x) = 0$  ✓
- E.  $f^{(4)}(0) = 0$      $f^{(4)}(x) = 0$  ✓

**Which of the following is  
always true of a CUBIC  
polynomial  $f(x)$ ?**

- A.  $f(0) = 0$
- B.  $f'(0) = 0$
- C.  $f''(0) = 0$
- D.  $f'''(0) = 0$
- E.  $f^{(4)}(0) = 0$

## CONCEPT CHECK

**True or False:** If  $f'(1) = 18$ , then  $f''(1) = 0$ ,  
since the  $\frac{d}{dx}\{18\} = 0$ .

**False:** for example,  $f(x) = 9x^2$ .

**Which of the following is  
always true of a QUADRATIC  
polynomial  $f(x)$ ?**

- A.  $f(0) = 0$      $f(x) = ax^2 + bx + c$
- B.  $f'(0) = 0$      $f'(x) = 2ax + b$
- C.  $f''(0) = 0$      $f''(x) = 2a$
- D.  $f'''(0) = 0$      $f'''(x) = 0$  ✓
- E.  $f^{(4)}(0) = 0$      $f^{(4)}(x) = 0$  ✓

**Which of the following is  
always true of a CUBIC  
polynomial  $f(x)$ ?**

- A.  $f(0) = 0$      $f(x) = ax^3 + bx^2 + cx + d$
- B.  $f'(0) = 0$      $f'(x) = 3ax^2 + 2bx + c$
- C.  $f''(0) = 0$      $f''(x) = 6ax + 2b$
- D.  $f'''(0) = 0$      $f'''(x) = 6a$
- E.  $f^{(4)}(0) = 0$      $f^{(4)}(x) = 0$  ✓



# IMPLICIT DIFFERENTIATION

Suppose  $y(x)$  is a function such that

$$y(x) = y^3x + x^2 - 1$$

Find  $y''(x)$  at the point  $(-2, 1)$ .

# IMPLICIT DIFFERENTIATION

Suppose  $y(x)$  is a function such that

$$y(x) = y^3x + x^2 - 1$$

Find  $y''(x)$  at the point  $(-2, 1)$ . We start by differentiating both sides of the function. Remember that  $y$  is a function, not a variable.

$$y(x) = y(x)^3x + x^2 + 1$$

$$\frac{dy}{dx}(x) \stackrel{\text{prod}}{=} y(x)^3 + 3xy(x)^2 \frac{dy}{dx}(x) + 2x \quad (*)$$

Let's differentiate both sides again. Remember we have a rule for the product of three functions.

$$\frac{d^2y}{dx^2} = 3y^2 \frac{dy}{dx} + 3 \left( y^2 \frac{dy}{dx} + x \cdot 2y \frac{dy}{dx} \cdot \frac{dy}{dx} + xy^2 \frac{d^2y}{dx^2} \right) + 2 \quad (**)$$

When  $x = -2$  and  $y = 1$ , using (\*), we find

$$\left. \frac{dy}{dx} \right|_{(-2,1)} = 1^3 + 3(-2)(1^2) \left. \frac{dy}{dx} \right|_{(-2,1)} + 2(-2) = -3 - 6 \left. \frac{dy}{dx} \right|_{(-2,1)}$$

$$\left. \frac{dy}{dx} \right|_{(-2,1)} = -\frac{3}{7}$$

We set  $x = -2$ ,  $y = 1$ , and  $\frac{dy}{dx} = -\frac{3}{7}$  in equation (\*\*). Now by  $\frac{d^2y}{dx^2}$ , we actually mean

$\left. \frac{d^2y}{dx^2} \right|_{(-2,1)}$ , but to avoid clutter we don't write it that way until the end.

$$\frac{d^2y}{dx^2} = 3(1) \left( -\frac{3}{7} \right) + 3 \left( (1) \left( -\frac{3}{7} \right) + (-2) \cdot 2(1) \left( -\frac{3}{7} \right) \cdot \left( -\frac{3}{7} \right) + (-2)(1) \frac{d^2y}{dx^2} \right) + 2$$

$$= -\frac{9}{7} + 3 \left( -\frac{3}{7} - \frac{36}{49} - 2 \frac{d^2y}{dx^2} \right) + 2$$

$$= \left( 2 - \frac{18}{7} - \frac{108}{49} \right) - 6 \frac{d^2y}{dx^2}$$

$$7 \frac{d^2y}{dx^2} = -\frac{136}{7^2}$$

$$\left. \frac{d^2y}{dx^2} \right|_{(-2,1)} = -\frac{136}{7^3}$$

## Included Work



'Recursive Blanket Flower' by [Alexandre Duret-Lutz](#) is licensed under [CC BY-NC-SA 2.0](#) (accessed 20 July 2021), 2