
TABLE OF CONTENTS



The derivative of f(x) at a, written f'(a), is the instantaneous rate of change of f(x) when x = a.



The derivative of f(x) at a, written f'(a), is the instantaneous rate of change of f(x) when x = a.

Suppose P(t) gives the number of people in the world at t minutes past midnight, January 1, 2012. Suppose further that P'(0) = 156. How do you interpret P'(0) = 156?



The derivative of f(x) at a, written f'(a), is the instantaneous rate of change of f(x) when x = a.

Suppose P(t) gives the number of people in the world at t minutes past midnight, January 1, 2012. Suppose further that P'(0) = 156. How do you interpret P'(0) = 156?

At midnight of January 1, 2012, the world population was increasing at a rate of 156 people each minute



The derivative of f(x) at a, written f'(a), is the instantaneous rate of change of f(x) when x = a.

Suppose P(n) gives the total profit, in dollars, earned by selling n widgets. How do you interpret P'(100)?



The derivative of f(x) at a, written f'(a), is the instantaneous rate of change of f(x) when x = a.

Suppose P(n) gives the total profit, in dollars, earned by selling n widgets. How do you interpret P'(100)?

How fast your profit is increasing as you sell more widgets, measured in dollars per widget, at the time you sell Widget #100. So, roughly the profit earned from the sale of the 101st widget.



The derivative of f(x) at a, written f'(a), is the instantaneous rate of change of f(x) when x = a.

Suppose h(t) gives the height of a rocket t seconds after liftoff. What is the interpretation of h'(t)?



The derivative of f(x) at a, written f'(a), is the instantaneous rate of change of f(x) when x = a.

Suppose h(t) gives the height of a rocket t seconds after liftoff. What is the interpretation of h'(t)?

The speed at which the rocket is rising at time t.



The derivative of f(x) at a, written f'(a), is the instantaneous rate of change of f(x) when x = a.

Suppose M(t) is the number of molecules of a chemical in a test tube t seconds after a reaction starts. Interpret M'(t).



The derivative of f(x) at a, written f'(a), is the instantaneous rate of change of f(x) when x = a.

Suppose M(t) is the number of molecules of a chemical in a test tube t seconds after a reaction starts. Interpret M'(t).

The rate (measured in molecules per second) at which the number of molecules of a certain type is changing. Roughly, how many molecules of that type are being added (or taken away, if negative) per second at time t.



The derivative of f(x) at a, written f'(a), is the instantaneous rate of change of f(x) when x = a.

Suppose G(w) gives the diameter in millimetres of steel wire needed to safely support a load of w kg. Suppose further that G'(100) = 0.01. How do you interpret G'(100) = 0.01?

The derivative of f(x) at a, written f'(a), is the instantaneous rate of change of f(x) when x = a.

Suppose G(w) gives the diameter in millimetres of steel wire needed to safely support a load of w kg. Suppose further that G'(100) = 0.01. How do you interpret G'(100) = 0.01?

When your load is about 100 kg, you need to increase the diameter of your wire by about 0.01 mm for each kg increase in your load.



A paper¹ on the impacts of various factors in average life expectancy contains the following:

The only statistically significant variable in the model is physician density. The coefficient for this variable 20.67 indicating that a one unit increase in physician density leads to a 20.67 unit increase in life expectancy. This variable is also statistically significant at the 1% level demonstrating that this variable is very strongly and positively correlated with quality of healthcare received. This denotes that access to healthcare is very impactful in terms of increasing the quality of health in the country.

¹Natasha Deshpande, Anoosha Kumar, Rohini Ramaswami, *The Effect of National Healthcare Expenditure on Life Expectancy*, page 12.


Remark: physician density is measured as number of doctors per 1000 members of the population.

If L(p) is the average life expectancy in an area with a density p of physicians, write the statement as a derivative: "a one unit increase in physician density leads to a 20.67 unit increase in life expectancy."


If L(p) is the average life expectancy in an area with a density p of physicians, write the statement as a derivative: "a one unit increase in physician density leads to a 20.67 unit increase in life expectancy."

$$L'(p) = 20.67$$


The tangent line to f(x) at a has slope f'(a) and passes through the point (a, f(a)).



The tangent line to f(x) at a has slope f'(a) and passes through the point (a, f(a)).



The tangent line to f(x) at a has slope f'(a) and passes through the point (a, f(a)).



Slope of the tangent line:

$$f'(a) = \frac{\text{rise}}{\text{run}} = \frac{y - f(a)}{x - a}$$

The tangent line to f(x) at a has slope f'(a) and passes through the point (a, f(a)).

Slope of the tangent line:

$$f'(a) = \frac{\text{rise}}{\text{run}} = \frac{y - f(a)}{x - a}$$

The tangent line to f(x) at a has slope f'(a) and passes through the point (a, f(a)).

Slope of the tangent line:

$$f'(a) = \frac{\text{rise}}{\text{run}} = \frac{y - f(a)}{x - a}$$

Rearranging: y - f(a) = f'(a)(x - a) (equation of tangent line)

The tangent line to the function f(x) at point a is:

$$(y - f(a)) = f'(a)(x - a)$$

The tangent line to the function f(x) at point a is:

$$(y - f(a)) = f'(a)(x - a)$$

Point-Slope Formula

In general, a line with slope m passing through point (x_1, y_1) has the equation:

$$(y-y_1)=m(x-x_1)$$

The tangent line to the function f(x) at point a is:

$$(y - f(a)) = f'(a)(x - a)$$

Point-Slope Formula

In general, a line with slope m passing through point (x_1, y_1) has the equation:

$$(y-y_1)=m(x-x_1)$$

Find the equation of the tangent line to the curve $f(x) = \sqrt{x}$ at x = 9. (Recall $\frac{d}{dx} \left[\sqrt{x} \right] = \frac{1}{2\sqrt{x}}$).

23/27

The tangent line to the function f(x) at point a is:

$$(y - f(a)) = f'(a)(x - a)$$

Point-Slope Formula

In general, a line with slope m passing through point (x_1, y_1) has the equation:

$$(y-y_1)=m(x-x_1)$$

Find the equation of the tangent line to the curve $f(x) = \sqrt{x}$ at x = 9. (Recall $\frac{d}{dx} \left[\sqrt{x} \right] = \frac{1}{2\sqrt{x}}$).

$$a = 9$$
, $f(a) = 3$, $f'(a) = \frac{1}{2\sqrt{9}} = \frac{1}{6}$
 $y - 3 = \frac{1}{6}(x - 9)$

Memorize

The tangent line to the function f(x) at point a is:

$$(y - f(a)) = f'(a)(x - a)$$



Let $s(t) = 3 - 0.8t^2$. Then s'(t) = -1.6t. Find the

equation for the tangent line to the function s(t) when t = 1.

Let $s(t) = 3 - 0.8t^2$. Then s'(t) = -1.6t. Find the

equation for the tangent line to the function s(t) when t = 1.

$$a = 1$$
, $s(a) = 2.2$, $s'(a) = -1.6$
 $y - 2.2 = -1.6(x - 1)$

Included Work



'Brain' by Eucalyp is licensed under CC BY 3.0 (accessed 8 June 2021), 26, 27

Natasha Deshpande, Anoosha Kumar, Rohini Ramaswami. (2014). The Effect of National Healthcare Expenditure on Life Expectancy, page 12. College of Liberal Arts -Ivan Allen College (IAC), School of Economics: Econometric Analysis Undergraduate Research Papers. https://smartech.gatech.edu/handle/1853/51648 (accessed July 2021), 13