


TABLE OF CONTENTS



RADIOACTIVE DECAY

The number of atoms in a sample that decay in a given time interval is proportional to the number of atoms in the sample.

Differential Equation

Let $Q = Q(t)$ be the amount of a radioactive substance at time t . Then for some positive constant k :

$$\frac{dQ}{dt} = -kQ$$

RADIOACTIVE DECAY

The number of atoms in a sample that decay in a given time interval is proportional to the number of atoms in the sample.

Differential Equation

Let $Q = Q(t)$ be the amount of a radioactive substance at time t . Then for some positive constant k :

$$\frac{dQ}{dt} = -kQ$$

Solution – Theorem 3.3.2

Let $Q(t) = Ce^{-kt}$, where k and C are constants. Then:

$$\frac{dQ}{dt}(t) = C \cdot e^{-kt} \cdot (-k) = -kCe^{-kt} = -kQ(t)$$

RADIOACTIVE DECAY

Quantity of a Radioactive Isotope

$$Q(t) = Ce^{-kt}$$

$Q(t)$: quantity at time t

Every 30 seconds, the size of the sample halves. What are C and k ?



A sample of radioactive matter is stored in a lab in 2000. In the year 2002, it is tested and found to contain 10 units of a particular radioactive isotope. In the year 2005, it is tested and found to contain only 2 units of that same isotope. How many units of the isotope were present in the year 2000?



$$Q'(t) = kQ(t)$$

The number of atoms in a sample that decay in a given time interval is proportional to the number of atoms in the sample.

$$Q'(t) = kQ(t)$$

The number of atoms in a sample that decay in a given time interval is proportional to the number of atoms in the sample.

The rate of growth of a population in a given time interval is proportional to the number of individuals in the population, when the population has ample resources.

$$Q'(t) = kQ(t)$$

The number of atoms in a sample that decay in a given time interval is proportional to the number of atoms in the sample.

The rate of growth of a population in a given time interval is proportional to the number of individuals in the population, when the population has ample resources.

The amount of interest a bank account accrues in a given time interval is proportional to the balance in that bank account.

Exponential Growth – Theorem 3.3.2

Let $Q = Q(t)$ satisfy:

$$\frac{dQ}{dt} = kQ$$

for some constant k . Then for some constant $C = Q(0)$,

$$Q(t) = Ce^{kt}$$

$$\frac{dy}{dt} + 3y = 0 \quad \text{and} \quad y(1) = 2.$$

What is $y(t)$?

POPULATION GROWTH

Suppose a petri dish starts with a culture of 100 bacteria cells and a limited amount of food and space. The population of the culture at different times is given in the table below. At approximately what time did the culture start to show signs of limited resources?

time	population
0	100
1	1000
3	100000
5	1000000

Newton's Law of Cooling – Equation 3.3.7

The rate of change of temperature of an object is proportional to the difference in temperature between that object and its surroundings.

Newton's Law of Cooling – Equation 3.3.7

The rate of change of temperature of an object is proportional to the difference in temperature between that object and its surroundings.

$$\frac{dT}{dt}(t) = K[T(t) - A]$$

where $T(t)$ is the temperature of the object at time t , A is the (constant) ambient temperature of the surroundings, and K is some constant depending on the object.

$$\frac{dT}{dt}(t) = K[T(t) - A]$$

$T(t)$ is the temperature of the object, A is the ambient temperature, K is some constant.

What is true of K ?

- A. $K \geq 0$
B. $K \leq 0$
C. $K = 0$
D. K could be positive, negative, or zero, depending on the object
E. I don't know

$$\frac{dT}{dt}(t) = K[T(t) - A]$$

$T(t)$ is the temperature of the object, A is the ambient temperature, K is some constant.

What is true of K ?

- A. $K \geq 0$
- B. $K \leq 0$
- C. $K = 0$
- D. K could be positive, negative, or zero, depending on the object
- E. I don't know

Newton's Law of Cooling – Equation 3.3.7

$$\frac{dT}{dt}(t) = K[T(t) - A]$$

$T(t)$ is the temperature of the object, A is the ambient temperature, and K is some constant.

$$T(t) = [T(0) - A]e^{Kt} + A$$

is the only function satisfying Newton's Law of Cooling

Newton's Law of Cooling – Equation 3.3.7

$$\frac{dT}{dt}(t) = K[T(t) - A]$$

$T(t)$ is the temperature of the object, A is the ambient temperature, and K is some constant.

$$T(t) = [T(0) - A]e^{Kt} + A$$

is the only function satisfying Newton's Law of Cooling

If $T(10) < A$, then:

- A. $K > 0$
- B. $T(0) > 0$
- C. $T(0) > A$
- D. $T(0) < A$

Newton's Law of Cooling – Equation 3.3.7

$$\frac{dT}{dt}(t) = K[T(t) - A]$$

$T(t)$ is the temperature of the object, A is the ambient temperature, and K is some constant.

$$T(t) = [T(0) - A]e^{Kt} + A$$

is the only function satisfying Newton's Law of Cooling

If $T(10) < A$, then:

- A. $K > 0$
- B. $T(0) > 0$
- C. $T(0) > A$
- D. $T(0) < A$

Evaluate $\lim_{t \rightarrow \infty} T(t)$.

- A. A
- B. 0
- C. ∞
- D. $T(0)$

Newton's Law of Cooling – Equation 3.3.7

$$\frac{dT}{dt}(t) = K[T(t) - A]$$

$T(t)$ is the temperature of the object, A is the ambient temperature, and K is some constant.

$$T(t) = [T(0) - A]e^{Kt} + A$$

is the only function satisfying Newton's Law of Cooling

If $T(10) < A$, then:

- A. $K > 0$
- B. $T(0) > 0$
- C. $T(0) > A$
- D. $T(0) < A$

Evaluate $\lim_{t \rightarrow \infty} T(t)$.

- A. A
- B. 0
- C. ∞
- D. $T(0)$



What assumptions are we making that might not square with the real world?

Newton's Law of Cooling – Equation 3.3.7

$$\frac{dT}{dt} = K[T(t) - A]$$

$T(t)$ is the temperature of the object, A is the ambient temperature, and K is some constant.

Temperature of a Cooling Body – Corollary 3.3.8

$$T(t) = [T(0) - A]e^{Kt} + A$$

A farrier forms a horseshoe heated to 400°C , then dunks it in a river at room-temperature (25°C). The water boils for 30 seconds. The horseshoe is safe for the horse when it's 40°C . When can the farrier put on the horseshoe?



A farrier forms a horseshoe heated to 400°C , then dunks it in a river at room-temperature (25°C). The water boils for 30 seconds. The horseshoe is safe for the horse when it's 40°C . When can the farrier put on the horseshoe?

$$T(t) = [T(0) - A]e^{Kt} + A$$



We know: $T(0) = 400$, $T(30) = 100$, and $A = 25$. We want to find K .

$$100 = T(30) = [T(0) - A]e^{30K} + A = 375e^{30K} + 25$$

$$\Rightarrow 75 = 375e^{30K} \Rightarrow \frac{1}{5} = e^{30K} \Rightarrow K = \frac{-\log 5}{30}$$

Now, we set $T(t) = 40$ and solve for t :

$$40 = T(t) = 375e^{\frac{-\log 5}{30}t} + 25$$

$$15 = 375e^{\frac{-\log 5}{30}t} = 375 \cdot 5^{-t/30}$$

$$\frac{1}{25} = 5^{-t/30}$$

$$25 = 5^{t/30}$$

$$2 = t/30$$

So the farrier can put the shoe on after 60 seconds in the water.



A glass of just-boiled tea is put on a porch outside. After ten minutes, the tea is 40° , and after 20 minutes, the tea is 25° . What is the temperature outside?



$$T(0) = 100, \text{ so}$$

$$T(10) = [100 - A]e^{10K} + A = 100e^{10K} + A(1 - e^{10K}) = 40$$

$$T(20) = [100 - A]e^{20K} + A = 100e^{20K} + A(1 - e^{20K}) = 25$$

$$\text{Solving both for } A, \text{ we get } A = \frac{40 - 100e^{10K}}{1 - e^{10K}} = \frac{25 - 100e^{20K}}{1 - e^{20K}}$$

Although this looks complicated, if we set $x = e^{10k}$, it simplifies to something we can easily solve.



$$A = \frac{40 - 100e^{10K}}{1 - e^{10K}} = \frac{25 - 100e^{20K}}{1 - e^{20K}}$$

$$A = \frac{40 - 100x}{1 - x} = \frac{25 - 100x^2}{1 - x^2}$$

$$(40 - 100x)(1 - x^2) = (25 - 100x^2)(1 - x)$$

$$(40 - 100x)(1 + x)(1 - x) = (25 - 100x^2)(1 - x)$$

$$(40 - 100x)(1 + x) = 25 - 100x^2$$

$$40 - 60x - 100x^2 = 25 - 100x^2$$

$$40 - 60x = 25$$

$$x = \frac{1}{4}$$

$$A = \frac{40 - 100x}{1 - x} = \frac{40 - \frac{100}{4}}{1 - \frac{1}{4}} = 20$$

It is 20 degrees outside.



In 1963, the US Fish and Wildlife Service recorded a bald eagle population of 487 breeding pairs. In 1993, that number was 4015. How many breeding pairs would you expect there were in 2006? What about 2015?

○○○○○○○○○○○○○○○○●○○○○○○○○


○○○○○○○○○○○○

○○○○○○○

○○○○○○○○○○○

○○○○○○○○○○○○○○○○

○○○○○○○



Since we don't have a better model, let's assume the population P of nesting pairs follows:

$$P(t) = P(0)e^{Kt}$$

for some constant K .

To fit the data we have, let $t = 0$ represent 1963, so $P(0) = 487$. Then

$$4015 = P(30) = 487e^{30K}$$

$$\text{so } e^K = \left(\frac{4015}{487}\right)^{1/30}.$$

Now we use this to predict $P(43)$ (since 2006 is 43 years after 1963) and $P(52)$ (since 2015 is 52 years after 1963).

$$P(43) = 487(e^K)^{43} = 487 \left(\frac{4015}{487}\right)^{43/30} \approx 10016$$

So we guess in 2016 there were about 10,016 breeding pairs in the lower 48.

$$P(52) = 487(e^K)^{52} = 487 \left(\frac{4015}{487}\right)^{52/30} \approx 18860$$

link: Wood Bison Restoration in Alaska, Alaska Department of Fish and Game

Excerpt:

Based on experience with reintroduced populations elsewhere, wood bison would be expected to increase at a rate of 15%-25% annually after becoming established.... With an average annual growth rate of 20%, an initial precalving population of 50 bison would increase to 500 in approximately 13 years.

NOW
YOU



Are they using our same model?

link: Wood Bison Restoration in Alaska, Alaska Department of Fish and Game

Excerpt:

Based on experience with reintroduced populations elsewhere, wood bison would be expected to increase at a rate of 15%-25% annually after becoming established.... With an average annual growth rate of 20%, an initial precalving population of 50 bison would increase to 500 in approximately 13 years.

NOW
YOU



Are they using our same model?

Our model gives the same result.

COMPOUND INTEREST

Suppose you invest \$10,000 in an account that accrues interest each month. After one month, your balance (with interest) is \$10,100. How much money will be in your account after a year?

COMPOUND INTEREST

Suppose you invest \$10,000 in an account that accrues interest each month. After one month, your balance (with interest) is \$10,100. How much money will be in your account after a year?

Compound interest is calculated according to the formula Pe^{rt} , where r is the interest rate and t is time.



COMPOUND INTEREST

Suppose you invest \$10,000 in an account that accrues interest each month. After one month, your balance (with interest) is \$10,100. How much money will be in your account after a year?

Compound interest is calculated according to the formula Pe^{rt} , where r is the interest rate and t is time.

Measuring time in months,

$$10000e^{r \cdot 1} = 10100$$

$$e^r = \frac{10100}{10000} = 1.01$$

$$10000e^{12r} = 10000 \cdot (e^r)^{12} = 10000 \cdot 1.01^{12} \approx 11268.25$$



CARRYING CAPACITY

For a population of size P with unrestricted access to resources, let β be the average number of offspring each breeding pair produces per generation, where a generation has length t_g . Then $b = \frac{\beta-2}{2t_g}$ is the net birthrate (births minus deaths) per member per unit time. This yields $\frac{dP}{dt}(t) = bP(t)$, hence:



CARRYING CAPACITY

For a population of size P with unrestricted access to resources, let β be the average number of offspring each breeding pair produces per generation, where a generation has length t_g . Then $b = \frac{\beta-2}{2t_g}$ is the net birthrate (births minus deaths) per member per unit time. This yields $\frac{dP}{dt}(t) = bP(t)$, hence:

$$P(t) = P(0)e^{bt}$$



CARRYING CAPACITY

For a population of size P with unrestricted access to resources, let β be the average number of offspring each breeding pair produces per generation, where a generation has length t_g . Then $b = \frac{\beta-2}{2t_g}$ is the net birthrate (births minus deaths) per member per unit time. This yields $\frac{dP}{dt}(t) = bP(t)$, hence:

$$P(t) = P(0)e^{bt}$$


But as resources grow scarce, b might change.

CARRYING CAPACITY

b is the net birthrate (births minus deaths) per member per unit time.

If K is the carrying capacity of an ecosystem, we can model

$$b = b_0 \left(1 - \frac{P}{K}\right).$$




CARRYING CAPACITY

b is the net birthrate (births minus deaths) per member per unit time.

If K is the carrying capacity of an ecosystem, we can model

$$b = b_0 \left(1 - \frac{P}{K}\right).$$



NOW
YOU



Describe to your neighbour what the following mean in

terms of the model:

- ▶ $b > 0, b = 0, b < 0$
- ▶ $P = 0, P > 0, P < 0$



CARRYING CAPACITY

Then:

$$\frac{dP}{dt}(t) = b_0 \underbrace{\left(1 - \frac{P(t)}{K}\right)}_{\text{per capita birthrate}} P(t)$$

CARRYING CAPACITY

Then:

$$\frac{dP}{dt}(t) = b_0 \underbrace{\left(1 - \frac{P(t)}{K}\right)}_{\text{per capita birthrate}} P(t)$$

This is an example of a differential equation that we don't have the tools to solve. (If you take more calculus, though, you'll learn how!) It's also an example of a way you might tweak a model so its assumptions better fit what you observe.

RADIOCARBON DATING

Researchers at Charlie Lake in BC have found evidence¹ of habitation dating back to around 8500 BCE. For instance, a butchered bison bone was radiocarbon dated to about 10,500 years ago.

Suppose a comparable bone of a bison alive today contains $1\mu\text{g}$ of ^{14}C . If the half-life of ^{14}C is about 5730 years, roughly how much ^{14}C do you think the researchers found in the sample?

¹<http://pubs.aina.ucalgary.ca/arctic/Arctic49-3-265.pdf>

RADIOCARBON DATING

Researchers at Charlie Lake in BC have found evidence of habitation dating back to around 8500 BCE. For instance, a butchered bison bone was radiocarbon dated to about 10,500 years ago.

Suppose a comparable bone of a bison alive today contains $1\mu\text{g}$ of ^{14}C . If the half-life of ^{14}C is about 5730 years, roughly how much ^{14}C do you think the researchers found in the sample?

- A. About $\frac{1}{10,500} \mu\text{g}$
- B. About $\frac{1}{4} \mu\text{g}$
- C. About $\frac{1}{2} \mu\text{g}$

- D. About $1 \mu\text{g}$
- E. I'm not sure how to estimate this



RADIOCARBON DATING

First, an estimate; 10500 is not so far off from $2(5730)$, i.e. two half-lives, so we might guess that there is roughly a $(\frac{1}{2})^2 = \frac{1}{4}$ of a microgram left.

We know $Q(t) = Ce^{-kt} = e^{-kt} \mu\text{g}$. We want to find $Q(10500)$, so we need to solve for k . Since we know the half-life: to do this, solve

$$\frac{1}{2} = e^{-k \cdot 5730} \quad \text{to get} \quad k = \frac{\log 2}{5730}$$

Now:

$$Q(10500) = e^{-\frac{\log 2}{5730} \cdot 10500} = 2^{-\frac{10500}{5730}} \approx 0.28 \mu\text{g}$$



Suppose a body is discovered at 3:45 pm, in a room held at 20° , and the body's temperature is 27° , not the normal 37° . At 5:45 pm, the temperature of the body has dropped to 25.3° . When did the inhabitant of the body die?



Set our time so that $t = 0$ is 3:45pm and $t = 2$ is 5:45pm. Then $T(0) = 27$, $T(2) = 25.4$, and $A = 20$. Now:

$$T(t) = [27 - 20]e^{Kt} + 20 = 7e^{Kt} + 20$$

Using what we know about 5:45pm:

$$7e^{2K} + 20 = T(2) = 25.3$$

so

$$7e^{2K} = 5.3 \implies e^{2K} = \frac{5.3}{7} \implies e^K = \left(\frac{5.3}{7}\right)^{1/2}$$

Now:

$$T(t) = 7e^{Kt} + 20 = 7\left(\frac{5.3}{7}\right)^{t/2} + 20$$

So we set $T(t) = 37$ and solve for t .



$$7 \left(\frac{5.3}{7} \right)^{t/2} + 20 = 37$$

$$7 \left(\frac{5.3}{7} \right)^{t/2} = 17$$

$$\left(\frac{5.3}{7} \right)^{t/2} = \frac{17}{7}$$


$$\frac{t}{2} = \frac{\log(17/7)}{\log(5.3/7)}$$

$$t = 2 \frac{\log(17/7)}{\log(5.3/7)} \approx -6.4$$


So the person died about 6.4 hours before 3:45pm. Now 0.4 hours is 24 minutes. So 6 hours and 24 minutes before 3:45 pm is 6 hours before 3:21pm, which is 9:21 am.




TABLE OF CONTENTS




APPROXIMATING A FUNCTION




APPROXIMATING A FUNCTION



APPROXIMATING A FUNCTION



APPROXIMATING A FUNCTION




Constant Approximation – Equation 3.4.1

We can approximate $f(x)$ near a point a by

$$f(x) \approx f(a)$$

APPROXIMATING A FUNCTION



Constant Approximation – Equation 3.4.1


We can approximate $f(x)$ near a point a by

$$f(x) \approx f(a)$$

Google: $\sin(0.2) \approx 0.198669\dots$

Constant approx: $\sin(0.2) \approx 0$

APPROXIMATING A FUNCTION



Constant Approximation – Equation 3.4.1


We can approximate $f(x)$ near a point a by

$$f(x) \approx f(a)$$


Google: $\sin(0.2) \approx 0.198669\dots$

Constant approx: $\sin(0.2) \approx 0$


APPROXIMATING A FUNCTION



APPROXIMATING A FUNCTION



APPROXIMATING A FUNCTION



Linear Approximation (Linearization) – Equation 3.4.3

We can approximate $f(x)$ near a point a by the tangent line to $f(x)$ at a , namely


$$f(x) \approx L(x) = f(a) + f'(a)(x - a)$$

Google:

$$\sin(0.2) \approx 0.198669...$$

Linear approx:

$$\sin(0.2) \approx 0 + 1(0.2 - 0) = 0.2$$

$$f(x) \approx L(x) = f(a) + f'(a)(x - a)$$


○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○

○ ○ ○ ● ○ ○ ○ ○ ○ ○ ○ ○ ○ ○

○○○○○○

○○○○○○○○○○

○○○○○○○○○○○○○○○○○○

○ ○ ○ ○ ○ ○ ○ ○ ○ ○

To find a linear approximation of $f(x)$ at a particular point x , pick a point a **near to x** , such that $f(a)$ and $f'(a)$ are **easy to calculate**.

$$f(x) \approx L(x) = f(a) + f'(a)(x - a)$$

Let $f(x) = \sqrt{x}$. Approximate $f(8.9)$.

○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○

☐ ☐ ☒ ☐ ☐ ☐ ☐ ☐ ☐

○○○○○

○○○○○○○○○○○○○○

○○○○○○○○○○○○○○○○○○○○

○○○○○○○○

To find a linear approximation of $f(x)$ at a particular point x , pick a point a **near to x** , such that $f(a)$ and $f'(a)$ are **easy to calculate**.

$$f(x) \approx L(x) = f(a) + f'(a)(x - a)$$

Let $f(x) = \sqrt{x}$. Approximate $f(8.9)$.

First we note that $8.9 \approx 9$ and we can easily calculate $f(9) = 3$.

○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○

☐ ☐ ☒ ☐ ☐ ☐ ☐ ☐ ☐

○○○○○

○○○○○○○○○○○○

○○○○○○○○○○○○○○○○

○○○○○○○○○○

To find a linear approximation of $f(x)$ at a particular point x , pick a point a **near to x** , such that $f(a)$ and $f'(a)$ are **easy to calculate**.

$$f(x) \approx L(x) = f(a) + f'(a)(x - a)$$

Let $f(x) = \sqrt{x}$. Approximate $f(8.9)$.

First we note that $8.9 \approx 9$ and we can easily calculate $f(9) = 3$.

Constant approximation: $8.9 \approx 9$, so $f(8.9) \approx f(9) = \boxed{3}$



☐ ☐ ☒ ☐ ☐ ☐ ☐ ☐ ☐

○○○○○○

○○○○○○○○○○

○○○○○○○○○○○○○○○○○○○○

○○○○○○○○

To find a linear approximation of $f(x)$ at a particular point x , pick a point a **near to x** , such that $f(a)$ and $f'(a)$ are **easy to calculate**.

$$f(x) \approx L(x) = f(a) + f'(a)(x - a)$$

Let $f(x) = \sqrt{x}$. Approximate $f(8.9)$.

First we note that $8.9 \approx 9$ and we can easily calculate $f(9) = 3$.

Constant approximation: $8.9 \approx 9$, so $f(8.9) \approx f(9) = \boxed{3}$

Linear approximation: Using $a = 9$,

$$f'(a) = \frac{1}{2\sqrt{a}} = \frac{1}{2\sqrt{9}} = \frac{1}{6}$$

$$f(8.9) \approx f(9) + f'(9)(8.9 - 9) = 3 + \frac{1}{6}(-.1)$$

$$f(8.9) \approx 3 - \frac{1}{60} = 2.98\overline{33}$$

○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○

☐ ☐ ☒ ☐ ☐ ☐ ☐ ☐ ☐

○○○○○

○○○○○○○○○○○○

○○○○○○○○○○○○○○○○○○○○

○○○○○○○○

To find a linear approximation of $f(x)$ at a particular point x , pick a point a **near to x** , such that $f(a)$ and $f'(a)$ are **easy to calculate**.

$$f(x) \approx L(x) = f(a) + f'(a)(x - a)$$

Let $f(x) = \sqrt{x}$. Approximate $f(8.9)$.

First we note that $8.9 \approx 9$ and we can easily calculate $f(9) = 3$.

Constant approximation: $8.9 \approx 9$, so $f(8.9) \approx f(9) = \boxed{3}$

Linear approximation: Using $a = 9$,

$$f'(a) = \frac{1}{2\sqrt{a}} = \frac{1}{2\sqrt{9}} = \frac{1}{6}$$

$$f(8.9) \approx f(9) + f'(9)(8.9 - 9) = 3 + \frac{1}{6}(-.1)$$

$$f(8.9) \approx 3 - \frac{1}{60} = 2.98\overline{33}$$

Google: $\sqrt{8.9} = 2.98328677804\dots$

81 / 197


Accurate

Accurate

Possible to calculate: add, subtract, multiply, divide. Use integers or known constants

LINEAR APPROXIMATION

Approximate $\sin(3)$ using a linear approximation. You may leave your answer in terms of π .



Let $f(x) = \sin x$ and $a = \pi$. Then

$$f(3) \approx f(\pi) + f'(\pi)(3 - \pi) = \sin(\pi) + \cos(\pi)(3 - \pi) = \boxed{\pi - 3} \approx 0.14159$$

Google: $\sin(3) = 0.14112000806\dots$

Approximate $e^{1/10}$ using a linear approximation.

LINEAR APPROXIMATION

Approximate $e^{1/10}$ using a linear approximation.

If $f(x) = e^x$ and $a = 0$:

$$f'(x) = e^x$$

$$f(1/10) \approx f(0) + f'(0)(1/10 - 0) = e^0 + e^0(1/10 - 0) = 1 + 1/10 = 1.1$$

Google: $e^{1/10} = 1.10517091808\dots$

LINEAR APPROXIMATION

Approximate $e^{1/10}$ using a linear approximation.

If $f(x) = e^x$ and $a = 0$:

$$f'(x) = e^x$$

$$f(1/10) \approx f(0) + f'(0)(1/10 - 0) = e^0 + e^0(1/10 - 0) = 1 + 1/10 = 1.1$$


If $g(x) = x^{1/10}$:

Google: $e^{1/10} = 1.10517091808\dots$



LINEAR APPROXIMATION WRAP-UP


Let $L(x) = f(a) + f'(a)(x - a)$, so $L(x)$ is the linear approximation (linearization) of $f(x)$ at a .



LINEAR APPROXIMATION WRAP-UP

Let $L(x) = f(a) + f'(a)(x - a)$, so $L(x)$ is the linear approximation (linearization) of $f(x)$ at a .

What is $L(a)$?




LINEAR APPROXIMATION WRAP-UP

Let $L(x) = f(a) + f'(a)(x - a)$, so $L(x)$ is the linear approximation (linearization) of $f(x)$ at a .

What is $L(a)$?

What is $L'(a)$?




LINEAR APPROXIMATION WRAP-UP

Let $L(x) = f(a) + f'(a)(x - a)$, so $L(x)$ is the linear approximation (linearization) of $f(x)$ at a .

What is $L(a)$?

What is $L'(a)$?

What is $L''(a)$? (Recall $L''(x)$ is the derivative of $L'(x)$.)



LINEAR APPROXIMATION WRAP-UP

Let $L(x) = f(a) + f'(a)(x - a)$, so $L(x)$ is the linear approximation (linearization) of $f(x)$ at a .


What is $L(a)$?

$$L(a) = f(a)$$

What is $L'(a)$?

$$L'(a) = f'(a)$$

What is $L''(a)$? (Recall $L''(x)$ is the derivative of $L'(x)$.)



LINEAR APPROXIMATION WRAP-UP

Let $L(x) = f(a) + f'(a)(x - a)$, so $L(x)$ is the linear approximation (linearization) of $f(x)$ at a .

What is $L(a)$?


$$L(a) = f(a)$$

What is $L'(a)$?

$$L'(a) = f'(a)$$


What is $L''(a)$? (Recall $L''(x)$ is the derivative of $L'(x)$.)

$$L''(a) = 0$$




QUADRATIC APPROXIMATION

Imagine we approximate $f(x)$ at $x = a$ with a **parabola**, $P(x)$.



QUADRATIC APPROXIMATION

Imagine we approximate $f(x)$ at $x = a$ with a **parabola**, $P(x)$.



$P(x) = A + Bx + Cx^2$	$P(a) = A + Ba + Ca^2$	$f(a)$
$P'(x) = B + 2Cx$	$P'(a) = B + 2Ca$	$f'(a)$
$P''(x) = 2C$	$P''(a) = 2C$	$f''(a)$

108/197

Constant:

$$f(x) \approx f(a)$$

Linear:

$$f(x) \approx f(a) + f'(a)(x - a)$$

Quadratic:

$$f(x) \approx f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2$$

QUADRATIC APPROXIMATION

$$P(x) = f(a) + f'(a)(x - a) + \frac{1}{2}f''(a)(x - a)^2$$

Approximate $\sqrt[3]{28}$ using a quadratic approximation.

You may leave your answer unsimplified, as long as it is an expression you could figure out from integers using only plus, minus, times, and divide.

$$(2.01)^6$$

Determine what $f(x)$ and a should be so that you can approximate the following using a quadratic approximation.

$$\log(.9) \quad f(x) = \log(x), a = 1$$

$$e^{-1/30} \quad f(x) = e^x, a = 0$$

$$\sqrt[5]{30} \quad f(x) = \sqrt[5]{x}, a = 32 = 2^5$$

$$(2.01)^6 \quad f(x) = x^6, a = 2$$

It is possible to compute the last one without an approximation, but an approximation might save time while being sufficiently accurate for your purposes.



	Constant	Linear	Quadratic	degree n
match $f(a)$	✓	✓	✓	✓
match $f'(a)$	✗	✓	✓	✓
match $f''(a)$	✗	✗	✓	✓
...				
match $f^{(n)}(a)$	✗	✗	✗	✓
match $f^{(n+1)}(a)$	✗	✗	✗	✗

$$f(x) \approx f(a)$$

Linear:

$$f(x) \approx f(a) + f'(a)(x - a)$$

Quadratic:

$$f(x) \approx f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2$$

Degree- n :

$$f(x) \approx f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2 + \dots?$$

BRIEF DETOUR: SIGMA (SUMMATION) NOTATION

$$\sum_{i=a}^b f(i)$$

BRIEF DETOUR: SIGMA (SUMMATION) NOTATION

$$\sum_{i=a}^b f(i)$$

- a, b (integers) “bounds”

BRIEF DETOUR: SIGMA (SUMMATION) NOTATION

$$\sum_{i=a}^b f(i)$$

- ▶ a, b (integers) “bounds”
- ▶ i “index”: runs over integers from a to b

BRIEF DETOUR: SIGMA (SUMMATION) NOTATION

$$\sum_{i=a}^b f(i)$$

- ▶ a, b (integers) “bounds”
- ▶ i “index”: runs over integers from a to b
- ▶ $f(i)$ “summand”: compute for every i , add

$$\sum_{i=2}^4 (2i + 5)$$

SIGMA NOTATION

$$\sum_{i=2}^4 (2i + 5)$$

$$\begin{aligned}\sum_{i=2}^4 (2i+5) &= \underbrace{(2 \cdot 2 + 5)}_{i=2} + \underbrace{(2 \cdot 3 + 5)}_{i=3} + \underbrace{(2 \cdot 4 + 5)}_{i=4} \\ &= 9 + 11 + 13 = 33\end{aligned}$$

SIGMA NOTATION

$$\sum_{i=1}^4 (i + (i-1)^2)$$

SIGMA NOTATION

$$\sum_{i=1}^4 (i + (i-1)^2)$$

$$= \underbrace{(1 + 0^2)}_{i=1} + \underbrace{(2 + 1^2)}_{i=2} + \underbrace{(3 + 2^2)}_{i=3} + \underbrace{(4 + 3^2)}_{i=4}$$
$$= 1 + 3 + 7 + 13 = 24$$

1. $3 + 4 + 5 + 6 + 7$
2. $8 + 8 + 8 + 8 + 8$
3. $1 + (-2) + 4 + (-8) + 16$

$$T_n(a) = \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k$$

$$\begin{aligned} T_n(a) &= \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k \\ &= \underbrace{f(a)}_{k=0} + \underbrace{f'(a)(x-a)}_{k=1} + \underbrace{\frac{1}{2!}f''(a)(x-a)^2}_{k=2} + \\ &\quad \underbrace{\frac{1}{3!}f'''(a)(x-a)^3}_{k=3} + \underbrace{\frac{1}{4!}f^{(4)}(a)(x-a)^4}_{k=4} + \\ &\quad \cdots + \underbrace{\frac{1}{n!}f^{(n)}(a)(x-a)^n}_{k=n} \end{aligned}$$




SMALL DEGREE TAYLOR POLYNOMIALS

$$\begin{aligned} T_0(a) &= \sum_{k=0}^0 \frac{f^{(k)}(a)}{k!} (x-a)^k \\ &= f(a) \end{aligned}$$

The 0th degree Taylor polynomial is the **constant** approximation.

Find the 7th degree Maclaurin² polynomial for e^x .

Q



$$T_n(a) = f(a) + f'(a)(x-a) + \frac{1}{2!}f''(a)(x-a)^2 + \cdots + \frac{1}{n!}f^{(n)}(a)(x-a)^n$$

Find the 7th degree Taylor polynomial for $f(x) = \log x$, centered at $a = 1$.

$$f(x) = \log x$$

$$f(1) = 0$$

$$f^{(4)}(x) = -3!x^{-4}$$

$$f^{(4)}(1) = -3!$$

$$f'(x) = x^{-1}$$

$$f'(1) = 1$$

$$f^{(5)}(x) = 4!x^{-5}$$

$$f^{(5)}(1) = 4!$$

$$f''(x) = -x^{-2}$$

$$f''(1) = -1$$

$$f^{(6)}(x) = -5!x^{-6}$$

$$f^{(6)}(1) = -5!$$

$$f'''(x) = 2x^{-3}$$

$$f'''(1) = 2$$

$$f^{(7)}(x) = 6!x^{-7}$$

$$f^{(7)}(1) = 6!$$

$$\begin{aligned} T_8(x) &= f(1) + f'(1)(x-1) + \frac{1}{2}f''(1)(x-1)^2 + \cdots + \frac{1}{7!}f^{(7)}(1)(x-1)^7 \\ &= 0 + (1)(x-1) + (-1)\frac{1}{2}(x-1)^2 + (2)\frac{1}{3!}(x-1)^3 - 3!\frac{1}{4!}(x-1)^4 \\ &\quad + 4!\frac{1}{5!}(x-1)^5 - 5!\frac{1}{6!}(x-1)^6 + 6!\frac{1}{7!}(x-1)^7 \\ &= (x-1) - \frac{(x-1)^2}{2} + \frac{(x-1)^3}{3} - \frac{(x-1)^4}{4} + \frac{(x-1)^5}{5} - \frac{(x-1)^6}{6} + \frac{(x-1)^7}{7!} \\ &= \sum_{k=1}^7 (-1)^{k+1} \frac{(x-1)^k}{k} \end{aligned}$$

Notation 3.4.18

Let x, y be variables related such that $y = f(x)$. Then we denote a small change in the variable x by Δx (read as “delta x ”). The corresponding small change in the variable y is denoted Δy (read as “delta y ”).

$$\Delta y = f(x + \Delta x) - f(x)$$

Thinking about change in this way can lead to convenient approximations.

○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○○

○○○○○○○○○○

○○○○○○

○○○○○○○○○○○○○○○○

○●○○○○○○○○○○○○○○○○○○

0 00000000

Let $y = f(x)$ be the amount of water needed to produce x apples in an orchard.

A farmer wants to know how much water is needed to increase their crop yield.



A farmer wants to know how much water is needed to increase their crop yield. Δx is shorthand for some change in the number of apples, and Δy is shorthand for some change in the amount of water.



Let $y = f(x)$ be the amount of water needed to produce x apples in an orchard.

A farmer wants to know how much water is needed to increase their crop yield. Δx is shorthand for some change in the number of apples, and Δy is shorthand for some change in the amount of water.

- Consider changing the number of apples grown from a to $a + \Delta x$



LINEAR APPROXIMATION OF Δy

- Using a linear approximation, setting $x = a + \Delta x$:

QUADRATIC APPROXIMATION OF Δy

If we wanted a more accurate approximation, we can use other Taylor polynomials. For example, let's try the quadratic approximation.

Your answer may consist of the sum, difference, product, and quotient of integers, roots of integers, and π .

$$f''\left(\frac{\pi}{3}\right) = \frac{\sqrt{3}}{(1/2)^3} = 8\sqrt{3}$$

$$f\left(\frac{13\pi}{36}\right) \approx f\left(\frac{\pi}{3}\right) = \sqrt{3}$$

$$\begin{aligned} f\left(\frac{13\pi}{36}\right) &\approx f\left(\frac{\pi}{3}\right) + f'\left(\frac{\pi}{3}\right)\left(\frac{13\pi}{36} - \frac{\pi}{3}\right) \\ &= \sqrt{3} + 4\left(\frac{\pi}{36}\right) \end{aligned}$$

$$\begin{aligned} f\left(\frac{13\pi}{36}\right) &\approx f\left(\frac{\pi}{3}\right) + f'\left(\frac{\pi}{3}\right)\left(\frac{13\pi}{36} - \frac{\pi}{3}\right) + \frac{1}{2}f''\left(\frac{\pi}{3}\right)\left(\frac{13\pi}{36} - \frac{\pi}{3}\right)^2 \\ &= \sqrt{3} + 4\left(\frac{\pi}{36}\right) + \frac{1}{2}(8\sqrt{3})\left(\frac{\pi}{36}\right)^2 \\ &= \sqrt{3} + \frac{\pi}{9} + \frac{4\sqrt{3}\pi^2}{6^4} \end{aligned}$$

type	approx	decimal
constant	$\sqrt{3}$	1.732...
linear	$\sqrt{3} + \frac{\pi}{9}$	2.081...
quadratic	$\sqrt{3} + \frac{\pi}{9} + \frac{4\sqrt{3}\pi^2}{6^4}$	2.134...
actual	—	2.145...

Definition 3.4.25

Let Q_0 be the exact value of a quantity and let $Q_0 + \Delta Q$ be the measured value. We call

$|\Delta Q|$

the **absolute error** of the measurement, and

$$100 \frac{|\Delta Q|}{Q_0}$$

the **percentage error** of the measurement.

Definition 3.4.25

Let Q_0 be the exact value of a quantity and let $Q_0 + \Delta Q$ be the measured value. We call

$|\Delta Q|$

the **absolute error** of the measurement, and

$$100 \frac{|\Delta Q|}{Q_0}$$


the **percentage error** of the measurement.

Suppose a bottle of water is labelled as having 500 mL of water, but in fact contains 502.

The absolute error of the labelling is 2, and the percent error is $100 \frac{2}{502} = 0.4$, or less than one-half of one percent.


Use a linear approximation.

ERROR: WHAT “CAUSES” ERROR IN AN ESTIMATION?




Constant approximation: We assume the function doesn't change, but in fact the function does change (its derivative is not always zero).

ERROR: WHAT “CAUSES” ERROR IN AN ESTIMATION?




Linear approximation: We assume the function changes at a constant rate, but in fact the function changes at different rates (its second derivative is not always zero).

ERROR: WHAT “CAUSES” ERROR IN AN ESTIMATION?



Quadratic approximation: We assume the function's derivative changes at a constant rate, but in fact the function's derivative changes at different rates (its third derivative is not always zero).


CONTROLLING THE “CAUSE” OF THE ERROR



Constant approximation: We assume the function doesn't change, but in fact the function does change (its derivative is not always zero).


BUT: suppose we know the max and min values of the function's slope.

CONTROLLING THE “CAUSE” OF THE ERROR




Linear approximation: We assume the function changes at a constant rate, but in fact the function changes at different rates (its first derivative is not always zero). **BUT:** suppose we know the max and min values of the function's second derivative.

CONTROLLING THE “CAUSE” OF THE ERROR



Linear approximation: We assume the function changes at a constant rate, but in fact the function changes at different rates (its first derivative is not always zero). **BUT:** suppose we know the max and min values of the function's second derivative.

CONTROLLING THE “CAUSE” OF THE ERROR



In general, if the “thing that causes the error” is big, then our error is big. We find the largest and smallest possible errors.

Third degree Maclaurin polynomial for $f(x) = e^x$:

Third degree Maclaurin polynomial for $f(x) = e^x$:

$$\begin{aligned} T_3(x) &= f(0) + f'(0)(x-0) + \frac{1}{2!}f''(0)(x-0)^2 + \frac{1}{3!}f'''(0)(x-0)^3 \\ &= e^0 + e^0x + \frac{1}{2!}e^0x^2 + \frac{1}{3!}e^0x^3 \\ &= 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} \end{aligned}$$

Third degree Maclaurin polynomial for $f(x) = e^x$:

$$\begin{aligned} T_3(x) &= f(0) + f'(0)(x-0) + \frac{1}{2!}f''(0)(x-0)^2 + \frac{1}{3!}f'''(0)(x-0)^3 \\ &= e^0 + e^0x + \frac{1}{2!}e^0x^2 + \frac{1}{3!}e^0x^3 \\ &= 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} \end{aligned}$$

Bound the error associated with using $T_3(x)$ to approximate $e^{1/10}$.

For some c in $(0, 0.1)$:

$$\underbrace{f(0.1) - T_3(0.1)}_{\text{error}} = \frac{1}{4!} f^{(4)}(c) (.1 - 0)^4$$

$$= \frac{1}{4!} (0.0001) e^c$$

For c in $(0, 0.1)$, $1 \leq e^c < e^1 < 3$, so

$$\begin{aligned} \frac{1}{4!}(0.0001) &\leq \underbrace{f(.1) - T_3(.1)}_{\text{error}} \\ &\leq \frac{1}{4!}(0.0001)3 \\ 4.2 \times 10^{-6} &\leq \underbrace{f(0.1) - T_3(0.1)}_{\text{error}} \leq 1.3 \times 10^{-5} \end{aligned}$$

For some c in $(\pi/2, 2)$:

$$\underbrace{f(2) - T_5(2)}_{\text{error}} = \frac{1}{6!} f^{(6)}(c) (2 - \pi/2)^6$$

Note $f^{(6)}(x)$ is going to be plus or minus sine or cosine, so $-1 \leq f^{(6)}(c) \leq 1$. Also, $0 < 2 - \pi/2 < 1$. Now:

$$\frac{-1}{6!} = \frac{1}{6!}(-1)(1)^6 \leq f(2) - T_5(2) \leq \frac{1}{6!}(1)(1)^6 = \frac{1}{6!}$$

And $\frac{1}{6!} \approx 0.0014$. Be very careful with positives and negatives here :)

We don't actually have to compute $T_5(x)$, but if you want to as an exercise, click [here](#) to see the result.

For some c in $(0, 1)$:

$$|f(1) - T_4(1)| = \left| \frac{1}{5!} f^{(5)}(c)(1-0)^5 \right| = \frac{1}{5!} e^c \leq \frac{1}{5!} e^1 < \frac{3}{5!} = 0.025$$



A reasonable question to ask is: which approximation will be good enough to keep our error within some fixed error tolerance?

WHICH DEGREE?

Suppose you want to approximate e^5 using a Maclaurin polynomial of $f(x) = e^x$. If the magnitude of your error must be less than 0.001, what degree Maclaurin polynomial should you use?

Included Work



'Water Drop' by [hunotika](#) is licensed under [CC BY 3.0](#) (accessed 21 July 2021), 144–147



'Brain' by [Eucalyp](#) is licensed under [CC BY 3.0](#) (accessed 8 June 2021), 46, 47, 54, 55, 141



'old tree' by [FayraLovers](#) is licensed under [CC BY 3.0](#) (accessed 21 July 2021), 144–147



'Tree' by [FayraLovers](#) is licensed under [CC BY 3.0](#) (accessed 21 July 2021), 145–147



'Tree' by [FayraLovers](#) is licensed under [CC BY 3.0](#) (accessed 21 July 2021), 146, 147



'Tree' by [FayraLovers](#) is licensed under [CC BY 3.0](#) (accessed 21 July 2021) / cropped from original, 147



'Chart of Bald Eagle Breeding Pairs in Lower 48 States'. Eagle nesting data, US Fish and Wildlife Service Midwest Region (accessed 19 October 2016), 44



'Notebook' by [Iconic](#) is licensed under [CC BY 3.0](#) (accessed 9 June 2021), 11, 20, 37, 178–180



U.S. WHO/NREVSS Collaborating Laboratories and ILNet. '[Stacked Column Chart WHO//NREVSS](#)' Centers for Disease Control and Prevention. No longer available from <http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html> (accessed 20 October 2015), 26

Alaska Department of Fish and Game, Division of Wildlife Conservation. (April 2007). Wood Bison Restoration in Alaska: A Review of Environmental and Regulatory Issues and Proposed Decisions for Project Implementation, p. 11. http://www.adfg.alaska.gov/static/species/speciesinfo/woodbison/pdfs/er_no_appendices.pdf (accessed 2015 or 2016), 46, 47

Driver et.al. Stratigraphy, Radiocarbon Dating, and Culture History of Charlie Lake Cave, British Columbia. *ARCTICVOL.* 49, no. 3 (September 1996) pp. 265 – 277. <http://pubs.aina.ucalgary.ca/arctic/Arctic49-3-265.pdf> (accessed 2015 or 2016), 58, 59

Public Domain by Man vyi via https://commons.wikimedia.org/wiki/File:West_Show_Jersey_2010_farrier_f.jpg, accessed October 2015, 37, 38