


TABLE OF CONTENTS

[▶ SKIP REVIEW OF INVERSE FUNCTIONS AND LOGARITHMS](#)

INVERTIBILITY GAME

- ▶ A function $y = f(x)$ is known to both players

INVERTIBILITY GAME

- ▶ A function $y = f(x)$ is known to both players
- ▶ **Player A** chooses a secret value x in the domain of $f(x)$

INVERTIBILITY GAME

- ▶ A function $y = f(x)$ is known to both players
- ▶ **Player A** chooses a secret value x in the domain of $f(x)$
- ▶ **Player A** tells **Player B** what $f(x)$ is

INVERTIBILITY GAME

- ▶ A function $y = f(x)$ is known to both players
- ▶ **Player A** chooses a secret value x in the domain of $f(x)$
- ▶ **Player A** tells **Player B** what $f(x)$ is
- ▶ **Player B** tries to guess **Player A**'s x -value.

INVERTIBILITY GAME

- ▶ A function $y = f(x)$ is known to both players
- ▶ **Player A** chooses a secret value x in the domain of $f(x)$
- ▶ **Player A** tells **Player B** what $f(x)$ is
- ▶ **Player B** tries to guess **Player A**'s x -value.

INVERTIBILITY GAME

- ▶ A function $y = f(x)$ is known to both players
- ▶ **Player A** chooses a secret value x in the domain of $f(x)$
- ▶ **Player A** tells **Player B** what $f(x)$ is
- ▶ **Player B** tries to guess **Player A**'s x -value.

Round 1: $f(x) = 2x$

INVERTIBILITY GAME

- ▶ A function $y = f(x)$ is known to both players
- ▶ **Player A** chooses a secret value x in the domain of $f(x)$
- ▶ **Player A** tells **Player B** what $f(x)$ is
- ▶ **Player B** tries to guess **Player A**'s x -value.

Round 1: $f(x) = 2x$

Round 2: $f(x) = \sqrt[3]{x}$

INVERTIBILITY GAME

- ▶ A function $y = f(x)$ is known to both players
- ▶ **Player A** chooses a secret value x in the domain of $f(x)$
- ▶ **Player A** tells **Player B** what $f(x)$ is
- ▶ **Player B** tries to guess **Player A**'s x -value.

Round 1: $f(x) = 2x$

Round 2: $f(x) = \sqrt[3]{x}$

Round 3: $f(x) = |x|$

INVERTIBILITY GAME

- ▶ A function $y = f(x)$ is known to both players
- ▶ **Player A** chooses a secret value x in the domain of $f(x)$
- ▶ **Player A** tells **Player B** what $f(x)$ is
- ▶ **Player B** tries to guess **Player A**'s x -value.


Round 1: $f(x) = 2x$

Round 2: $f(x) = \sqrt[3]{x}$


Round 3: $f(x) = |x|$

Round 4: $f(x) = \sin x$


FUNCTIONS ARE MAPS




FUNCTIONS ARE MAPS




FUNCTIONS ARE MAPS




FUNCTIONS ARE MAPS




FUNCTIONS ARE MAPS




FUNCTIONS ARE MAPS




FUNCTIONS ARE MAPS




FUNCTIONS ARE MAPS




FUNCTIONS ARE MAPS




FUNCTIONS ARE MAPS






A. invertible

B. not invertible




A. invertible

B. not invertible




A. invertible

B. not invertible




A. invertible

B. not invertible



A. invertible

B. not invertible



A. invertible

B. not invertible

RELATIONSHIP BETWEEN $f(x)$ AND $f^{-1}(x)$

Let f be an invertible function.

What is $f^{-1}(f(x))$?


- A. x
- B. 1
- C. 0
- D. not sure

RELATIONSHIP BETWEEN $f(x)$ AND $f^{-1}(x)$

Let f be an invertible function.

What is $f^{-1}(f(x))$?

- A. x
- B. 1
- C. 0
- D. not sure




RELATIONSHIP BETWEEN $f(x)$ AND $f^{-1}(x)$

Let f be an invertible function.

What is $f^{-1}(f(x))$?

- A. x
- B. 1
- C. 0
- D. not sure



Invertibility

In order for a function to be invertible

Invertibility

In order for a function to be invertible, different x values cannot map to the same y value.

Invertibility

In order for a function to be invertible, different x values cannot map to the same y value.

We call such a function **one-to-one**, or **injective**.

Invertibility

In order for a function to be invertible, different x values cannot map to the same y value.

We call such a function **one-to-one**, or **injective**.

Suppose $f(x) = \sqrt[3]{19 + x^3}$. What is $f^{-1}(3)$? (simplify your answer)

Invertibility

In order for a function to be invertible, different x values cannot map to the same y value.

We call such a function **one-to-one**, or **injective**.

Suppose $f(x) = \sqrt[3]{19 + x^3}$. What is $f^{-1}(3)$? (simplify your answer)

What is $f^{-1}(10)$? (do not simplify)

Invertibility

In order for a function to be invertible, different x values cannot map to the same y value.

We call such a function **one-to-one**, or **injective**.

Suppose $f(x) = \sqrt[3]{19 + x^3}$. What is $f^{-1}(3)$? (simplify your answer)

What is $f^{-1}(10)$? (do not simplify)

What is $f^{-1}(x)$?



Invertibility

In order for a function to be invertible, different x values cannot map to the same y value.

We call such a function **one-to-one**, or **injective**.


Suppose $f(x) = \sqrt[3]{19 + x^3}$. What is $f^{-1}(3)$? (simplify your answer)
 $f(2) = 3$, so $f^{-1}(3) = 2$


What is $f^{-1}(10)$? (do not simplify)
 $\sqrt[3]{19 + y^3} = 10$ tells us $f^{-1}(10) = \sqrt[3]{10^3 - 19}$

What is $f^{-1}(x)$?
 $\sqrt[3]{19 + y^3} = x$ tells us $f^{-1}(x) = \sqrt[3]{x^3 - 19}$


$$\text{Let } f(x) = x^2 - x.$$

1. Sketch a graph of $f(x)$, and choose a (large) domain over which it is invertible.
2. For the domain you chose, evaluate $f^{-1}(20)$.
3. For the domain you chose, evaluate $f^{-1}(x)$.
4. What are the domain and range of $f^{-1}(x)$? What are the (restricted) domain and range of $f(x)$?






Domain: $(-\infty, \frac{1}{2}]$




Domain: $[\frac{1}{2}, \infty)$




$$f^{-1}(20) =$$

$$\text{Domain: } \left[\frac{1}{2}, \infty\right)$$



$$f^{-1}(20) =$$




$$f^{-1}(20) = 5$$


$$20 = x^2 - x$$

$$0 = x^2 - x - 20 = (x - 5)(x + 4)$$


$$x = 5$$




$$f^{-1}(x) =$$



$$f^{-1}(x) =$$



$$f^{-1}(x) =$$




$$f^{-1}(x) = \frac{1+\sqrt{1+4x}}{2}$$


$$a^2 - a = x, \text{ find } a$$

$$a^2 - a - x = 0$$


$$a = \frac{1 \pm \sqrt{1+4x}}{2}$$

$$f^{-1}(x) = \frac{1+\sqrt{1+4x}}{2}$$







$$f(x) = x^2 - x, \text{ domain: } \left[\frac{1}{2}, \infty\right)$$



$$f(x) = x^2 - x, \text{ domain: } \left[\frac{1}{2}, \infty\right)$$




$$f(x) = x^2 - x, \text{ domain: } \left[\frac{1}{2}, \infty\right)$$



$$f(x) = x^2 - x, \text{ domain: } \left[\frac{1}{2}, \infty\right)$$

$$f^{-1}(x) = \frac{1 + \sqrt{1 + 4x}}{2}$$



INVERTIBILITY GAME: $f(x) = e^x$

- I'm thinking of an x . Your clue: $f(x) = e$. What is my x ?

INVERTIBILITY GAME: $f(x) = e^x$

- ▶ I'm thinking of an x . Your clue: $f(x) = e$. What is my x ?
- ▶ I'm thinking of an x . Your clue: $f(x) = 1$. What is my x ?

INVERTIBILITY GAME: $f(x) = e^x$

- ▶ I'm thinking of an x . Your clue: $f(x) = e$. What is my x ?
- ▶ I'm thinking of an x . Your clue: $f(x) = 1$. What is my x ?
- ▶ I'm thinking of an x . Your clue: $f(x) = \frac{1}{e}$. What is my x ?

INVERTIBILITY GAME: $f(x) = e^x$

- ▶ I'm thinking of an x . Your clue: $f(x) = e$. What is my x ?
- ▶ I'm thinking of an x . Your clue: $f(x) = 1$. What is my x ?
- ▶ I'm thinking of an x . Your clue: $f(x) = \frac{1}{e}$. What is my x ?
- ▶ I'm thinking of an x . Your clue: $f(x) = e^3$. What is my x ?

INVERTIBILITY GAME: $f(x) = e^x$

- ▶ I'm thinking of an x . Your clue: $f(x) = e$. What is my x ?
- ▶ I'm thinking of an x . Your clue: $f(x) = 1$. What is my x ?
- ▶ I'm thinking of an x . Your clue: $f(x) = \frac{1}{e}$. What is my x ?
- ▶ I'm thinking of an x . Your clue: $f(x) = e^3$. What is my x ?
- ▶ I'm thinking of an x . Your clue: $f(x) = 0$. What is my x ?

INVERTIBILITY GAME: $f(x) = e^x$

- ▶ I'm thinking of an x . Your clue: $f(x) = e$. What is my x ? $x = 1$
- ▶ I'm thinking of an x . Your clue: $f(x) = 1$. What is my x ? $x = 0$
- ▶ I'm thinking of an x . Your clue: $f(x) = \frac{1}{e}$. What is my x ? $x = -1$
- ▶ I'm thinking of an x . Your clue: $f(x) = e^3$. What is my x ? $x = 3$
- ▶ I'm thinking of an x . Your clue: $f(x) = 0$. What is my x ? **Trick question: no x gives $f(x) = 0$.**

INVERTIBILITY GAME: $f(x) = e^x$

$$f^{-1}(x) = \log_e x$$

- ▶ I'm thinking of an x . Your clue: $f(x) = e$. What is my x ? $x = 1$
- ▶ I'm thinking of an x . Your clue: $f(x) = 1$. What is my x ? $x = 0$
- ▶ I'm thinking of an x . Your clue: $f(x) = \frac{1}{e}$. What is my x ? $x = -1$
- ▶ I'm thinking of an x . Your clue: $f(x) = e^3$. What is my x ? $x = 3$
- ▶ I'm thinking of an x . Your clue: $f(x) = 0$. What is my x ? **Trick question: no x gives $f(x) = 0$.**

INVERTIBILITY GAME: $f(x) = e^x$

$$f^{-1}(x) = \log_e x$$

- ▶ I'm thinking of an x . Your clue: $f(x) = e$. What is my x ? $x = 1$
 $\log_e(e) = 1$
- ▶ I'm thinking of an x . Your clue: $f(x) = 1$. What is my x ? $x = 0$
 $\log_e(1) = 0$
- ▶ I'm thinking of an x . Your clue: $f(x) = \frac{1}{e}$. What is my x ? $x = -1$
 $\log_e\left(\frac{1}{e}\right) = -1$
- ▶ I'm thinking of an x . Your clue: $f(x) = e^3$. What is my x ? $x = 3$
 $\log_e(e^3) = 3$
- ▶ I'm thinking of an x . Your clue: $f(x) = 0$. What is my x ? **Trick question: no x gives $f(x) = 0$.**
 $\log_e(x)$ is undefined at $x = 0$

1. Suppose $0 < x < 1$. Then $\log_e(x)$ is...
2. Suppose $-1 < x < 0$. Then $\log_e(x)$ is...
3. Suppose $e < x$. Then $\log_e(x)$ is...
 - A. positive
 - B. negative
 - C. greater than one
 - D. less than one
 - E. undefined

EXPONENTS AND LOGARITHMS

$$f(x) = e^x$$

$$f^{-1}(x) = \log_e(x) = \ln(x) = \log(x)$$

EXPONENTS AND LOGARITHMS

$$f(x) = e^x$$

$$f^{-1}(x) = \log_e(x) = \ln(x) = \log(x)$$

x	e^x	
0	1	
1	e	
-1	$\frac{1}{e}$	
n	e^n	

EXPONENTS AND LOGARITHMS

$$f(x) = e^x$$

$$f^{-1}(x) = \log_e(x) = \ln(x) = \text{log}(x)$$

x	e^x	$e \text{ fact} \leftrightarrow \log_e \text{ fact}$	x	$\log_e(x)$
0	1			
1	e			
-1	$\frac{1}{e}$			
n	e^n			

EXPONENTS AND LOGARITHMS

$$f(x) = e^x$$

$$f^{-1}(x) = \log_e(x) = \ln(x) = \text{log}(x)$$

x	e^x	$e \text{ fact} \leftrightarrow \log_e \text{ fact}$	x	$\log_e(x)$
0	1	$e^0 = 1 \leftrightarrow \log_e(1) = 0$	1	0
1	e			
-1	$\frac{1}{e}$			
n	e^n			

EXPONENTS AND LOGARITHMS

$$f(x) = e^x$$

$$f^{-1}(x) = \log_e(x) = \ln(x) = \text{log}(x)$$

x	e^x	$e \text{ fact} \leftrightarrow \log_e \text{ fact}$	x	$\log_e(x)$
0	1	$e^0 = 1 \leftrightarrow \log_e(1) = 0$	1	0
1	e	$e^1 = e \leftrightarrow \log_e(e) = 1$	e	1
-1	$\frac{1}{e}$			
n	e^n			

EXPONENTS AND LOGARITHMS

$$f(x) = e^x$$

$$f^{-1}(x) = \log_e(x) = \ln(x) = \text{log}(x)$$


x	e^x	$e \text{ fact} \leftrightarrow \log_e \text{ fact}$	x	$\log_e(x)$
0	1	$e^0 = 1 \leftrightarrow \log_e(1) = 0$	1	0
1	e	$e^1 = e \leftrightarrow \log_e(e) = 1$	e	1
-1	$\frac{1}{e}$	$e^{-1} = \frac{1}{e} \leftrightarrow \log_e(\frac{1}{e}) = -1$	$\frac{1}{e}$	-1
n	e^n			


EXPONENTS AND LOGARITHMS


$$f(x) = e^x$$


$$f^{-1}(x) = \log_e(x) = \ln(x) = \text{log}(x)$$


x	e^x	$e \text{ fact} \leftrightarrow \log_e \text{ fact}$	x	$\log_e(x)$
0	1	$e^0 = 1 \leftrightarrow \log_e(1) = 0$	1	0
1	e	$e^1 = e \leftrightarrow \log_e(e) = 1$	e	1
-1	$\frac{1}{e}$	$e^{-1} = \frac{1}{e} \leftrightarrow \log_e(\frac{1}{e}) = -1$	$\frac{1}{e}$	-1
n	e^n	$e^n = e^n \leftrightarrow \log_e(e^n) = n$	e^n	n




















LOGS OF OTHER BASES: $\log_n(x)$ IS THE INVERSE OF n^x

$$\log_{10} 10^8 =$$

A. 0

B. 8

C. 10

D. other

LOGS OF OTHER BASES: $\log_n(x)$ IS THE INVERSE OF n^x

$$\log_{10} 10^8 =$$

A. 0

B. 8 ✓

C. 10

D. other

LOGS OF OTHER BASES: $\log_n(x)$ IS THE INVERSE OF n^x

$$\log_{10} 10^8 =$$

A. 0

B. 8 ✓

C. 10

D. other

$$\log_2 16 =$$

A. 1

B. 2

C. 3

D. other

LOGS OF OTHER BASES: $\log_n(x)$ IS THE INVERSE OF n^x

$$\log_{10} 10^8 =$$

A. 0

B. 8 ✓

C. 10

D. other

$$\log_2 16 =$$

A. 1

B. 2

C. 3

D. other ✓ $2^4 = 16$ so $\log_2 16 = 4$

Logarithm Rules

Let A and B be positive, and let n be any real number.

Logarithm Rules

Let A and B be positive, and let n be any real number.

$$\log(A \cdot B) =$$

Logarithm Rules

Let A and B be positive, and let n be any real number.

$$\log(A \cdot B) = \log(A) + \log(B)$$

$$\text{Proof: } \log(A \cdot B) = \log(e^{\log A} e^{\log B}) = \log(e^{\log A + \log B}) = \log(A) + \log(B)$$

Logarithm Rules

Let A and B be positive, and let n be any real number.

$$\log(A \cdot B) = \log(A) + \log(B)$$

Proof: $\log(A \cdot B) = \log(e^{\log A} e^{\log B}) = \log(e^{\log A + \log B}) = \log(A) + \log(B)$

$$\log(A/B) =$$

Logarithm Rules

Let A and B be positive, and let n be any real number.

$$\log(A \cdot B) = \log(A) + \log(B)$$

$$\text{Proof: } \log(A \cdot B) = \log(e^{\log A} e^{\log B}) = \log(e^{\log A + \log B}) = \log(A) + \log(B)$$

$$\log(A/B) = \log(A) - \log(B)$$

$$\text{Proof: } \log(A/B) = \log\left(\frac{e^{\log A}}{e^{\log B}}\right) = \log(e^{\log A - \log B}) = \log A - \log B$$

Logarithm Rules

Let A and B be positive, and let n be any real number.

$$\log(A \cdot B) = \log(A) + \log(B)$$

Proof: $\log(A \cdot B) = \log(e^{\log A} e^{\log B}) = \log(e^{\log A + \log B}) = \log(A) + \log(B)$

$$\log(A/B) = \log(A) - \log(B)$$

Proof: $\log(A/B) = \log\left(\frac{e^{\log A}}{e^{\log B}}\right) = \log(e^{\log A - \log B}) = \log A - \log B$

$$\log(A^n)$$

Logarithm Rules

Let A and B be positive, and let n be any real number.

$$\log(A \cdot B) = \log(A) + \log(B)$$

Proof: $\log(A \cdot B) = \log(e^{\log A} e^{\log B}) = \log(e^{\log A + \log B}) = \log(A) + \log(B)$

$$\log(A/B) = \log(A) - \log(B)$$

Proof: $\log(A/B) = \log\left(\frac{e^{\log A}}{e^{\log B}}\right) = \log(e^{\log A - \log B}) = \log A - \log B$

$$\log(A^n) = n \log(A)$$

Proof: $\log(A^n) = \log\left((e^{\log A})^n\right) = \log(e^{n \log A}) = n \log A$

Logarithm Rules

Let A and B be positive, and let n be any real number.

$$\log(A \cdot B) = \log(A) + \log(B)$$

$$\log(A/B) = \log(A) - \log(B)$$

$$\log(A^n) = n \log(A)$$

Logarithm Rules

Let A and B be positive, and let n be any real number.

$$\log(A \cdot B) = \log(A) + \log(B)$$

$$\log(A/B) = \log(A) - \log(B)$$

$$\log(A^n) = n \log(A)$$

Write as a single logarithm:

$$f(x) = \log\left(\frac{10}{x^2}\right) + 2 \log x + \log(10 + x)$$

Logarithm Rules

Let A and B be positive, and let n be any real number.

$$\log(A \cdot B) = \log(A) + \log(B)$$

$$\log(A/B) = \log(A) - \log(B)$$

$$\log(A^n) = n \log(A)$$

$$\begin{aligned} f(x) &= \log\left(\frac{10}{x^2}\right) + 2 \log x + \log(10 + x) \\ &= \log 10 - \log(x^2) + 2 \log x + \log(10 + x) \\ &= \log 10 - 2 \log x + 2 \log x + \log(10 + x) \\ &= \log 10 + \log(10 + x) = \log(10(10 + x)) \\ &= \log(100 + 10x) \end{aligned}$$

BASE CHANGE

Fact: $b^{\log_b(a)} = a$

BASE CHANGE

$$\text{Fact: } b^{\log_b(a)} = a$$

$$\Rightarrow \log(b^{\log_b(a)}) = \log(a)$$

$$\Rightarrow \log_b(a) \log(b) = \log(a)$$

$$\Rightarrow \log_b(a) = \frac{\log(a)}{\log(b)}$$

BASE CHANGE

Fact: $b^{\log_b(a)} = a$

$$\Rightarrow \log(b^{\log_b(a)}) = \log(a)$$

$$\Rightarrow \log_b(a) \log(b) = \log(a)$$

$$\Rightarrow \log_b(a) = \frac{\log(a)}{\log(b)}$$

In general, for positive a , b , and c :

$$\boxed{\log_b(a) = \frac{\log_c(a)}{\log_c(b)}}$$

In general, for positive a , b , and c :

$$\log_b(a) = \frac{\log_c(a)}{\log_c(b)}$$

Suppose your calculator can only compute logarithms base 10. What would you enter to calculate $\log(17)$?

Suppose your calculator can only compute natural logarithms. What would you enter to calculate $\log_2(57)$?

Suppose your calculator can only compute logarithms base 2. What would you enter to calculate $\log(2)$?


In general, for positive a , b , and c :

$$\log_b(a) = \frac{\log_c(a)}{\log_c(b)}$$

Suppose your calculator can only compute logarithms base 10. What would you enter to calculate $\log(17)$? $\frac{\log_{10}(17)}{\log_{10}(e)}$

Suppose your calculator can only compute natural logarithms. What would you enter to calculate $\log_2(57)$? $\frac{\log(57)}{\log(2)}$

Suppose your calculator can only compute logarithms base 2. What would you enter to calculate $\log(2)$? $\frac{\log_2 2}{\log_2 e} = \frac{1}{\log_2 e}$



Decibels: For a particular measure of the power P of a sound wave, the decibels of that sound is:

$$10 \log_{10}(P)$$

So, every ten decibels corresponds to a sound being ten **times** louder.

Decibels: For a particular measure of the power P of a sound wave, the decibels of that sound is:

$$10 \log_{10}(P)$$

So, every ten decibels corresponds to a sound being ten **times** louder.

A lawnmower emits a 100dB sound. How much sound will two lawnmowers make?

- A. 100 dB
- B. 110 dB
- C. 200 dB
- D. other

Decibels: For a particular measure of the power P of a sound wave, the decibels of that sound is:


$$10 \log_{10}(P)$$

So, every ten decibels corresponds to a sound being ten **times** louder.

A lawnmower emits a 100dB sound. How much sound will two lawnmowers make?

- A. 100 dB
- B. 110 dB
- C. 200 dB
- D. other ✓ more than 100, less than 110

TABLE OF CONTENTS



DIFFERENTIATING THE NATURAL LOGARITHM

Calculate $\frac{d}{dx} \{\log_e x\}$.

DIFFERENTIATING THE NATURAL LOGARITHM

Calculate $\frac{d}{dx} \{\log_e x\}$.

One Weird Trick:

$$x = e^{\log_e x}$$

DIFFERENTIATING THE NATURAL LOGARITHM

Calculate $\frac{d}{dx} \{\log_e x\}$.

One Weird Trick:

$$x = e^{\log_e x}$$

$$\frac{d}{dx} \{x\} = \frac{d}{dx} \{e^{\log_e x}\}$$

$$1 = e^{\log_e x} \cdot \frac{d}{dx} \{\log_e x\} = x \cdot \frac{d}{dx} \{\log_e x\}$$

$$\frac{1}{x} = \frac{d}{dx} \{\log_e x\}$$

DIFFERENTIATING THE NATURAL LOGARITHM

Calculate $\frac{d}{dx}\{\log_e x\}$.

Derivative of Natural Logarithm – Theorem 2.10.1

$$\frac{d}{dx}\{\log_e x\} = \frac{1}{x} \quad (x > 0)$$

DIFFERENTIATING THE NATURAL LOGARITHM

Calculate $\frac{d}{dx} \{\log_e x\}$.

Derivative of Natural Logarithm – Theorem 2.10.1

$$\frac{d}{dx} \{\log_e x\} = \frac{1}{x} \quad (x > 0)$$

$$\frac{d}{dx} \{\log_e |x|\} =$$

DIFFERENTIATING THE NATURAL LOGARITHM

Calculate $\frac{d}{dx} \{\log_e x\}$.

Derivative of Natural Logarithm – Theorem 2.10.1

$$\frac{d}{dx} \{\log_e x\} = \frac{1}{x} \quad (x > 0)$$

$$\frac{d}{dx} \{\log_e |x|\} = \frac{1}{x} \quad (x \neq 0)$$

Derivative of Natural Logarithm

$$\frac{d}{dx} \{\log_e |x|\} = \frac{1}{x} \quad (x \neq 0)$$

Differentiate: $f(x) = \log_e |x^2 + 1|$

Derivative of Natural Logarithm

$$\frac{d}{dx} \{\log_e |x|\} = \frac{1}{x} \quad (x \neq 0)$$

Differentiate: $f(x) = \log_e |x^2 + 1|$

We use the chain rule:

$$\begin{aligned} \frac{d}{dx} \left\{ \log_e \left| \boxed{x^2 + 1} \right| \right\} &= \frac{1}{x^2 + 1} \cdot (2x) \\ &= \frac{2x}{x^2 + 1} \end{aligned}$$

Derivatives of Logarithms – Corollary 2.10.6

For $a > 0$:

$$\frac{d}{dx} [\log_a |x|] = \frac{1}{x \log a}$$

In particular:

$$\frac{d}{dx} [\log |x|] = \frac{1}{x}$$

Derivatives of Logarithms – Corollary 2.10.6

For $a > 0$:

$$\frac{d}{dx} [\log_a |x|] = \frac{1}{x \log a}$$

In particular:

$$\frac{d}{dx} [\log |x|] = \frac{1}{x}$$

Differentiate: $f(x) = \log_e |\cot x|$

Derivatives of Logarithms – Corollary 2.10.6

For $a > 0$:

$$\frac{d}{dx} [\log_a |x|] = \frac{1}{x \log a}$$

In particular:

$$\frac{d}{dx} [\log |x|] = \frac{1}{x}$$

Differentiate: $f(x) = \log_e |\cot x|$

We use the chain rule:

$$\frac{d}{dx} [\log_e |\boxed{\cot x}|] = \frac{1}{\cot x} \cdot (-\csc^2 x) = \frac{-\csc^2 x}{\cot x}$$

LOGARITHMIC DIFFERENTIATION - A FANCY TRICK

$$\blacktriangleright \log(f \cdot g) = \log f + \log g$$

$$\blacktriangleright \log\left(\frac{f}{g}\right) = \log f - \log g$$

$$\blacktriangleright \log(f^g) = g \log f$$

LOGARITHMIC DIFFERENTIATION - A FANCY TRICK

► $\log(f \cdot g) = \log f + \log g$

multiplication turns into addition

► $\log\left(\frac{f}{g}\right) = \log f - \log g$

division turns into subtraction

► $\log(f^g) = g \log f$

exponentiation turns into multiplication

LOGARITHMIC DIFFERENTIATION - A FANCY TRICK

► $\log(f \cdot g) = \log f + \log g$

multiplication turns into addition

► $\log\left(\frac{f}{g}\right) = \log f - \log g$

division turns into subtraction

► $\log(f^g) = g \log f$

exponentiation turns into multiplication


We can exploit these properties to differentiate!

Logarithmic Differentiation

In general, if $f(x) \neq 0$, $\frac{d}{dx} [\log |f(x)|] = \frac{f'(x)}{f(x)}$.

$$f(x) = \left(\frac{(2x+5)^4(x^2+1)}{x+3} \right)^5$$

Find $f'(x)$.



LOGARITHMIC DIFFERENTIATION - A FANCY TRICK

$$f(x) = \left(\frac{(2x+5)^4(x^2+1)}{x+3} \right)^5$$

LOGARITHMIC DIFFERENTIATION - A FANCY TRICK

$$f(x) = \left(\frac{(2x+5)^4(x^2+1)}{x+3} \right)^5$$

$$\log(f(x)) = \log \left[\left(\frac{(2x+5)^4(x^2+1)}{x+3} \right)^5 \right]$$

$$= 5 \log \left[\frac{(2x+5)^4(x^2+1)}{x+3} \right]$$

$$= 5 \left[4 \log(2x+5) + \log(x^2+1) - \log(x+3) \right]$$

$$\frac{f'(x)}{f(x)} = 5 \left[4 \frac{2}{2x+5} + \frac{2x}{x^2+1} - \frac{1}{x+3} \right]$$

$$f'(x) = \left(\frac{(2x+5)^4(x^2+1)}{x+3} \right)^5 \cdot 5 \left[4 \frac{2}{2x+5} + \frac{2x}{x^2+1} - \frac{1}{x+3} \right]$$

LOGARITHMIC DIFFERENTIATION - A FANCY TRICK

Differentiate:

$$f(x) = x^x$$

LOGARITHMIC DIFFERENTIATION - A FANCY TRICK

Differentiate:

$$f(x) = x^x$$

$$\log(f(x)) = \log[x^x]$$

$$= x \log x$$

$$\frac{f'(x)}{f(x)} = x \cdot \frac{1}{x} + \log x \cdot 1$$

$$= 1 + \log x$$

$$f'(x) = x^x [1 + \log x]$$

LOGARITHMIC DIFFERENTIATION - A FANCY TRICK

Differentiate:

$$f(x) = \left(\frac{(x^{15} - 9x^2)^{10}(x + x^2 + 1)}{(x^7 + 7)(x + 1)(x + 2)(x + 3)} \right)^5$$

LOGARITHMIC DIFFERENTIATION - A FANCY TRICK

Differentiate:

$$f(x) = \left(\frac{(x^{15} - 9x^2)^{10}(x + x^2 + 1)}{(x^7 + 7)(x + 1)(x + 2)(x + 3)} \right)^5$$

$$\log(f(x)) = \log \left[\left(\frac{(x^{15} - 9x^2)^{10}(x + x^2 + 1)}{(x^7 + 7)(x + 1)(x + 2)(x + 3)} \right)^5 \right]$$

$$= 5 \log \left[\frac{(x^{15} - 9x^2)^{10}(x + x^2 + 1)}{(x^7 + 7)(x + 1)(x + 2)(x + 3)} \right]$$

$$= 5 \left[10 \log(x^{15} - 9x^2) + \log(x + x^2 + 1) - \log(x^7 + 7) - \log(x + 1) - \log(x + 2) - \log(x + 3) \right]$$

$$\frac{f'(x)}{f(x)} = 5 \left[10 \frac{15x^{14} - 18x}{x^{15} - 9x^2} + \frac{1 + 2x}{x + x^2 + 1} - \frac{7x^6}{x^7 + 7} - \frac{1}{x + 1} - \frac{1}{x + 2} - \frac{1}{x + 3} \right]$$

$$f'(x) = \left(\frac{(x^{15} - 9x^2)^{10}(x + x^2 + 1)}{(x^7 + 7)(x + 1)(x + 2)(x + 3)} \right)^5 \cdot 5 \left[10 \frac{15x^{14} - 18x}{x^{15} - 9x^2} + \frac{1 + 2x}{x + x^2 + 1} - \frac{7x^6}{x^7 + 7} - \frac{1}{x + 1} - \frac{1}{x + 2} - \frac{1}{x + 3} \right]$$

$$f(x) = \frac{(x^8 - e^x)(\sqrt{x} + 5)}{\csc^5 x}$$

$$f(x) = \frac{(x^8 - e^x)(\sqrt{x} + 5)}{\csc^5 x}$$

$$\frac{(x^8 - e^x)(\sqrt{x} + 5)}{\csc^5 x} = f(x)$$

$$\log \left| \frac{(x^8 - e^x)(\sqrt{x} + 5)}{\csc^5 x} \right| = \log |f(x)|$$

$$\left[\log |x^8 - e^x| + \log |x^{1/2} + 5| - 5 \log |\csc x| \right] = \log |f(x)|$$

$$\frac{d}{dx} \left[\log |x^8 - e^x| + \log |x^{1/2} + 5| - 5 \log |\csc x| \right] = \frac{d}{dx} \log |f(x)|$$

$$\frac{8x^7 - e^x}{x^8 - e^x} + \frac{\frac{1}{2}x^{-1/2}}{x^{1/2} + 5} - 5 \frac{-\csc x \cot x}{\csc x} = \frac{f'(x)}{f(x)}$$

$$\frac{(x^8 - e^x)(\sqrt{x} + 5)}{\csc^5 x} \left(\frac{8x^7 - e^x}{x^8 - e^x} + \frac{\frac{1}{2}x^{-1/2}}{x^{1/2} + 5} + 5 \frac{\csc x \cot x}{\csc x} \right) = f'(x)$$

$$f(x) = (x^2 + 17)(32x^5 - 8)(x^{98} - x^{57} + 32x^2)^4(32x^{10} - 10x^{32})$$

Find $f'(x)$.



$$f(x) = (x^2 + 17)(32x^5 - 8)(x^{98} - x^{57} + 32x^2)^4(32x^{10} - 10x^{32})$$

Find $f'(x)$.

$$(x^2 + 17)(32x^5 - 8)(x^{98} - x^{57} + 32x^2)^4(32x^{10} - 10x^{32}) = f(x)$$

$$\log \left| (x^2 + 17)(32x^5 - 8)(x^{98} - x^{57} + 32x^2)^4(32x^{10} - 10x^{32}) \right| = \log |f(x)|$$

$$\log |x^2 + 17| + \log |32x^5 - 8| + 4 \log |x^{98} - x^{57} + 32x^2| + \log |32x^{10} - 10x^{32}| = \log |f(x)|$$

$$\frac{d}{dx} \left[\log |x^2 + 17| + \log |32x^5 - 8| + 4 \log |x^{98} - x^{57} + 32x^2| + \log |32x^{10} - 10x^{32}| \right] = \frac{d}{dx} [\log |f(x)|]$$

$$\frac{2x}{x^2 + 17} + \frac{160x^4}{32x^5 - 8} + 4 \frac{98x^{97} - 57x^{56} + 64x}{x^{98} - x^{57} + 32x^2} + \frac{320x^9 - 320x^{31}}{32x^{10} - 10x^{32}} = \frac{f'(x)}{f(x)}$$

$$\left((x^2 + 17)(32x^5 - 8)(x^{98} - x^{57} + 32x^2)^4(32x^{10} - 10x^{32}) \right) \cdot$$

$$\left(\frac{2x}{x^2 + 17} + \frac{160x^4}{32x^5 - 8} + 4 \frac{98x^{97} - 57x^{56} + 64x}{x^{98} - x^{57} + 32x^2} + \frac{320x^9 - 320x^{31}}{32x^{10} - 10x^{32}} \right) = f'(x)$$



Included Work



Anonymous. (2012) [Decibel Scale of Frequently Heard Sounds](#).
Biology Forums. <http://biology-forums.com/index.php?action=gallery;sa=view;id=6156> (accessed 7 October 2015) , 97



'Slide rule – KEUFFEL & ESSER CO. N.Y.' by s58y is licensed under
[CC BY 2.0](#) (accessed 20 July 2021) , 115



'Notebook' by [Iconic](#) is licensed under [CC BY 3.0](#) (accessed 9 June 2021), 37, 115