
Table of Contents



Implicitly Defined Functions

$$y^2 + x^2 + xy + x^2y = 1$$

Implicitly Defined Functions

$$y^2 + x^2 + xy + x^2y = 1$$

Which of the following points are on the curve?

$(0, 1)$, $(0, -1)$, $(0, 0)$, $(1, 1)$

Q

ans

Implicitly Defined Functions

$$y^2 + x^2 + xy + x^2y = 1$$

Which of the following points are on the curve?

$(0, 1)$, $(0, -1)$, $(0, 0)$, $(1, 1)$

$(0, 1)$ and $(0, -1)$

A

Implicitly Defined Functions

$$y^2 + x^2 + xy + x^2y = 1$$

Which of the following points are on the curve?

$(0,1)$, $(0,-1)$, $(0,0)$, $(1,1)$

$(0,1)$ and $(0,-1)$

If $x = -3$, what is y?

Q

ans

Implicitly Defined Functions

$$y^2 + x^2 + xy + x^2y = 1$$

Which of the following points are on the curve?


$(0,1)$, $(0,-1)$, $(0,0)$, $(1,1)$

$(0,1)$ and $(0,-1)$


If $x = -3$, what is y? $y = -2$ and $y = -4$

A

$$y^2 + x^2 + xy + x^2y = 1$$




$$y^2 + x^2 + xy + x^2y = 1$$



Still has a slope: $\frac{\Delta y}{\Delta x}$

$$y^2 + x^2 + xy + x^2y = 1$$



Still has a slope: $\frac{\Delta y}{\Delta x}$
Locally, y is still a function of x.

$$y^2 + x^2 + xy + x^2y = 1$$

Consider y as a function of x. Can we find $\frac{dy}{dx}$?

$\frac{d}{dx}[y] =$

ans

$$y^2 + x^2 + xy + x^2y = 1$$

Consider y as a function of x. Can we find $\frac{dy}{dx}$?

$$\frac{d}{dx}[y] = \frac{dy}{dx} = y'$$

$$y^2 + x^2 + xy + x^2y = 1$$

Consider y as a function of x. Can we find $\frac{dy}{dx}$?

$\frac{d}{dx}[y] = \frac{dy}{dx} = y'$ $\qquad$ $\frac{d}{dx}[x] =$

$$y^2 + x^2 + xy + x^2y = 1$$

Consider y as a function of x. Can we find $\frac{dy}{dx}$?

$$\frac{d}{dx}[y] = \frac{dy}{dx} = y' \qquad \frac{d}{dx}[x] = 1$$

$$y^2 + x^2 + xy + x^2y = 1$$

Consider y as a function of x. Can we find $\frac{dy}{dx}$?

$\frac{d}{dx}[y] = \frac{dy}{dx} = y'$ $\frac{d}{dx}[x] = 1$ $\frac{d}{dx}[1] =$

ans

$$y^2 + x^2 + xy + x^2y = 1$$

Consider y as a function of x. Can we find $\frac{dy}{dx}$?

$$\frac{d}{dx}[y] = \frac{dy}{dx} = y' \qquad \frac{d}{dx}[x] = 1 \qquad \frac{d}{dx}[1] = 0$$

ans

$$y^2 + x^2 + xy + x^2y = 1$$

Consider y as a function of x. Can we find $\frac{dy}{dx}$?

$$\frac{d}{dx}[y] = \frac{dy}{dx} = y' \qquad \frac{d}{dx}[x] = 1 \qquad \frac{d}{dx}[1] = 0$$

Differentiate both sides with respect to x.

$$0 = 2y\frac{dy}{dx} + 2x + \left(x\frac{dy}{dx} + (1)y\right) + \left(x^2\frac{dy}{dx} + 2xy\right)$$

$$0 = \frac{dy}{dx}\left(2y + x + x^2\right) + (2x + y + 2xy)$$

$$-(2x + y + 2xy) = \frac{dy}{dx}\left(2y + x + x^2\right)$$

$$-\frac{2x + y + 2xy}{2y + x + x^2} = \frac{dy}{dx}$$

$$y^2 + x^2 + xy + x^2y = 1$$


$$\frac{\mathrm{d}y}{\mathrm{d}x} = -\frac{2x + y + 2xy}{2y + x + x^2}$$

ans

$$y^2 + x^2 + xy + x^2y = 1$$

$$\frac{dy}{dx} = -\frac{2x + y + 2xy}{2y + x + x^2}$$

Necessarily, $\frac{dy}{dx}$ depends on **both** y and x. Why?



ans

$$y^2 + x^2 + xy + x^2y = 1$$

$$\frac{dy}{dx} = -\frac{2x + y + 2xy}{2y + x + x^2}$$

Necessarily, $\frac{dy}{dx}$ depends on **both** y and x. Why?

$$\left.\frac{dy}{dx}\right|_{(1,0)} =$$

$$\left.\frac{dy}{dx}\right|_{(1,-2)} =$$

ans

$$y^2 + x^2 + xy + x^2y = 1$$

$$\frac{dy}{dx} = -\frac{2x + y + 2xy}{2y + x + x^2}$$

Necessarily, $\frac{dy}{dx}$ depends on **both** y and x. Why?

$$\left.\frac{dy}{dx}\right|_{(1,0)} = -\frac{2(1) + 0 + 2(1)(0)}{2(0) + 1 + 1}$$

$$= -\frac{2}{2} = -1$$

$$\left.\frac{dy}{dx}\right|_{(1,-2)} = -\frac{2(1) - 2 + 2(1)(-2)}{2(-2) + 1 + 1}$$

$$= -2$$

$$y^2 + x^2 + xy + x^2y = 1$$

$$\frac{dy}{dx} = -\frac{2x + y + 2xy}{2y + x + x^2}$$

Necessarily, $\frac{dy}{dx}$ depends on **both** y and x. Why?


$$\left.\frac{dy}{dx}\right|_{(1,0)} = -\frac{2(1) + 0 + 2(1)(0)}{2(0) + 1 + 1} = -\frac{2}{2} = -1$$

$$\left.\frac{dy}{dx}\right|_{(1,-2)} = -\frac{2(1) - 2 + 2(1)(-2)}{2(-2) + 1 + 1} = -2$$

Points with the same x-value may have different slopes. We need both the x-value and the y-value to figure out which point we're talking about.

$$y^2 + x^2 + xy + x^2y = 1$$

$$\frac{dy}{dx} = -\frac{2x + y + 2xy}{2y + x + x^2}$$



Points with the same x-value may have different slopes. We need both the x-value and the y-value to figure out which point we're talking about.

NOW YOU 

Suppose $x^4y + y^4x = 2$. Find $\frac{dy}{dx}$ at the point $(1, 1)$.

Q

ans

NOW YOU 

Suppose $x^4y + y^4x = 2$. Find $\frac{dy}{dx}$ at the point $(1, 1)$.

$$x^4y(x) + y(x)^4x = 2$$

$$4x^3y(x) + x^4\frac{dy}{dx}(x) + y(x)^4 + 4y(x)^3\frac{dy}{dx}(x)\,x = 0$$

We may only replace variables with constants *after* differentiating. When $x = 1$ and $y(1) = 1$,


$$4(1)^3y(1) + (1)^4\frac{dy}{dx}(1) + y(1)^4 + 4y(1)^3\frac{dy}{dx}(1) = 0$$

$$4 + \frac{dy}{dx}(1) + 1 + 4\frac{dy}{dx}(1) = 0$$

$$5\frac{dy}{dx}(1) = -5$$

$$\frac{dy}{dx}(1) = -1$$

A



$$x^4y + y^4x = 2$$

NOW YOU  Suppose $\dfrac{3y^2+2y+y^3}{x^2+1} = x$. Find $\dfrac{dy}{dx}$ when $x = 0$, and the equations of the associated tangent line(s).

Q

ans

NOW YOU  Suppose $\dfrac{3y^2+2y+y^3}{x^2+1}=x$. Find $\dfrac{\mathrm{d}y}{\mathrm{d}x}$ when $x=0$, and the equations of the associated tangent line(s).

To avoid the quotient rule, we start by simplifying our expression.

$$\frac{3y(x)^2+2y(x)+y(x)^3}{x^2+1}=x$$

$$3y(x)^2+2y(x)+y(x)^3=x^3+x$$

$$6y(x)\frac{\mathrm{d}y}{\mathrm{d}x}(x)+2\frac{\mathrm{d}y}{\mathrm{d}x}(x)+3y(x)^2\frac{\mathrm{d}y}{\mathrm{d}x}(x)=3x^2+1$$

When $x=0$:

$$\frac{\mathrm{d}y}{\mathrm{d}x}(0)=\frac{1}{6y(0)+2+3y(0)^2}$$

A

We need to know y to find $\frac{dy}{dx}$. We want all points where $x = 0$.

$$3y(0)^2 + 2y(0) + y(0)^3 = 0$$
$$y(0)\left(y(0)^2 + 3y(0) + 2\right) = 0$$
$$y(0)\left(y(0) + 1\right)\left(y(0) + 2\right) = 0$$
$$y(0) = 0,\ y(0) = -1,\ y(0) = -2$$

$$\frac{dy}{dx} = \frac{1}{6y + 2 + 3y^2}$$

$(0, 0)$

$$\left.\frac{dy}{dx}\right|_{(0,0)} = \frac{1}{2}$$

$$y - 0 = \frac{1}{2}(x - 0)$$

$$y = \frac{1}{2}x$$

$(0, -1)$

$$\left.\frac{dy}{dx}\right|_{(0,-1)} = \frac{1}{-6+2+3}$$

$$= -1$$

$$y - (-1) = -1(x - 0)$$

$$y = -x - 1$$


$(0, -2)$

$$\left.\frac{dy}{dx}\right|_{(0,-2)} = \frac{1}{-12+2+12}$$

$$= \frac{1}{2}$$


$$y - (-2) = \frac{1}{2}(x - 0)$$

$$y = \frac{1}{2}x - 2$$



$$\frac{3y^2 + 2y + y^3}{x^2 + 1} = x$$

A



$$\frac{3y^2 + 2y + y^3}{x^2 + 1} = x$$

A

Use implicit differentiation to differentiate $\log(x)$, $x > 0$.

ans

Use implicit differentiation to differentiate $\log(x)$, $x > 0$.

$$\log x = y(x)$$
$$x = e^{y(x)}$$

ans

Use implicit differentiation to differentiate $\log(x)$, $x > 0$.

$$\log x = y(x)$$

$$x = e^{y(x)}$$

$$1 = e^{y(x)} \cdot \frac{\mathrm{d}y}{\mathrm{d}x}(x)$$

$$\frac{\mathrm{d}y}{\mathrm{d}x}(x) = \frac{1}{e^{y(x)}} = \frac{1}{x}$$

Use implicit differentiation to differentiate $\log(x)$, $x > 0$.

$$\log x = y(x)$$
$$x = e^{y(x)}$$
$$1 = e^{y(x)} \cdot \frac{dy}{dx}(x)$$
$$\frac{dy}{dx}(x) = \frac{1}{e^{y(x)}} = \frac{1}{x}$$

Use implicit differentiation to differentiate $\log|x|$, $x < 0$.

Use implicit differentiation to differentiate $\log(x)$, $x > 0$.

Use implicit differentiation to differentiate $\log|x|$, $x < 0$.

Use implicit differentiation to differentiate $\log(x)$, $x > 0$.

Use implicit differentiation to differentiate $\log|x|$, $x < 0$.

$$\log|x| = y(x)$$
$$\log(-x) = y(x)$$
$$-x = e^{y(x)}$$
$$-1 = e^{y(x)} \cdot \frac{dy}{dx}(x)$$
$$\frac{dy}{dx}(x) = \frac{-1}{e^{y(x)}} = \frac{-1}{-x} = \frac{1}{x}$$

Use implicit differentiation to differentiate $\log_a(x)$, where $a > 0$ is a constant and $x > 0$.

Use implicit differentiation to differentiate $\log_a |x|$, $a > 0$.

Use implicit differentiation to differentiate $\log_a(x)$, where $a > 0$ is a constant and $x > 0$.

Use implicit differentiation to differentiate $\log_a(x)$, where $a > 0$ is a constant and $x > 0$.

$$\begin{aligned}
\log_a x &= y(x) \\
x &= a^{y(x)} \\
1 &= a^{y(x)} \cdot \log_e a \cdot \frac{\mathrm{d}y}{\mathrm{d}x}(x) \\
\frac{\mathrm{d}y}{\mathrm{d}x}(x) &= \frac{1}{a^{y(x)} \cdot \log_e a} = \frac{1}{x \log_e a}
\end{aligned}$$

Use implicit differentiation to differentiate $\log_a |x|$, $a > 0$.


Use implicit differentiation to differentiate $\log_a |x|$, $a > 0$.


If $x > 0$, it's what we just computed. So assume $x < 0$.


$$\begin{aligned}
\log_a |x| &= y(x) \\
\log_a (-x) &= y(x) \\
-x &= a^{y(x)} \\
-1 &= a^{y(x)} \cdot \log_e a \cdot \frac{dy}{dx}(x) \\
\frac{dy}{dx}(x) &= \frac{-1}{a^{y(x)} \cdot \log_e a} = \frac{-1}{x \log_e a}
\end{aligned}$$


Included Work

 ‘Brain’ by Eucalyp is licensed under CC BY 3.0 (accessed 8 June 2021), 23, 24, 26, 27

 screenshot of graph using Desmos Graphing Calculator, https://www.desmos.com/calculator (accessed 19 October 2017), 18, 22

 screenshot of graph using Desmos Graphing Calculator, https://www.desmos.com/calculator (accessed 19 October 2017), 7–9

 screenshot of graph using Desmos Graphing Calculator, https://www.desmos.com/calculator (accessed 24 October 2018), 25

 screenshot of graph using Desmos Graphing Calculator, https://www.desmos.com/calculator (accessed 24 October 2018), 30

 screenshot of graph using Desmos Graphing Calculator, https://www.desmos.com/calculator (accessed 24 October 2018), 31

 ‘Notebook’ by Iconic is licensed under CC BY 3.0 (accessed 9 June 2021), 35, 38