
Table of Contents



Implicitly Defined Functions


$$y^2 + x^2 + xy + x^2y = 1$$

Which of the following points are on the curve?

$(0, 1)$, $(0, -1)$, $(0, 0)$, $(1, 1)$

If $x = -3$, what is y?

$$y^2 + x^2 + xy + x^2y = 1$$



Still has a slope: $\frac{\Delta y}{\Delta x}$
Locally, y is still a function of x.

$$y^2 + x^2 + xy + x^2y = 1$$


Consider y as a function of x. Can we find $\frac{dy}{dx}$?

$\frac{d}{dx}[y] =$ $\frac{d}{dx}[x] =$ $\frac{d}{dx}[1] =$

$$y^2 + x^2 + xy + x^2y = 1$$

$$\frac{dy}{dx} = -\frac{2x + y + 2xy}{2y + x + x^2}$$

Necessarily, $\frac{dy}{dx}$ depends on **both** y and x. Why?



NOW YOU 

Suppose $x^4y + y^4x = 2$. Find $\frac{dy}{dx}$ at the point $(1, 1)$.

NOW YOU  Suppose $\dfrac{3y^2+2y+y^3}{x^2+1}=x$. Find $\dfrac{\mathrm{d}y}{\mathrm{d}x}$ when $x=0$, and the equations of the associated tangent line(s).

Use implicit differentiation to differentiate $\log(x)$, $x > 0$.

$$\log x = y(x)$$
$$x = e^{y(x)}$$


Use implicit differentiation to differentiate $\log|x|$, $x < 0$.


Use implicit differentiation to differentiate $\log_a(x)$, where $a > 0$ is a constant and $x > 0$.

Use implicit differentiation to differentiate $\log_a |x|$, $a > 0$.

Included Work

 'Brain' by Eucalyp is licensed under CC BY 3.0 (accessed 8 June 2021), 6, 7

 screenshot of graph using Desmos Graphing Calculator, https://www.desmos.com/calculator (accessed 19 October 2017), 5

 screenshot of graph using Desmos Graphing Calculator, https://www.desmos.com/calculator (accessed 19 October 2017), 3